



HAL
open science

Boundary Conditions for Translation- Invariant Gibbs Measures of the Potts Model on Cayley Trees

Daniel Gandolfo, M. M. Rakhmatullaev, U.A. Rozikov

► **To cite this version:**

Daniel Gandolfo, M. M. Rakhmatullaev, U.A. Rozikov. Boundary Conditions for Translation- Invariant Gibbs Measures of the Potts Model on Cayley Trees. *Journal of Statistical Physics*, 2017, 167 (5), pp.1164. 10.1007/s10955-017-1771-5 . hal-01591298

HAL Id: hal-01591298

<https://hal.science/hal-01591298>

Submitted on 26 Apr 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Boundary conditions for translation-invariant Gibbs measures of the Potts model on Cayley trees

D. Gandolfo

Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France.
gandolfo@cpt.univ-mrs.fr

M. M. Rahmatullaev and U. A. Rozikov

Institute of mathematics, 29, Do'rmon Yo'li str., 100125, Tashkent, Uzbekistan.
mrahmatullaev@rambler.ru rozikovu@yandex.ru

Abstract

We consider translation-invariant splitting Gibbs measures (TISGMs) for the q -state Potts model on a Cayley tree of order two. Recently a full description of the TISGMs was obtained, and it was shown in particular that at sufficiently low temperatures their number is $2^q - 1$. In this paper for each TISGM μ we explicitly give the set of boundary conditions such that limiting Gibbs measures with respect to these boundary conditions coincide with μ .

Mathematics Subject Classifications (2010). 82B26; 60K35.

Key words. Cayley tree, Potts model, boundary condition, Gibbs measure.

1 Introduction.

The analysis of translational invariant splitting Gibbs measures of the q -state Potts model on Cayley trees is based on the classification of translation-invariant boundary laws which are in one-to-one correspondence with the TISGMs. Recall that boundary laws are length- q vectors which satisfy a non-linear fixed-point equation (tree recursion).

It has been known for a long time that for the anti-ferromagnetic Potts model there exists a unique TISGM [11] and for the ferromagnetic Potts model at sufficiently low temperatures there are at least $q + 1$ translation-invariant Gibbs measures [4], [5].

One of the $q + 1$ well-known measures mentioned above is obtained as infinite-volume limit of the finite-dimensional Gibbs measures with free boundary condition and each of the remaining q measures is obtained as the corresponding limit with the boundary conditions of homogeneous (constant) spin-configurations. While the q measures with homogeneous boundary conditions are always extremal in the set of all Gibbs measures [4], [5], for the free boundary condition measure there is a temperature, denoted by T_0 , which is below the transition temperature, such that the measure is an extremal Gibbs measure if $T \geq T_0$ and loses its extremality for even lower temperatures [10, Theorem 5.6].

Recently, in [7] all TISGMs for the Potts model were found on the Cayley tree of order $k \geq 2$, and it is shown that at sufficiently low temperatures their number is $2^q - 1$. We

note that the number of TISGMs does not depend on $k \geq 2$. In the case $k = 2$ the explicit formulae for the critical temperatures and all TISGMs are given.

In [8] some regions for the temperature ensuring that a given TISGM is (non-)extreme in the set of all Gibbs measures are found. In particular the existence of a temperature interval is shown for which there are at least $2^{q-1} + q$ extremal TISGMs.

The fact that these measures can never be nontrivial convex combinations of each other (i.e., they are extremal in the set of all TIGMs) is almost automatic (see [7, Theorem 2]). However it is not clear what kind of boundary conditions are needed to get the remaining $2^q - q - 2$ TISGMs as corresponding limits with the boundary conditions. In this paper we shall answer this question. It is non-trivial problem since the number of TISGMs (i.e., $2^q - 1$) is larger than the number (i.e., q) of translation-invariant configurations. Therefore one expects to need non-translation-invariant boundary conditions for some TISGMs. Concerning the Ising model, the dependence of TISGMs on boundary conditions has been studied in [9].

The paper is organized as follows. Section 2 contains preliminaries (necessary definitions and facts). In section 3 we will show how to connect boundary laws with boundary conditions, moreover we shall give the list of known TISGMs. Section 4 contains our main result, namely given any TISGM μ , we show how to compute explicitly a set of boundary conditions such that the limiting Gibbs measures with respect to these boundary conditions coincide with μ . In the last section we construct concrete boundary conditions.

2 Definitions

Let $\Gamma^k = (V, L)$ be the regular Cayley tree, where each vertex has $k + 1$ neighbors with V being the set of vertices and L the set of edges.

Two vertices $t, s \in V$, ($t \neq s$) are called *neighbors* if they are connected by an edge. In this case we write $\langle s, t \rangle$. Each vertex of Γ^k has $k + 1$ neighbors.

Fix an origin 0 of Γ^k . We write $s \rightarrow t$, if $t \neq s$ and the path connecting 0 and t passes through s . If $s \rightarrow t$ and s, t are neighbors, then t is called a *direct successor* of s and this we write as $s \rightarrow_1 t$.

For any finite $A \subset V$, the boundary ∂A of A is

$$\partial A = \{t \in V \setminus A : \exists x \in A, \langle x, t \rangle\}.$$

For every $A \subset V$, let $\Omega_A = \{1, 2, \dots, q\}^A$ be the set of all possible spin configurations on A . For brevity we write Ω instead of Ω_V .

For every $A \subset V$ we define the σ -algebra B_A by

$$B_A = \text{the } \sigma \text{- algebra generated by } \{X_t, t \in A\},$$

where $X_t(\sigma) = \sigma(t)$ for all $t \in A, \sigma \in \Omega$. For brevity we write B instead of B_V .

Let A be a finite subset of V , $\omega \in \Omega$ and $\sigma \in \Omega_A$. We define Potts interaction energy on A given the inner configuration σ and the boundary condition ω by

$$E_A^\omega(\sigma) = -J \sum_{\substack{t, s \in A: \\ \langle t, s \rangle}} \delta_{\sigma(t)\sigma(s)} - J \sum_{\substack{t \in A, s \in \partial A: \\ \langle t, s \rangle}} \delta_{\sigma(t)\omega(s)}, \quad (2.1)$$

where $J \in \mathbb{R}$ and δ is the Kronecker's delta.

A finite Gibbs measure P_A^ω on Ω_A corresponding to E_A^ω is defined by

$$P_A^\omega(\sigma) = [Z_A^\omega]^{-1} \exp[-E_A^\omega(\sigma)], \sigma \in \Omega_A, \quad (2.2)$$

where $Z_A^\omega = \sum_{\hat{\sigma} \in \Omega_A} \exp[-E_A^\omega(\hat{\sigma})]$. As usual P_A^ω can be considered as a probability measure on (Ω, B) .

For fixed J , if there is an increasing sequence of finite subsets $\{V_n\}$ such that $V_n \nearrow V$ as $n \rightarrow \infty$ and $P^\omega = w - \lim_{n \rightarrow \infty} P_{V_n}^\omega$ (the weak convergence of measures) exists for suitable fixed $\omega \in \Omega$, then P^ω is called a *limiting Gibbs measure* with boundary condition ω for J . On the other hand, a Gibbs measure P for J is defined as a probability measure on (Ω, B) such that for every M in B_A

$$P(M|B_{A^c})(\omega) = P_A^\omega(M). \quad a.s.(P) \quad (2.3)$$

It is known ([6], [9]) that the set $\mathfrak{S}(J)$ of all Gibbs measures for a fixed J is a non-empty, compact convex set. A limiting Gibbs measure is a Gibbs measure for the same J . Conversely, every extremal point of $\mathfrak{S}(J)$ is a limiting Gibbs measure with a suitable boundary condition for the same J . It is known (see page 241 of [6]) that any extreme Gibbs measure of a Hamiltonian with nearest-neighbor interactions is a *splitting* Gibbs measure (which is equivalently called a tree-indexed Markov chain [6]). Consequently, any non-splitting Gibbs measure is not extreme. However, any splitting Gibbs measure (not necessary extreme) is a limiting Gibbs measure, because it corresponds to a (generalized)¹ boundary condition satisfying a compatibility (tree recursion) condition of Kolmogorov's theorem. In [1] it was shown that for non-extremal Gibbs measures on \mathbb{Z}^d a Gibbs measure need not be a limiting Gibbs measure (see [1] and [2] for more details).

3 Translation-invariant limiting Gibbs measures

Let $|t|$ denote the distance between 0 and $t \in V$, i.e. $|t| = n$ if there exists a chain $0 \rightarrow_1 u_1 \rightarrow_1 u_2 \rightarrow_1 u_3 \rightarrow_1 \dots \rightarrow_1 u_{n-1} \rightarrow_1 t$. We only consider the sequence of boxes

$$V_n = \{t \in V : |t| \leq n\}, n \geq 1.$$

For every $s \in V$ we define

$$\Gamma_s^k = \{s\} \cup \{t \in V : s \rightarrow t\}, \quad \text{and} \quad V_{n,s} = \Gamma_s^k \cap V_n, \quad n \geq 1.$$

Elements of the set $V_1 \setminus \{0\}$ are the nearest neighbors of the origin 0, since this set contains $k+1$ elements, we number them by $1, 2, \dots, k+1$. We note that the subtrees $V_{n,i}$, $i = 1, 2, \dots, k+1$ are similar to each other, i.e., for any $i, j \in \{1, \dots, k+1\}$ the subtree $V_{n,i}$ can be obtained from the $V_{n,j}$ by a rotation around 0. Moreover we have

$$V_n = \{0\} \cup \bigcup_{i=1}^{k+1} V_{n,i}, \quad \text{and} \quad V_{n,i} = \{i\} \cup \bigcup_{j:i \rightarrow_1 j} V_{n-1,j}. \quad (3.1)$$

Using (3.1), from (2.1) we get

$$E_{V_n}^\omega(\sigma) = \sum_{i=1}^{k+1} \left(E_{V_{n,i}}^\omega(\sigma) - J\delta_{\sigma(0)\sigma(i)} \right). \quad (3.2)$$

¹Adding a boundary field at each site of the boundary is called a generalized boundary condition [3] or boundary law [6]

For every $\omega \in \Omega$, $s \in V \setminus \{0\}$ and $n \geq |s|$ define

$$W_{n,s}^\omega(l) = \sum_{\sigma \in \Omega_{V_n, s} : \sigma(s)=l} \exp[-E_{V_n, s}^\omega(\sigma) - J\delta_{l\omega(t)}], \quad l = 1, 2, \dots, q, \quad (3.3)$$

$$R_{n,s}^l(\omega) = \frac{W_{n,s}^\omega(l)}{W_{n,s}^\omega(q)}, \quad l = 1, 2, \dots, q, \quad (3.4)$$

here t is the unique vertex such that $t \rightarrow_1 s$.

Now by (2.2) and (3.2)-(3.4) we get

$$\begin{aligned} \frac{P_{V_n}^\omega(\sigma(0) = l)}{P_{V_n}^\omega(\sigma(0) = q)} &= \frac{\sum_{\sigma: \sigma(0)=l} \exp(-E_{V_n}^\omega(\sigma))}{\sum_{\sigma: \sigma(0)=q} \exp(-E_{V_n}^\omega(\sigma))} \\ &= \prod_{i=1}^{k+1} \frac{(\exp(J) - 1)R_{n,i}^l(\omega) + \sum_{p=1}^{q-1} R_{n,i}^p(\omega) + 1}{\exp(J) + \sum_{p=1}^{q-1} R_{n,i}^p(\omega)}, \quad l = 1, 2, \dots, q, \end{aligned} \quad (3.5)$$

By (3.1) and (3.3) we obtain

$$W_{n,s}^\omega(l) = \prod_{u: s \rightarrow_1 u} [(\exp(J) - 1)W_{n,u}^\omega(l) + \sum_{p=1}^q W_{n,u}^\omega(p)], \quad l = 1, 2, \dots, q, \quad (3.6)$$

and for $n > m$, $\eta \in \Omega_{V_m}$, we get

$$P_{V_n}^\omega(\{\sigma(s) = \eta(s), s \in V_m\}) = \frac{\exp[-E_{V_{m-1}}^\eta(\eta)] \prod_{s \in \partial V_{m-1}} W_{n,s}^\omega(\eta(s))}{\sum_{\xi \in \Omega_{V_m}} \exp[-E_{V_{m-1}}^\xi(\xi)] \prod_{s \in \partial V_{m-1}} W_{n,s}^\omega(\xi(s))}. \quad (3.7)$$

From the above equalities we obtain the following

Lemma 1. *Let $\omega \in \Omega$ be given. If there is $N > 0$ such that $R_{n,s}^l(\omega)$ converges as $n \rightarrow \infty$ for every $s \in V \setminus V_N$ and for every $l = 1, \dots, q$, then $P^\omega = w - \lim_{n \rightarrow \infty} P_{V_n}^\omega$ exists.*

For $n \geq 1$, $p = 1, 2, \dots, q$, $i = 1, \dots, k+1$ we denote

$$A_n = \{t \in V : |t| = n\}, \quad N_{n,i}^{(p)}(\sigma) = |\{x \in A_n \cap V_{n,i} : \sigma(x) = p\}|.$$

Lemma 2. *Let $l = 1, \dots, q$ and ω be a configuration such that²*

$$c^l(\omega) = \sum_{s: t \rightarrow_1 s} \delta_{l\omega(s)}$$

is independent of $t \in V \setminus \{0\}$. Then $R_{n,i}^l(\omega) = R_{n,j}^l(\omega)$ for any $i, j = 1, 2, \dots, k+1$.

Proof. Since $R_{n,i}^l(\omega) = \frac{W_{n,i}^\omega(l)}{W_{n,i}^\omega(q)}$, it suffices to prove that $W_{n,i}^\omega(l) = W_{n,j}^\omega(l)$ for any $i, j = 1, 2, \dots, k+1$.

For the Hamiltonian we have

$$E_{V_n, i}^\omega(\sigma) = E_{V_{n-1}, i}^\sigma(\sigma) - J \sum_{x \in A_n \cap V_{n,i}} \sum_{x \rightarrow_1 y} \delta_{\sigma(x)\omega(y)}. \quad (3.8)$$

²The sum in the RHS of $c^l(\omega)$ is taken over all *direct successors* of t , it should not be confused with the sum over all neighbors of t , i.e., $\sum_{s: (t,s)}$.

By the condition of lemma 2 we obtain

$$E_{V_{n,i}}^\omega(\sigma) = E_{V_{n-1,i}}^\sigma(\sigma) - J \sum_{x \in A_n \cap V_{n,i}} c^{\sigma(x)}(\omega) = E_{V_{n-1,i}}^\sigma(\sigma) - J \sum_{p=1}^q N_{n,i}^{(p)}(\sigma) c^p(\omega).$$

For $W_{n,i}^\omega(l)$ we have

$$W_{n,i}^\omega(l) = \sum_{\sigma \in V_{n,i} : \sigma(i)=l} \exp[E_{V_{n-1,i}}^\sigma(\sigma) - J \sum_{p=1}^q N_{n,i}^{(p)}(\sigma) c^p(\omega) - J \delta_{l\omega(t)}].$$

Since $V_{n,i}$ is similar to $V_{n,j}$, for any $i, j \in \{1, \dots, k+1\}$, $n \geq 1$, there is an one-to-one correspondence γ between sets $\Omega_{V_{n,i}}$ and $\Omega_{V_{n,j}}$, which can be obtained by a rotation of the $V_{n,i}$ on the set $V_{n,j}$. We note that the Potts interaction energy (2.1) is translation-invariant and by the condition of the lemma the quantity $c^l(\omega)$ also does not depend on vertices of the tree. Therefore, if $\gamma(\sigma) = \varphi$ then

$$E_{V_{n-1,i}}^\sigma(\sigma) = E_{V_{n-1,j}}^{\gamma(\sigma)}(\gamma(\sigma)) = E_{V_{n-1,j}}^\varphi(\varphi),$$

$$N_{n,i}^{(p)}(\sigma) = N_{n,j}^{(p)}(\varphi), \quad \forall i, j = 1, \dots, k+1, p = 1, \dots, q.$$

Using these equalities we get

$$\begin{aligned} W_{n,i}^\omega(l) &= \sum_{\sigma \in \Omega_{V_{n,i}} : \sigma(i)=l} \exp[E_{V_{n-1,i}}^\sigma(\sigma) - J \sum_{p=1}^q N_{n,i}^{(p)}(\sigma) c^p(\omega) - J \delta_{l\omega(t)}] = \\ &= \sum_{\varphi \in \Omega_{V_{n,j}} : \varphi(j)=l} \exp[E_{V_{n-1,j}}^\varphi(\varphi) - J \sum_{p=1}^q N_{n,j}^{(p)}(\varphi) c^p(\omega) - J \delta_{l\omega(t)}] = W_{n,j}^\omega(l). \end{aligned}$$

Thus $R_{n,i}^l(\omega) = R_{n,j}^l(\omega)$ for any $i, j = 1, 2, \dots, k+1$. □

From the above proof it follows that $R_{n,s}^l(\omega)$ depends only on $n - |s|$, i.e., we have

$$R_{n,s}^l(\omega) = R_{n-|s|+1}^l(\omega), \quad l = 1, 2, \dots, q. \quad (3.9)$$

Then from (3.6) we get the following

$$Y_n^l(\omega) = k F_l(Y_{n-1}^1(\omega), Y_{n-1}^2(\omega), \dots, Y_{n-1}^{q-1}(\omega)), \quad (3.10)$$

where $l = 1, \dots, q-1$, $n \geq 2$, $Y_n^l(\omega) = \ln R_n^l(\omega)$ and $F = (F_1, \dots, F_{q-1})$ with coordinates

$$F_l(x^1, x^2, \dots, x^{q-1}) = \ln \frac{(\exp(J) - 1) \exp(x^l) + \sum_{p=1}^{q-1} \exp(x^p) + 1}{\exp(J) + \sum_{p=1}^{q-1} \exp(x^p)}. \quad (3.11)$$

It is clear that if $Y^i(\omega)$ is a limit point of $Y_n^i(\omega)$, as $n \rightarrow \infty$ then by (3.10) we get

$$Y^l(\omega) = k F_l(Y^1(\omega), Y^2(\omega), \dots, Y^{q-1}(\omega)), \quad l = 1, 2, \dots, q-1. \quad (3.12)$$

For convenience we denote

$$\theta = \exp(J), \quad h_l = Y^l(\omega), \quad l = 1, 2, \dots, q-1.$$

Then the system (3.12) becomes

$$h_i = k \ln \left(\frac{(\theta - 1)e^{h_i} + \sum_{j=1}^{q-1} e^{h_j} + 1}{\theta + \sum_{j=1}^{q-1} e^{h_j}} \right), \quad i = 1, \dots, q-1. \quad (3.13)$$

In [7] it is proven that to each solution of (3.13) corresponds a unique Gibbs measure which is called a translation-invariant splitting Gibbs measure (TISGM).

In [7] all solutions of the equation (3.13) are given. By these solutions the full set of TISGMs is described. In particular, it is shown that any TISGM of the Potts model corresponds to a solution of the following equation

$$h = f_m(h) \equiv k \ln \left(\frac{(\theta + m - 1)e^h + q - m}{me^h + q - m - 1 + \theta} \right), \quad (3.14)$$

for some $m = 1, \dots, q-1$.

Denote

$$\theta_m = 1 + 2\sqrt{m(q-m)}, \quad m = 1, \dots, q-1. \quad (3.15)$$

It is easy to see that

$$\theta_m = \theta_{q-m} \quad \text{and} \quad \theta_1 < \theta_2 < \dots < \theta_{\lfloor \frac{q}{2} \rfloor - 1} < \theta_{\lfloor \frac{q}{2} \rfloor} \leq q+1. \quad (3.16)$$

Proposition 1. [7] *Let $k = 2$, $J > 0$.*

1. *If $\theta < \theta_1$ then there exists a unique TISGM;*
2. *If $\theta_m < \theta < \theta_{m+1}$ for some $m = 1, \dots, \lfloor \frac{q}{2} \rfloor - 1$ then there are $1 + 2 \sum_{s=1}^m \binom{q}{s}$ TISGMs which correspond to the solutions $h_i \equiv h_i(\theta, s) = 2 \ln[x_i(s, \theta)]$, $i = 1, 2$ $s = 1, \dots, m$ of (3.14), where*

$$\begin{aligned} x_1(s, \theta) &= \frac{\theta - 1 - \sqrt{(\theta - 1)^2 - 4s(q-s)}}{2s}, \\ x_2(s, \theta) &= \frac{\theta - 1 + \sqrt{(\theta - 1)^2 - 4s(q-s)}}{2s}. \end{aligned} \quad (3.17)$$

3. *If $\theta_{\lfloor \frac{q}{2} \rfloor} < \theta \neq q+1$ then there are $2^q - 1$ TISGMs;*
4. *If $\theta = q+1$ the number of TISGMs is as follows*

$$\begin{cases} 2^{q-1}, & \text{if } q \text{ is odd} \\ 2^{q-1} - \binom{q-1}{q/2}, & \text{if } q \text{ is even;} \end{cases}$$

5. *If $\theta = \theta_m$, $m = 1, \dots, \lfloor \frac{q}{2} \rfloor$, $(\theta_{\lfloor \frac{q}{2} \rfloor} \neq q+1)$ then the number of TISGMs is*

$$1 + \binom{q}{m} + 2 \sum_{s=1}^{m-1} \binom{q}{s}.$$

The number of TISGMs does not depend on $k \geq 2$ see [7, Theorem 1]. But for $k \geq 3$ explicit formulas for the solutions are not known. Therefore in this paper we consider only the case $k = 2$.

Following [7] we note that each TISGM corresponds to a solution of (3.14) with some $m \leq \lfloor \frac{q}{2} \rfloor$. Moreover, for a given $m \leq \lfloor \frac{q}{2} \rfloor$, a fixed solution $h_i(\theta, m)$ to (3.14) generates $\binom{q}{m}$

vectors by permuting coordinates of the vector $(\underbrace{h_i, h_i, \dots, h_i}_m, \underbrace{0, 0, \dots, 0}_{q-m})$ and giving $\binom{q}{m}$

TISGMs. Thus without loss of generality we can only consider the measure $\mu_i(\theta, m)$ corresponding to vector $\mathbf{h}(m, i) = (\underbrace{h_i, h_i, \dots, h_i}_m, \underbrace{0, 0, \dots, 0}_{q-m-1})$, i.e., normalized on q th coordinate

(see Remark 2 and Corollary 2 of [7]). Denote by $\mu_0 \equiv \mu_0(\theta)$ the TISGM corresponding to solution $h_i \equiv 0$ and by $\mu_i \equiv \mu_i(\theta, m)$ the TISGM corresponding to the solution $h_i(\theta, m)$, $i = 1, 2$, $m = 1, \dots, \lfloor \frac{q}{2} \rfloor$ (given in Proposition 1). In this paper our aim is to obtain measures μ_i by changing boundary conditions.

4 Boundary conditions for TISGMs

The following lemma can be proved by simple analysis.

Lemma 3. *i. For $k \geq 2$ and $\theta > 1$ the function $f_m(h)$, $h \in \mathbb{R}$ defined in (3.14) has the following properties:*

- a) $\{h : f_m(h) = h\} = \{0, h_1, h_2\}$, if $\theta > \theta_m, m \leq \lfloor q/2 \rfloor$;
- b) $a < f_m(h) < A$, with $a = k \ln \frac{q-m}{q+\theta-m-1}$, $A = k \ln \frac{\theta+m-1}{m}$;
- c) $\frac{d}{dh} f_m(h) = \frac{k(\theta-1)(\theta+q-1)e^h}{(me^h+\theta+q-m-1)((\theta+m-1)e^h+q-m)} > 0$;

ii. If $k = 2$ and $m \leq q/2$ then for solutions h_1 and h_2 mentioned in Proposition 1 the following statements hold

$$\begin{aligned} 0 < h_1 = h_2, & \quad \text{if } \theta = \theta_m \\ 0 < h_1 < h_2, & \quad \text{if } \theta_m < \theta < \theta_c, \quad \text{with } \theta_c = q + 1 \\ 0 = h_1 < h_2, & \quad \text{if } \theta = q + 1 \\ h_1 < 0 < h_2, & \quad \text{if } q + 1 < \theta. \end{aligned}$$

For each solution $h_i(\theta, m)$ we want to find $\omega = \omega(h_i) \in \Omega$, such that $\mu_i(\theta, m) = P^\omega$, where P^ω is defined in Lemma 1.

Consider the dynamical system (3.10) for $k = 2$. Denote $G(h) = 2F(h)$. For a given initial vector $v^{(0)} = (v_1^{(0)}, \dots, v_{q-1}^{(0)})$, we shall study the limit

$$\lim_{n \rightarrow \infty} G^{(n)}(v^{(0)}), \tag{4.1}$$

here $G^{(n)}(v) = \underbrace{G(G(\dots G(v)\dots))}_n$.

Figures 1-4 show the streamlines of the vector field $G^{(n)}(v)$ for $k = 2$, $q = 3$. These figures also illustrate the limit points of (4.1).

Denote

$$I_m = \{v \in \mathbb{R}^{q-1} : v_1 = \dots = v_m, \quad v_{m+1} = \dots = v_{q-1} = 0\}.$$

It is easy to see that the set I_m is invariant with respect to G , i.e. $G(I_m) \subset I_m$.

The following lemma gives the limits of (4.1) on the invariant I_m (compare with Fig.1-4).

Lemma 4. 1) If $\theta = \theta_m$, for some $m = 1, \dots, \lfloor \frac{q}{2} \rfloor$ then

$$\lim_{n \rightarrow \infty} G^{(n)}(v^{(0)}) = \begin{cases} \mathbf{h}(m, 1), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} \geq h_1 \\ (0, \dots, 0), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} < h_1 \end{cases} \quad (4.2)$$

2) If $\theta_m < \theta < \theta_c = q + 1$ then

$$\lim_{n \rightarrow \infty} G^{(n)}(v^{(0)}) = \begin{cases} \mathbf{h}(m, 2), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} > h_1 \\ \mathbf{h}(m, 1), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} = h_1 \\ (0, \dots, 0), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} < h_1 \end{cases} \quad (4.3)$$

3) If $\theta = \theta_c$ then

$$\lim_{n \rightarrow \infty} G^{(n)}(v^{(0)}) = \begin{cases} \mathbf{h}(m, 2), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} > 0 \\ (0, \dots, 0), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} \leq 0 \end{cases} \quad (4.4)$$

4) If $\theta > \theta_c$ then

$$\lim_{n \rightarrow \infty} G^{(n)}(v^{(0)}) = \begin{cases} \mathbf{h}(m, 2), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} > 0 \\ \mathbf{h}(m, 1), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} < 0 \\ (0, \dots, 0), & \text{if } v^{(0)} \in I_m \text{ and } v_1^{(0)} = 0 \end{cases} \quad (4.5)$$

Proof. Restrict function $G(h)$ to I_m , then we get the j th coordinate of $G(h)$ (for any $j = 1, \dots, m$) is equal to $f_m(h)$ which is introduced in (3.14). Other coordinates of $G(h)$ are equal to 0. By Lemma 3 we have that f_m is an increasing function. Here we consider the case when the function f_m has three fixed points $0, h_1, h_2$. This proof is more simple for cases when f_m has two fixed points. We prove the part 2), other parts are similar. In case 2), by Lemma 3 we have that $0 < h_1 < h_2$ and the point h_1 is a repeller, i.e., $f'_m(h_1) > 1$ and the points $0, h_2$ are attractive, i.e., $f'_m(0) < 1, f'_m(h_2) < 1$. Now we shall take arbitrary $x_0 > 0$ and prove that $x_n = f_m(x_{n-1}), n \geq 1$ converges as $n \rightarrow \infty$. Consider the following partition $(-\infty, +\infty) = (-\infty, 0) \cup \{0\} \cup (0, h_1) \cup \{h_1\} \cup (h_1, h_2) \cup \{h_2\} \cup (h_2, +\infty)$. For any $x \in (-\infty, 0)$ we have $x < f_m(x) < 0$, since f_m is an increasing function, from the last inequalities we get $x < f_m(x) < f_m^2(x) < f_m(0) = 0$. Iterating this argument we obtain $f_m^{n-1}(x) < f_m^n(x) < 0$, which for any $x_0 \in (-\infty, 0)$ gives $x_{n-1} < x_n < 0$, i.e., x_n converges and its limit is a fixed point of f_m , since f_m has unique fixed point 0 in $(-\infty, 0]$ we conclude that the limit is 0. For $x \in (0, h_1)$ we have $h_1 > x > f(x) > 0$, consequently $x_n > x_{n+1}$, i.e., x_n converges and its limit is again 0. Similarly, one can show that if $x_0 > h_1$ then $x_n \rightarrow h_2$ as $n \rightarrow \infty$. \square

By (3.10) the asymptotic behavior of the vector $Y_n(\omega) = (Y_n^1(\omega), \dots, Y_n^{q-1}(\omega))$ depends only on the vector $Y_1(\omega)$, where

$$Y_1^l(\omega) = J \left(c^l(\omega) - c^q(\omega) \right), \quad l = 1, \dots, q-1. \quad (4.6)$$

For a given $m \in \{1, \dots, \lfloor \frac{q}{2} \rfloor\}$ and $J > 0$ we introduce the following sets of configurations:

$$\mathbb{B}_m = \{\omega \in \Omega : c^1(\omega) = \dots = c^m(\omega), \quad c^{m+1}(\omega) = \dots = c^{q-1}(\omega) = c^q(\omega)\},$$

$$\mathbb{B}_{m,0}^+ = \{\omega \in \mathbb{B}_m : c^1(\omega) > c^q(\omega)\},$$

$$\mathbb{B}_{m,0}^0 = \{\omega \in \mathbb{B}_m : c^1(\omega) = c^q(\omega)\},$$

$$\mathbb{B}_{m,0}^- = \{\omega \in \mathbb{B}_m : c^1(\omega) < c^q(\omega)\},$$

$$\mathbb{B}_{m,1}^+ = \{\omega \in \mathbb{B}_m : J (c^1(\omega) - c^q(\omega)) > h_1\},$$

$$\mathbb{B}_{m,1}^0 = \{\omega \in \mathbb{B}_m : J (c^1(\omega) - c^q(\omega)) = h_1\},$$

$$\mathbb{B}_{m,1}^- = \{\omega \in \mathbb{B}_m : J (c^1(\omega) - c^q(\omega)) < h_1\}.$$

Now taking the coordinates of an initial vector as in (4.6) by Lemma 1 and Lemma 4 we get the following

Theorem 1. 1) If $\theta = \theta_m$, for some $m = 1, \dots, \lfloor \frac{q}{2} \rfloor$ then

$$P^\omega = \begin{cases} \mu_1(\theta, m), & \text{if } \omega \in \mathbb{B}_{m,1}^+ \cup \mathbb{B}_{m,1}^0 \\ \mu_0(\theta), & \text{if } \omega \in \mathbb{B}_{m,1}^- \end{cases} \quad (4.7)$$

2) If $\theta_m < \theta < \theta_c = q + 1$ then

$$P^\omega = \begin{cases} \mu_2(\theta, m), & \text{if } \omega \in \mathbb{B}_{m,1}^+ \\ \mu_1(\theta, m), & \text{if } \omega \in \mathbb{B}_{m,1}^0 \\ \mu_0(\theta), & \text{if } \omega \in \mathbb{B}_{m,1}^- \end{cases} \quad (4.8)$$

3) If $\theta = \theta_c$ then

$$P^\omega = \begin{cases} \mu_2(\theta, m), & \text{if } \omega \in \mathbb{B}_{m,0}^+ \\ \mu_0(\theta), & \text{if } \omega \in \mathbb{B}_{m,0}^- \cup \mathbb{B}_{m,0}^0 \end{cases} \quad (4.9)$$

4) If $\theta > \theta_c$ then

$$P^\omega = \begin{cases} \mu_2(\theta, m), & \text{if } \omega \in \mathbb{B}_{m,0}^+ \\ \mu_1(\theta, m), & \text{if } \omega \in \mathbb{B}_{m,0}^- \\ \mu_0(\theta), & \text{if } \omega \in \mathbb{B}_{m,0}^0 \end{cases} \quad (4.10)$$

In the next section we use Theorem 1 to construct some concrete boundary conditions.

5 Construction of boundary conditions

In this section for $k = 2$, $J > 0$, $q \geq 3$ and $m \in \{1, \dots, \lfloor \frac{q}{2} \rfloor\}$ we shall give examples of boundary configurations.

For $k = 2$ by Proposition 1 there are up to $2^q - 1$ TISGMs. We shall consider only $\mu_0(\theta)$ corresponding to $h = (0, 0, \dots, 0)$ and $\mu_i(\theta, m)$ corresponding to vector $\mathbf{h}(m, i) = (\underbrace{h_i, h_i, \dots, h_i}_m, \underbrace{0, 0, \dots, 0}_{q-m-1})$, $i = 1, 2$ with

$$\begin{aligned} h_1 &= 2 \ln \frac{\theta - 1 - \sqrt{(\theta - 1)^2 - 4m(q - m)}}{2m}, \\ h_2 &= 2 \ln \frac{\theta - 1 + \sqrt{(\theta - 1)^2 - 4m(q - m)}}{2m}. \end{aligned} \quad (5.1)$$

Using Theorem 1 we shall give some boundary conditions for each measure μ_i . Boundary conditions for the remaining measures can be obtained by using the permutation symmetry of the Potts model.

Case μ_0 . If $\theta < \theta_1$ then μ_0 is a unique measure, and one can take any boundary configuration ω to have $P^\omega = \mu_0$. But for $\theta \geq \theta_1$ one has to check the conditions of Theorem 1 to have the limiting measure equal to μ_0 .

For example, if $\theta = \theta_m < q + 1$ for some $m \in \{1, \dots, \lfloor \frac{q}{2} \rfloor\}$ then we must take $\omega \in \mathbb{B}_{m,1}^-$, i.e.

$$J(c^1(\omega) - c^q(\omega)) < h_1, \quad c^1(\omega) = \dots = c^m(\omega), \quad c^{m+1}(\omega) = \dots = c^q(\omega). \quad (5.2)$$

Remark 1. For a given TISGM to find its boundary condition one has to construct configurations ω which satisfy the system (like (5.2)) derived by corresponding sufficient conditions of Theorem 1. Below we give several examples of such configurations. It will be clear from our examples that some TISGM may have an infinite set of boundary configurations³.

Since $h_1 > 0$ (see Lemma 3), the system (5.2) is satisfied, for example, if ω satisfies one of the following

- $c^i(\omega) = 0$, $i = 1, 2, \dots, m$, i.e., if $\omega(x) = i$, then on direct successors x_1, x_2 of x one has $\omega(x_1) \neq i$, $\omega(x_2) \neq i$; and $c^j(\omega) = 1$ for each $j = m + 1, \dots, q$, i.e., if $\omega(x) = j$ then one has $\omega(x_1) = j$ but $\omega(x_2) \neq j$. See Fig.5 for an example of such a configuration for $q = 5$ and $m = 2$.
- $c^i(\omega) = 1$, $i = 1, 2, \dots, m$, i.e., if $\omega(x) = i$ then $\omega(x_1) = i$ but $\omega(x_2) \neq i$ and it contains $j \in \{m + 1, \dots, q\}$ in such a way that if $\omega(x) = j$ then on direct successors x_1, x_2 of x one has $\omega(x_1) = \omega(x_2) = j$. In this case $c^i(\omega) = 2$ for each $i = m + 1, \dots, q$. See Fig.6 for an example of such a configuration for $q = 15$ and $m = 3$.

Case μ_1 . This measure exists for $\theta \geq \theta_m$.

Subcase $\theta = \theta_m$. By Theorem 1 for μ_1 we have the condition $\omega \in \mathbb{B}_{m,1}^+ \cup \mathbb{B}_{m,1}^0$, i.e.

$$J(c^1(\omega) - c^q(\omega)) \geq h_1, \quad c^1(\omega) = \dots = c^m(\omega), \quad c^{m+1}(\omega) = \dots = c^q(\omega). \quad (5.3)$$

Note that $J = \ln \theta$. Assume $\ln \theta_m \geq h_1 = 2 \ln \frac{\theta_m - 1}{2m}$ which is equivalent to the following

$$2m\sqrt{m^2 + 1}(\sqrt{m^2 + 1} - m) \leq q \leq 2m\sqrt{m^2 + 1}(\sqrt{m^2 + 1} + m). \quad (5.4)$$

³This remark and some examples below are added corresponding to a suggestion of a reviewer.

Under condition (5.4) the system (5.3) is satisfied *for example*, if $m \geq 1$ and ω is such that $c^i(\omega) = 1, i = 1, \dots, m$ and $c^j(\omega) = 0, j = m + 1, \dots, q$. See Fig.7 for an example of such a configuration for $q = 5$ and $m = 2$ and Fig.8 for configuration in case $q = 10, m = 4$.

Subcase $\theta_m < \theta < q + 1$. From Theorem 1 for μ_1 we have the condition

$$J(c^1(\omega) - c^q(\omega)) = h_1, \quad c^1(\omega) = \dots = c^m(\omega), \quad c^{m+1}(\omega) = \dots = c^q(\omega). \quad (5.5)$$

Assume θ is a solution to the equation $\ln \theta = h_1$. Computer analysis shows that this equation has a solution if for example $q = 17, m = 1$ or $q = 55, m = 2$. So assuming existence of such a solution $\theta = \theta^*$ one can take a boundary condition configuration as in the previous case (like in Fig.7)

Subcase $\theta = q + 1$. In this case we have $\mu_1 = \mu_0$. Therefore the boundary condition can be taken as in Case μ_0 .

Subcase $\theta > q + 1$. For μ_1 we should have

$$c^1(\omega) - c^q(\omega) < 0, \quad c^1(\omega) = \dots = c^m(\omega), \quad c^{m+1}(\omega) = \dots = c^q(\omega). \quad (5.6)$$

we can take a configuration ω such that $c^i(\omega) = 0, i = 1, \dots, m$ and $c^j(\omega) = 1, j = m + 1, \dots, q$. (See Fig.5 for such a configuration).

Remark 2. *From above examples one can see that depending on the temperature (equivalently depending on the parameter θ) a configuration may be the boundary condition for different TISGMs. For example, the configuration given in Fig.5 is a boundary condition for TISGM μ_0 if $\theta = \theta_2 < q + 1$, but the same configuration is the boundary condition for TISGM μ_1 if $\theta > q + 1$.*

Case μ_2 . Check the conditions of Theorem 1:

Subcase $\theta = \theta_m$. In this case we have $\mu_2 = \mu_1$, i.e. the boundary condition is constructed in the previous case.

Subcase $\theta_m < \theta < q + 1$. From Theorem 1 for μ_2 we have the condition

$$J(c^1(\omega) - c^q(\omega)) > h_1, \quad c^1(\omega) = \dots = c^m(\omega), \quad c^{m+1}(\omega) = \dots = c^q(\omega). \quad (5.7)$$

If $\ln \theta > h_1$ then it is easy to see that ω satisfies the condition (5.7) if $m \geq 1$ and ω is such that $c^i(\omega) = 1, i \in \{1, \dots, m\}$ and $c^j(\omega) = 0, j \in \{m + 1, \dots, q\}$ (like in Fig.7)

Subcase $\theta \geq q + 1$. For μ_2 we should have

$$c^1(\omega) - c^q(\omega) > 0, \quad c^1(\omega) = \dots = c^m(\omega), \quad c^{m+1}(\omega) = \dots = c^q(\omega). \quad (5.8)$$

Condition (5.8) is easily checkable. For example, configurations shown in Fig. 7 and Fig.8 satisfy this condition.

Acknowledgements

U.Rozikov thanks Aix-Marseille University Institute for Advanced Study IMÉRA (Marseille, France) for support by a residency scheme. We thank both referees for their helpful suggestions.

References

- [1] L. Coquille, *Examples of DLR states which are not weak limits of finite volume Gibbs measures with deterministic boundary conditions*, Jour. Stat. Phys. **159**(4) (2015), 958-971.
- [2] S. Friedli, Y. Velenik, *Statistical Mechanics of Lattice Systems: a Concrete Mathematical Introduction*. Cambridge University Press, 2017. <http://www.unige.ch/math/folks/velenik/smbook/index.html>
- [3] D.Gandolfo, M.M. Rakhmatullaev, U.A. Rozikov, J. Ruiz, *On free energies of the Ising model on the Cayley tree*. Jour. Stat. Phys. **150**(6) (2013), 1201–1217.
- [4] N.N. Ganikhodjaev, *On pure phases of the three-state ferromagnetic Potts model on the second-order Bethe lattice*. Theor. Math. Phys. **85**(2) (1990), 1125–1134.
- [5] N.N. Ganikhodzhaev, *On pure phases of the ferromagnetic Potts model Bethe lattices*, Dokl. AN Uzbekistan. No. 6-7 (1992), 4–7.
- [6] H.O. Georgii, *Gibbs measures and phase transitions*, [2nd ed.]. (Walter de Gruyter, Berlin, 2011).
- [7] C. Kuelske, U.A. Rozikov, R.M. Khakimov, *Description of all translation-invariant (splitting) Gibbs measures for the Potts model on a Cayley tree*, Jour. Stat. Phys. **156**(1) (2014), 189–200.
- [8] C. Kuelske, U.A. Rozikov, *Fuzzy transformations and extremality of Gibbs measures for the Potts model on a Cayley tree*. Random Structures and Algorithms. 2016. DOI 10.1002/rsa.20671. 43 pages.
- [9] Y. Higuchi, *Remarks on the limiting Gibbs states on a $(d+1)$ -tree*. Publ. RIMS, Kyoto Univ. **3** (1977), 335–348.
- [10] U.A. Rozikov, *Gibbs measures on Cayley trees*. World Sci. Publ. Singapore. 2013.
- [11] U.A. Rozikov, *On pure phase of the anti-ferromagnetic Potts model on the Cayley tree*, Uzbek Math. J. No. 1 (1999), 73–77 (Russian).




Figure 1: Here and in Figures 2-4 the streamlines of the vector field $G^{(n)}(v)$ for $k = 2$, $q = 3$ are shown. These figures also illustrate the limit points of (4.1). The plane is formed by a horizontal v_1 -axis, and a vertical v_2 -axis. This figure applies for the case $\theta = \theta_1 = 1 + 2\sqrt{2}$. Four fixed points. The origin is an attractor. There are 3 saddle fixed points.




Figure 2: $\theta = 3.9 > \theta_1$. Seven fixed points. Four of them (black dots) are attractors. Three (rectangular dots) points are saddles coming from the saddle points of Fig.1.




Figure 3: $\theta = 4$. Four fixed points. The origin is a repeller point. Other fixed points are attractors.




Figure 4: $\theta = 4.5$. Seven fixed points. The origin is a repeller, other rectangular dots are saddles. The black dots are attractors.




Figure 5: An example of a boundary condition for the TISGM μ_0 , $q = 5$, $m = 2$. Here $c^1(\omega) = c^2(\omega) = 0$, $c^3(\omega) = c^4(\omega) = c^5(\omega) = 1$. Note that the values 1, 2 occur sufficiently often keeping the conditions $c^1(\omega) = c^2(\omega) = 0$.




Figure 6: An example of a boundary condition for the TISGM μ_0 , for $q = 15$, $m = 3$. Here we have $c^i(\omega) = 1$, $i = 1, 2, 3$ and $c^j(\omega) = 2$, $j = 4, 5, \dots, 15$.




Figure 7: An example of a boundary condition for the TISGM μ_1 , $q = 5$, $m = 2$. Here we have $c^1(\omega) = c^2(\omega) = 1$, $c^3(\omega) = c^4(\omega) = c^5(\omega) = 0$.




Figure 8: An example of a boundary condition for the TISGM μ_1 , $q = 10$, $m = 4$. Here we have $c^i(\omega) = 1$, $i = 1, 2, 3, 4$; $c^j(\omega) = 0$, $j = 5, \dots, 10$.