



A new proof of Smorynski's theorem

Apoloniusz Tyszk

► To cite this version:

| Apoloniusz Tyszk. A new proof of Smorynski's theorem. 2017. hal-01591775v1

HAL Id: hal-01591775

<https://hal.science/hal-01591775v1>

Preprint submitted on 22 Sep 2017 (v1), last revised 11 Sep 2019 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A new proof of Smoryński's theorem

Apoloniusz Tyszk

Abstract. We prove: (1) the set of all Diophantine equations which have at most finitely many solutions in non-negative integers is not recursively enumerable, (2) the set of all Diophantine equations which have at most finitely many solutions in positive integers is not recursively enumerable, (3) the set of all Diophantine equations which have at most finitely many integer solutions is not recursively enumerable. Analogous theorems hold for Diophantine equations $D(x_1, \dots, x_p) = 0$, where $p \in \mathbb{N} \setminus \{0\}$ and for every $i \in \{1, \dots, p\}$ the polynomial $D(x_1, \dots, x_p)$ involves a monomial M with a non-zero coefficient such that x_i divides M .

Key words and phrases: Diophantine equation which has at most finitely many solutions in non-negative integers, Diophantine equation which has at most finitely many solutions in positive integers, Diophantine equation which has at most finitely many integer solutions, Hilbert's Tenth Problem, Matiyasevich's theorem, recursively enumerable set, Smoryński's theorem.

2010 Mathematics Subject Classification: 03D25, 11U05.

There is no algorithm to decide whether or not a given Diophantine equation has an integer solution ([2]). The same is true for solutions in non-negative integers and for solutions in positive integers ([2]). The set of all Diophantine equations which have at most finitely many solutions in non-negative integers is not recursively enumerable, see [3, p. 104, Corollary 1] and [4, p. 240]. Let \mathcal{E} denote the set of all Diophantine equations $D(x_1, \dots, x_p) = 0$ such that $p \in \mathbb{N} \setminus \{0\}$ and the polynomial $D(x_1, \dots, x_p)$ truly depends on all the variables x_1, \dots, x_p . The last phrase means that for every $i \in \{1, \dots, p\}$ the polynomial $D(x_1, \dots, x_p)$ involves a monomial M with a non-zero coefficient such that x_i divides M .

Lemma 1. *A Diophantine equation $D(x_1, \dots, x_p) = 0$ has no solutions in non-negative integers x_1, \dots, x_p if and only if the equation $D(x_1, \dots, x_p) + 0 \cdot x_{p+1} = 0$ has at most finitely many solutions in non-negative integers x_1, \dots, x_{p+1} .*

Lemma 1a. *A Diophantine equation $D(x_1, \dots, x_p) = 0$ has no solutions in non-negative integers x_1, \dots, x_p if and only if the equation $(2x_{p+1} + 1) \cdot D(x_1, \dots, x_p) = 0$ has at most finitely many solutions in non-negative integers x_1, \dots, x_{p+1} .*

Lemma 2. A Diophantine equation $D(x_1, \dots, x_p) = 0$ has no solutions in positive integers x_1, \dots, x_p if and only if the equation $D(x_1, \dots, x_p) + 0 \cdot x_{p+1} = 0$ has at most finitely many solutions in positive integers x_1, \dots, x_{p+1} .

Lemma 2a. A Diophantine equation $D(x_1, \dots, x_p) = 0$ has no solutions in positive integers x_1, \dots, x_p if and only if the equation $(2x_{p+1} + 1) \cdot D(x_1, \dots, x_p) = 0$ has at most finitely many solutions in positive integers x_1, \dots, x_{p+1} .


Lemma 3. A Diophantine equation $D(x_1, \dots, x_p) = 0$ has no solutions in integers x_1, \dots, x_p if and only if the equation $D(x_1, \dots, x_p) + 0 \cdot x_{p+1} = 0$ has at most finitely many solutions in integers x_1, \dots, x_{p+1} .

Lemma 3a. A Diophantine equation $D(x_1, \dots, x_p) = 0$ has no solutions in integers x_1, \dots, x_p if and only if the equation $(2x_{p+1} + 1) \cdot D(x_1, \dots, x_p) = 0$ has at most finitely many solutions in integers x_1, \dots, x_{p+1} .

Lemma 4. If a polynomial $D(x_1, \dots, x_p) \in \mathbb{Z}[x_1, \dots, x_p]$ truly depends on all the variables x_1, \dots, x_p , then the polynomial $(2x_{p+1} + 1) \cdot D(x_1, \dots, x_p)$ truly depends on all the variables x_1, \dots, x_{p+1} .

Theorem 1. If the set of all Diophantine equations which have at most finitely many solutions in non-negative integers is recursively enumerable, then there exists an algorithm which decides whether or not a given Diophantine equation has a solution in non-negative integers.

Proof. Suppose that $\{S_i = 0\}_{i=2}^\infty$ is a computable sequence of all Diophantine equations which have at most finitely many solutions in non-negative integers. The algorithm presented in Flowchart 1 uses a computable surjection from $\mathbb{N} \setminus \{0, 1\}$ onto \mathbb{N}^p . By this and Lemma 1, the execution of Flowchart 1 decides whether or not a Diophantine equation $D(x_1, \dots, x_p) = 0$ has a solution in non-negative integers.



Flowchart 1


□

Corollary 1. *By Matiyasevich's theorem, the set of all Diophantine equations which have at most finitely many solutions in non-negative integers is not recursively enumerable.*

The analogous reasoning with Lemmas 1a and 4 shows that the set of all equations from \mathcal{E} which have at most finitely many solutions in non-negative integers is not recursively enumerable.

Theorem 2. *If the set of all Diophantine equations which have at most finitely many solutions in positive integers is recursively enumerable, then there exists an algorithm which decides whether or not a given Diophantine equation has a solution in positive integers.*

Proof. Suppose that $\{\mathcal{T}_i = 0\}_{i=2}^\infty$ is a computable sequence of all Diophantine equations which have at most finitely many solutions in positive integers. The algorithm presented in Flowchart 2 uses a computable surjection from $\mathbb{N} \setminus \{0, 1\}$ onto $(\mathbb{N} \setminus \{0\})^p$. By this and Lemma 2, the execution of Flowchart 2 decides whether or not a Diophantine equation $D(x_1, \dots, x_p) = 0$ has a solution in positive integers.



Flowchart 2


□

Corollary 2. *By Matiyasevich's theorem, the set of all Diophantine equations which have at most finitely many solutions in positive integers is not recursively enumerable.*

The analogous reasoning with Lemmas 2a and 4 shows that the set of all equations from \mathcal{E} which have at most finitely many solutions in positive integers is not recursively enumerable.

Theorem 3. *If the set of all Diophantine equations which have at most finitely many integer solutions is recursively enumerable, then there exists an algorithm which decides whether or not a given Diophantine equation has an integer solution.*

Proof. Suppose that $\{\mathcal{U}_i = 0\}_{i=2}^{\infty}$ is a computable sequence of all Diophantine equations which have at most finitely many integer solutions. There are infinitely many prime numbers of the form $3k + 1$ and there are infinitely many prime numbers of the form $3k + 2$, see [1, p. 80]. Hence, the algorithm presented in Flowchart 3 uses a computable surjection from $\mathbb{N} \setminus \{0, 1\}$ onto \mathbb{Z}^p . By this and Lemma 3, the execution of Flowchart 3 decides whether or not a Diophantine equation $D(x_1, \dots, x_p) = 0$ has an integer solution.



Flowchart 3

□

Corollary 3. *By Matiyasevich's theorem, the set of all Diophantine equations which have at most finitely many integer solutions is not recursively enumerable.*

The analogous reasoning with Lemmas 3a and 4 shows that the set of all equations from \mathcal{E} which have at most finitely many integer solutions is not recursively enumerable.

References

- [1] B. Fine and G. Rosenberger, *Number theory: An introduction via the distribution of primes*, Birkhäuser, Boston, MA, 2007.
- [2] Yu. Matiyasevich, *Hilbert's tenth problem*, MIT Press, Cambridge, MA, 1993.
- [3] C. Smoryński, *A note on the number of zeros of polynomials and exponential polynomials*, J. Symbolic Logic 42 (1977), no. 1, 99–106.
- [4] C. Smoryński, *Logical number theory, vol. I*, Springer, Berlin, 1991.

Apoloniusz Tyszką
University of Agriculture
Faculty of Production and Power Engineering
Balicka 116B, 30-149 Kraków, Poland
E-mail: rttyszka@cyf-kr.edu.pl