



HAL
open science

On stationary two-dimensional flows around a fast rotating disk

Isabelle Gallagher, Mitsuo Higaki, Yasunori Maekawa

► **To cite this version:**

Isabelle Gallagher, Mitsuo Higaki, Yasunori Maekawa. On stationary two-dimensional flows around a fast rotating disk. *Mathematical News / Mathematische Nachrichten*, 2019. hal-01592269

HAL Id: hal-01592269

<https://hal.science/hal-01592269>

Submitted on 23 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On stationary two-dimensional flows around a fast rotating disk

Isabelle Gallagher* Mitsuo Higaki† Yasunori Maekawa‡

September 23, 2017

Abstract We study the two-dimensional stationary Navier-Stokes equations describing flows around a rotating disk. The existence of unique solutions is established for any rotating speed, and qualitative effects of a large rotation are described precisely by exhibiting a boundary layer structure and an axisymmetrization of the flow.

Keywords Navier-Stokes equations · two-dimensional exterior flows · flows around a rotating obstacle · boundary layer

Mathematics Subject Classification (2000) 35B35 · 35Q30 · 76D05 · 76D17

1 Introduction

Understanding the structure of flows generated by the motion of a rigid body is an important subject in fluid dynamics. The typical motions of a rigid body (called the obstacle below) are translation and rotation, and it is well known that the flow generated by an obstacle translating with constant speed possesses a wake structure behind it, and that this structure is well described by the Oseen approximation; see Galdi [9] for mathematical results on this topic. Flows around a rotating obstacle have been studied mathematically mainly in three space dimensions, while there are only few mathematical results for two dimensions. Moreover, in the two dimensional case most known mathematical results are restricted to the case when the Reynolds number is sufficiently small. In general, the motion of the obstacle leads to a drastic change of the decay structure of the flow, and in the two dimensional case it yields a significant localizing effect that enables one to construct corresponding steady state solutions when the motion of the obstacle is slow enough. Although on the one hand a faster motion of the obstacle might give a stronger localizing and stabilizing effect, on the other hand fast motion produces a rapid flow and creates a strong shear near the boundary that can be a source of instability. As a result, rigorous analysis becomes quite difficult for the nonlinear problem in general. Hence it is useful to study the problem under a simple geometrical setting and to understand a typical fluid structure that describes these two competitive mechanisms; localizing and stabilizing effects on the one hand, and the presence of a rapid flow and the boundary layer created by the fast motion of the obstacle on the other hand.

*IMJ-PRG, Université Paris-Diderot and DMA, Ecole Normale Supérieure de Paris, France. E-mail: gallagher@math.ens.fr

†Department of Mathematics, Kyoto University, Japan. E-mail: mhigaki@math.kyoto-u.ac.jp

‡Department of Mathematics, Kyoto University, Japan. E-mail: maekawa@math.kyoto-u.ac.jp

In this paper we study two dimensional flows around a rotating obstacle assuming that the obstacle is a unit disk centered at the origin, in the case when the rotation speed is sufficiently fast and the Reynolds number is high. Note that in a three dimensional setting, these flows are considered as a model for two dimensional flows around a rotating infinite cylinder with a uniform cross section which is a unit disk.

Let us formulate our problem mathematically. For notational convenience we first state the problem when the obstacle has a general shape. Let us consider the following Navier-Stokes equations for viscous incompressible flows in two dimensions:

$$\begin{cases} \partial_t w - \Delta w + w \cdot \nabla w + \nabla \phi = g, & \operatorname{div} w = 0, & t > 0, y \in \Omega(t), \\ w = \alpha y^\perp, & & t > 0, y \in \partial\Omega(t). \end{cases} \quad (1.1)$$

Here $w = w(y, t) = (w_1(y_1, y_2, t), w_2(y_1, y_2, t))^\top$ and $\phi = \phi(y, t)$ are respectively the unknown velocity field and pressure field, and $g(y, t) = (g_1(y, t), g_2(y, t))^\top$ is an external force. The vector y^\perp is defined as $y^\perp = (-y_2, y_1)^\top$. Here M^\top denotes the transpose of a matrix M . The domain $\Omega(t)$ is given by

$$\begin{aligned} \Omega(t) &:= \{y \in \mathbb{R}^2 \mid y = O(\alpha t)x, x := (x_1, x_2)^\top \in \Omega\}, \\ O(\alpha t) &:= \begin{pmatrix} \cos \alpha t & -\sin \alpha t \\ \sin \alpha t & \cos \alpha t \end{pmatrix}, \end{aligned} \quad (1.2)$$

where Ω is an exterior domain in \mathbb{R}^2 and its complement $\Omega^c := \mathbb{R}^2 \setminus \Omega$ describes the obstacle at initial time, while the real number α represents the rotation speed of the obstacle. The condition $w(t, y) = \alpha y^\perp$ on the boundary $\partial\Omega(t)$ represents the noslip boundary condition. Taking into account the rotation of the obstacle, we introduce the following change of variables and unknowns:

$$y = O(\alpha t)x, \quad u(x, t) = O(\alpha t)^\top w(y, t), \quad p(x, t) = \phi(y, t), \quad f(x, t) = O(\alpha t)^\top g(y, t).$$

Then (1.1) is equivalent to the equations in the time-independent domain Ω :

$$\begin{cases} \partial_t u - \Delta u - \alpha(x^\perp \cdot \nabla u - u^\perp) + \nabla p = -u \cdot \nabla u + f, & \operatorname{div} u = 0, & t > 0, x \in \Omega, \\ u = \alpha x^\perp, & & t > 0, x \in \partial\Omega. \end{cases}$$

We are interested in stationary solutions to this system when the obstacle Ω^c is a unit disk centered at the origin, and therefore we assume that f is independent of t and consider the elliptic system

$$\begin{cases} -\Delta u - \alpha(x^\perp \cdot \nabla u - u^\perp) + \nabla p = -u \cdot \nabla u + f, & x \in \Omega, \\ \operatorname{div} u = 0, & x \in \Omega, \\ u = \alpha x^\perp, & x \in \partial\Omega, \end{cases} \quad (\text{NS}_\alpha)$$

with $\Omega := \{x \in \mathbb{R}^2 \mid |x| > 1\}$. We note that in the original coordinates the stationary solution to (NS_α) gives a specific time periodic flow with a periodicity $\frac{2\pi}{|\alpha|}$. Due to the symmetry of the domain there is an explicit stationary solution to (NS_α) when $f = 0$:

$$(\alpha U, \alpha^2 \nabla P) \quad \text{with} \quad U(x) := \frac{x^\perp}{|x|^2}, \quad P(x) := -\frac{1}{2|x|^2}. \quad (1.3)$$

Thus it is natural to consider an expansion around this explicit solution. By using the identity $u \cdot \nabla u = \frac{1}{2} \nabla |u|^2 + u^\perp \text{rot } u$ with $\text{rot } u := \partial_1 u_2 - \partial_2 u_1$ and $\text{rot } U = 0$ for $x \neq 0$, the equations for $v := u - \alpha U$ are written as

$$\begin{cases} -\Delta v - \alpha(x^\perp \cdot \nabla v - v^\perp) + \nabla q + \alpha U^\perp \text{rot } v = -v^\perp \text{rot } v + f, & x \in \Omega, \\ \text{div } v = 0, & x \in \Omega, \\ v = 0, & x \in \partial\Omega. \end{cases} \quad (\widetilde{\text{NS}}_\alpha)$$

The goal of this paper is to show the existence and uniqueness of solutions to $(\widetilde{\text{NS}}_\alpha)$ for arbitrary $\alpha \in \mathbb{R} \setminus \{0\}$ under a suitable condition on the given external force f in terms of regularity and summability, and moreover, we shall give a detailed qualitative analysis for the fast rotation case $|\alpha| \gg 1$.

Before stating our results let us recall known mathematical results related to this problem. Flows around a rotating obstacle have been studied mainly for the three dimensional case. Borchers [2] proved the existence of global weak solutions for the nonstationary problem, and the existence and uniqueness of local in time regular solutions is shown by Hishida [15] and Geissert, Heck, and Hieber [11]. Global strong solutions for small data are obtained by Galdi and Silvestre [10]. The spectrum of the linear operator related to this problem is studied by Hishida [16], and Farwig and Neustupa [7]. The existence of stationary solutions to the three dimensional problem is proved in [2], Galdi [8], Silvestre [20], and Farwig and Hishida [4]. In particular, [8] constructs stationary flows with a decay of order $O(|x|^{-1})$, while [4] discusses the weak L^3 framework. The asymptotic profiles of these stationary flows at spatial infinity are studied by Farwig and Hishida [5, 6] and Farwig, Galdi, and Kyed [3], where it is proved that the asymptotic profiles are described by the Landau solutions, which are stationary self-similar solutions to the Navier-Stokes equations in $\mathbb{R}^3 \setminus \{0\}$. The stability of small stationary solutions has been well studied in the three-dimensional case. Indeed, the global L^2 stability is proved in [10] and local L^3 stability is obtained by Hishida and Shibata [18].

All the results mentioned above are for the three dimensional case. So far there have been only few results in the two dimensional case. Hishida [17] revealed the asymptotic behavior of the stationary Stokes flow around a rotating obstacle, and showed that the rotation of the obstacle leads to the resolution of the Stokes paradox as in the case of the translation of the obstacle. On the other hand, the linear result in [17] was not sufficient to solve the nonlinear problem due to the nature of the singular perturbation limit $\alpha \rightarrow 0$ which is specific to the two dimensional case. Recently the linear result of [17] was extended by Higaki, Maekawa, and Nakahara [12], where the existence and uniqueness of solutions to (NS_α) decaying at the scale-critical order $O(|x|^{-1})$ was also proved when the rotation speed α is sufficiently small and the external force f is of a divergence form $f = \text{div } F$ for some F which is small in a scale critical norm. Moreover, the leading profile at spatial infinity was shown to be $C \frac{x^\perp}{|x|^2}$ for some constant C under an additional decay condition on F such as $F = O(|x|^{-2-r})$ with $r > 0$, which is compatible with the work of [17] for the Stokes case. The stability of these stationary solutions, though they are small in a scale-critical norm, is a difficult problem and is still largely open in the two dimensional case. The only known result is by Maekawa [19] for a specific case, which shows the local L^2 stability of the explicit solution (1.3) in the original frame (1.1) when α is small enough and Ω is the exterior to the unit disk as assumed in this paper. Few results are known in the case of nonsmall α . As a related work in this direction, Hillairet and Wittwer [14] established the existence of stationary solutions to (1.1) for large $|\alpha|$ when $\Omega(t) = \{y \in \mathbb{R}^2 \mid |y| > 1\}$ as in

this paper, although in [14] the boundary condition is $\alpha y^\perp + b$ instead of αy^\perp , with some time-independent given data b and the external force f is assumed to be zero. In [14] the stationary solution is constructed around (1.3) when $|\alpha|$ is large and b is small. Our problem is in fact essentially different from the one discussed in [14]. Indeed, the stationary solution to (NS_α) is a time periodic solution in the original frame (1.1), and therefore, the result in [14] is not applicable to our problem and vice versa. The reader is also referred to [13] for a model problem of (NS_α) formulated in the whole plane \mathbb{R}^2 , where the existence and uniqueness of stationary solutions is proved when $\alpha \neq 0$. However, the argument in [13] relies on the absence of physical boundary, and the key mechanism originating from the boundary is not analyzed.

Let us now return to $(\widetilde{\text{NS}}_\alpha)$. The novelty of the results in this paper is the followings:

- (1) Existence and uniqueness of solutions to $(\widetilde{\text{NS}}_\alpha)$ for arbitrary $\alpha \in \mathbb{R} \setminus \{0\}$.
- (2) Relaxed summability condition on f and on the class of solutions, allowing slow spatial decay with respect to scaling.
- (3) Qualitative analysis of solutions in the fast rotation case $|\alpha| \gg 1$.

As for (1), the result is new compared with [12] in which the stationary solutions are obtained only for nonzero but small $|\alpha|$, though there is no restriction on the shape of the obstacle in [12]. The reason why we can construct solutions for all nonzero α is a remarkable coercive estimate for the term $-\alpha(x^\perp \cdot \nabla v - v^\perp) + \alpha U^\perp \text{rot } v$ in polar coordinates; see (1.18) below. As for (2), we note that the given data f and the class of solutions in [12] are in a scale critical space. A typical behavior for f assumed in [12] is that $f = \text{div } F$ with $F(x) = O(|x|^{-2})$, and then the solution v satisfies the estimate $|v(x)| \leq C|x|^{-1}$ for $|x| \gg 1$. In this paper the summability condition on f is much weaker than this scaling, see (1.6) below. Moreover, the radial part of the solution constructed in this paper only behaves like $o(1)$ as $|x| \rightarrow \infty$ in general, which is considerably slow, while the nonradial part of the solution belongs to $L^2(\Omega)$ which is just in the scale critical regime. The point (3) is important both physically and mathematically. Understanding the fluid structure around the fast rotating obstacle up to the boundary is one of the main subjects of this paper, and we show the appearance of a boundary layer as well as an axisymmetrization mechanism due to the fast rotation of the obstacle.

Let us state our functional setting. Due to the symmetry of the domain it is natural to introduce the relevant function spaces in terms of polar coordinates. As usual, we set

$$\begin{aligned} x_1 &= r \cos \theta, & x_2 &= r \sin \theta, & r &= |x| \geq 1, & \theta &\in [0, 2\pi), \\ \mathbf{e}_r &= \frac{x}{|x|}, & \mathbf{e}_\theta &= \frac{x^\perp}{|x|} = \partial_\theta \mathbf{e}_r, \end{aligned}$$

and

$$v = v_r \mathbf{e}_r + v_\theta \mathbf{e}_\theta, \quad v_r = v \cdot \mathbf{e}_r, \quad v_\theta = v \cdot \mathbf{e}_\theta.$$

Next, for each $n \in \mathbb{Z}$, we denote by \mathcal{P}_n the projection on the Fourier mode n with respect to the angular variable θ :

$$\mathcal{P}_n v := v_{r,n} e^{in\theta} \mathbf{e}_r + v_{\theta,n} e^{in\theta} \mathbf{e}_\theta, \quad (1.4)$$

where

$$\begin{aligned} v_{r,n}(r) &:= \frac{1}{2\pi} \int_0^{2\pi} v_r(r \cos \theta, r \sin \theta) e^{-in\theta} d\theta, \\ v_{\theta,n}(r) &:= \frac{1}{2\pi} \int_0^{2\pi} v_\theta(r \cos \theta, r \sin \theta) e^{-in\theta} d\theta. \end{aligned}$$

We also set for $m \in \mathbb{N} \cup \{0\}$,

$$\mathcal{Q}_m v := \sum_{|n|=m+1}^{\infty} \mathcal{P}_n v. \quad (1.5)$$

For notational convenience we will often write v_n for $\mathcal{P}_n v$. Each \mathcal{P}_n is an orthogonal projection in $L^2(\Omega)^2$, and the space $L_\sigma^2(\Omega) := \overline{\{f \in C_0^\infty(\Omega)^2 \mid \operatorname{div} f = 0\}}^{L^2(\Omega)^2}$ is invariant under the action of \mathcal{P}_n . Note that $v_0 := \mathcal{P}_0 v$ is the radial part of v , and thus, $\mathcal{Q}_0 v$ is the nonradial part of v . We will set $\mathcal{P}_n L^2(\Omega)^2 := \{f \in L^2(\Omega)^2 \mid f = \mathcal{P}_n f\}$, and similar notation will be used for $L_\sigma^2(\Omega)$ and \mathcal{Q}_0 . A vector field f in Ω is formally identified with the pair $(\mathcal{P}_0 f, \mathcal{Q}_0 f)$. Then, for the class of external forces we introduce the product space

$$Y := \mathcal{P}_0 L^1(\Omega)^2 \times \mathcal{Q}_0 L^2(\Omega)^2. \quad (1.6)$$

For the class of solutions we set

$$X := \mathcal{P}_0 W_0^{1,\infty}(\Omega)^2 \times \mathcal{Q}_0 W_0^{1,2}(\Omega)^2. \quad (1.7)$$

Here $W_0^{1,r}(\Omega) := \{f \in W^{1,r}(\Omega) \mid f = 0 \text{ on } \partial\Omega\}$ for $1 < r \leq \infty$. Our first result is stated as follows.

Theorem 1.1 *There exists $\gamma > 0$ such that the following statements hold.*

(i) *Let $0 < |\alpha| < 1$. Then for any external force $f = (\mathcal{P}_0 f, \mathcal{Q}_0 f) \in Y$ satisfying*

$$\|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} \leq \gamma |\alpha|, \quad \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \leq \gamma |\alpha|^2, \quad (1.8)$$

there exists a solution $v \in X \cap L^\infty(\Omega)^2 \cap W_{loc}^{2,1}(\overline{\Omega})^2$ to $(\widetilde{\mathcal{N}}\mathcal{S}_\alpha)$ with a suitable pressure $q \in W_{loc}^{1,1}(\overline{\Omega})$, and v satisfies

$$\|\mathcal{P}_0 v\|_{L^\infty(\Omega)} + \|\nabla \mathcal{P}_0 v\|_{L^\infty(\Omega)} \leq C \|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} + \frac{C}{|\alpha|^{\frac{3}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}^2, \quad (1.9)$$

$$\|\mathcal{Q}_0 v\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (1.10)$$

$$\sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{3}{4}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (1.11)$$

$$\|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (1.12)$$

This solution is unique in a suitable closed convex set in X (see Subsection 4.2 for the precise description).

(ii) *Let $|\alpha| \geq 1$. Then for any external force $f = (\mathcal{P}_0 f, \mathcal{Q}_0 f) \in Y$ satisfying*

$$\|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} \leq \gamma, \quad \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \leq \gamma, \quad (1.13)$$

there exists a solution $v \in X \cap L^\infty(\Omega)^2 \cap W_{loc}^{2,1}(\overline{\Omega})^2$ to $(\widetilde{\text{NS}}_\alpha)$ with a suitable pressure $q \in W_{loc}^{1,1}(\overline{\Omega})$, and v satisfies

$$\|\mathcal{P}_0 v\|_{L^\infty(\Omega)} + \|\nabla \mathcal{P}_0 v\|_{L^\infty(\Omega)} \leq C \|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} + \frac{C}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}^2, \quad (1.14)$$

$$\|\mathcal{Q}_0 v\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (1.15)$$

$$\sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{4}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (1.16)$$

$$\|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} \leq C \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (1.17)$$

This solution is unique in a suitable subset of X .

Note that the summability of f assumed in Theorem 1.1 is much weaker than the scale-critical one. For the radial part $\mathcal{P}_0 v$ we can show $\lim_{|x| \rightarrow \infty} |\mathcal{P}_0 v(x)| = 0$ but there is no rate in general under the assumptions of Theorem 1.1. Theorem 1.1 already exhibits the axisymmetrizing effect of the fast rotating obstacle in L^2 and L^∞ , which will be further extended in Theorems 1.2 and 1.3 below. The proof of Theorem 1.1 consists in two ingredients: the analysis of the linearized problem (S_α) (defined and studied in Section 3), and the estimate of the interaction between the radial part and the nonradial part in the nonlinear problem (see Section 4). The linear result used in Theorem 1.1 is stated in Proposition 3.1, and the proof is based on an energy method. Although the proof of the linear result is not so difficult, there is a key observation for the term $-\alpha(x^\perp \cdot \nabla v - v^\perp) + \alpha U^\perp \text{rot } v$. Indeed, for the linearized problem (S_α) the energy computation for $v_n = \mathcal{P}_n v$ with $n \neq 0$ gives the key identity

$$\begin{aligned} & \alpha n \left(\|v_{r,n}\|_{L^2(\Omega)}^2 - \left(1 - \frac{2}{n^2}\right) \left\| \frac{v_{r,n}}{r} \right\|_{L^2(\Omega)}^2 + \|v_{\theta,n}\|_{L^2(\Omega)}^2 - \left\| \frac{v_{\theta,n}}{r} \right\|_{L^2(\Omega)}^2 \right) \\ & = -\mathfrak{S} \langle f_n, v_n \rangle_{L^2(\Omega)}. \end{aligned} \quad (1.18)$$

Here f_n denotes $\mathcal{P}_n f$ and the norm $\|g\|_{L^2(\Omega)}$ for the function $g : [1, \infty) \rightarrow \mathbb{C}$ is defined as $(2\pi)^{\frac{1}{2}} \|g\|_{L^2((1, \infty); r dr)}$. The key point here is that the bracket in (1.18) is nonnegative and provides a bound for $\left\| \frac{\sqrt{|x|^2 - 1}}{|x|} v_n \right\|_{L^2(\Omega)}^2$ since $\Omega = \{|x| > 1\}$. Then by combining with an interpolation inequality of the form

$$\|g\|_{L^2(\Omega)} \leq C \|\partial_r g\|_{L^2(\Omega)}^{\frac{1}{3}} \left\| \frac{\sqrt{r^2 - 1}}{r} g \right\|_{L^2(\Omega)}^{\frac{2}{3}} + C \left\| \frac{\sqrt{r^2 - 1}}{r} g \right\|_{L^2(\Omega)} \quad (1.19)$$

for any scalar function $g \in W^{1,2}((1, \infty); r dr)$ and the dissipation from the Laplacian in the energy computation, we can close the energy estimate for all $\alpha \neq 0$. The proof of (1.19) is given in Appendix B. In solving the nonlinear problem the key observation is that the product of the radial parts in the nonlinear term can always be written in a gradient form and thus regarded as a pressure term, which yields the identity

$$v^\perp \text{rot } v = v_0^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } v_0 + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{Q}_0 v + \nabla \tilde{q} \quad (1.20)$$

for a suitable \tilde{q} . Since $\mathcal{P}_0(v_0^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } v_0) = 0$ as long as $\mathcal{Q}_0 v \in W_0^{1,2}(\Omega)^2$ the radial part of the velocity in the right-hand side of (1.20) (neglecting $\nabla \tilde{q}$) belongs to $L^1(\Omega)^2$,

which is the same summability as the space Y . This is a brief explanation for the reason why we can close the nonlinear estimate and solve $(\widetilde{\text{NS}}_\alpha)$ in X for a source $f \in Y$.

Our second result is focused on the fast rotation case $|\alpha| \gg 1$. In this regime there are three fundamental mechanisms in our system:

- (I) an axisymmetrization due to the fast rotation of the obstacle,
- (II) the presence of a boundary layer for the nonradial part of the flow due to the noslip boundary condition,
- (III) the diffusion in high angular frequencies due to the viscosity.

(I) and (II) are potentially in a competitive relation, for the noslip boundary condition and the boundary layer can suppress the effect of the fast rotation to some extent.

(II) and (III) are also competitive. Indeed, it is natural that if the viscosity is strong enough then the boundary layer is diffused and is no longer observable. The important task here is to determine the regime of angular frequencies in which the boundary layer appears, and to estimate the thickness of the boundary layer. We show that the boundary layer appears in the regime $1 \leq |n| \ll O(|\alpha|^{\frac{1}{2}})$, and the thickness of the boundary layer is $(2|\alpha n|)^{-\frac{1}{3}}$ for each n in this regime. In constructing the boundary layer the term $\alpha U^\perp \text{rot } v$ plays a crucial role as well as the term $-\alpha(x^\perp \cdot \nabla v - v^\perp)$. In fact, if we drop the term $\alpha U^\perp \text{rot } v$ as a model problem then the thickness of the boundary layer arising from the rotation term $-\alpha(x^\perp \cdot \nabla v - v^\perp)$ is $|\alpha n|^{-\frac{1}{2}}$, and the leading boundary layer profile is simply described by exponential functions. The term $\alpha U^\perp \text{rot } v$ leads to a significant change both in the thickness and in the profile of the boundary layer, and we need to introduce the Airy function to describe the profile of the boundary layer associated with the term $-\alpha(x^\perp \cdot \nabla v - v^\perp) + \alpha U^\perp \text{rot } v$.

By performing the boundary layer analysis we can improve the result stated in (ii) of Theorem 1.1 in the regime $|\alpha| \gg 1$, which is briefly described as follows.

Theorem 1.2 *There exists $\gamma > 0$ such that the following statement holds. For all sufficiently large $|\alpha| \geq 1$ and for any external force $f = (\mathcal{P}_0 f, \mathcal{Q}_0 f) \in Y$ satisfying*

$$\|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} \leq \gamma |\alpha|^{\frac{1}{3}}, \quad \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \leq \gamma |\alpha|^{\frac{1}{3}}, \quad (1.21)$$

there exists a solution $v \in X \cap L^\infty(\Omega)^2 \cap W_{loc}^{2,1}(\overline{\Omega})^2$ to $(\widetilde{\text{NS}}_\alpha)$ with a suitable pressure $q \in W_{loc}^{1,1}(\overline{\Omega})$, and v satisfies

$$\|\mathcal{P}_0 v\|_{L^\infty(\Omega)} + \|\nabla \mathcal{P}_0 v\|_{L^\infty(\Omega)} \leq C \|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} + \frac{C}{|\alpha|} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}^2, \quad (1.22)$$

$$\|\mathcal{Q}_0 v\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{2}{3}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (1.23)$$

$$\sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} \leq \frac{C (\log |\alpha|)^{\frac{1}{2}}}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (1.24)$$

$$\|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{3}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (1.25)$$

This solution is unique in a suitable subset of X .

By fixing the external force f we can state Theorem 1.2 in a different but more convenient way to understand the qualitative behavior of solutions in the fast rotation limit.

Theorem 1.3 *For any $f = (\mathcal{P}_0 f, \mathcal{Q}_0 f) \in Y$ there is $\alpha_0 = \alpha_0(\|f\|_Y) \geq 1$ such that the following statements hold. If $|\alpha| \geq \alpha_0$ then there exists a solution $v = v^{(\alpha)} \in X \cap W_{loc}^{2,1}(\overline{\Omega})^2$ to $(\widetilde{\text{NS}}_\alpha)$ with a suitable pressure $q^{(\alpha)} \in W_{loc}^{1,1}(\overline{\Omega})$, and $v^{(\alpha)}$ satisfies*

$$\|v^{(\alpha)} - v_0^{\text{linear}}\|_{L^\infty(\Omega)} \leq \frac{C(\log |\alpha|)^{\frac{1}{2}}}{|\alpha|^{\frac{1}{2}}}. \quad (1.26)$$

Here v_0^{linear} is the solution to the linearized problem (S_α) defined in page 11 with f replaced by $\mathcal{P}_0 f$ which is in fact independent of α , and C depends only on $\|f\|_Y$. Moreover, there exists $\kappa > 0$ independent of α and f such that if $1 \leq |n| \leq \kappa|\alpha|^{\frac{1}{2}}$ then $v_n^{(\alpha)} = \mathcal{P}_n v^{(\alpha)}$ is written in the form

$$v_n^{(\alpha)} = v_n^{(\alpha),\text{slip}} + v_n^{(\alpha),\text{slow}} + v_{n,\text{BL}}^{(\alpha)} + \widetilde{v}_n^{(\alpha)}. \quad (1.27)$$

Here $v_n^{(\alpha),\text{slip}}$ satisfies $v_{n,r}^{(\alpha),\text{slip}} = \text{rot } v_n^{(\alpha),\text{slip}} = 0$ on $\partial\Omega$, $v_n^{(\alpha),\text{slow}}$ is irrotational in Ω , and $v_{n,\text{BL}}^{(\alpha)}$ possesses a boundary layer structure with the boundary layer thickness $|2\alpha n|^{-\frac{1}{3}}$. Finally the following estimates hold:

$$\|v_n^{(\alpha),\text{slip}}\|_{L^2(\Omega)} + \|v_n^{(\alpha),\text{slow}}\|_{L^2(\Omega)} + \|v_{n,\text{BL}}^{(\alpha)}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha n|^{\frac{2}{3}}},$$

while $\widetilde{v}_n^{(\alpha)}$ is a remainder which satisfies

$$\|\widetilde{v}_n^{(\alpha)}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha n|}.$$

Here the constant C depends only on $\|f\|_Y$.

By going back to (NS_α) , Theorems 1.2 and 1.3 show that there exists a unique solution $u = u^{(\alpha)}$ which satisfies

$$\|u^{(\alpha)} - \alpha U - v_0^{\text{linear}}\|_{L^\infty(\Omega)} \leq \frac{C(\log |\alpha|)^{\frac{1}{2}}}{|\alpha|^{\frac{1}{2}}}, \quad |\alpha| \gg 1. \quad (1.28)$$

The expansion (1.28) verifies the axisymmetrizing effect (measured in L^∞) due to the fast rotation. The logarithmic factor $(\log |\alpha|)^{\frac{1}{2}}$ is simply due to the regularity of f , and if f has more regularity such as $\sum_{n \neq 0} \|\mathcal{P}_n f\|_{L^2(\Omega)}^s < \infty$ for some $s < 2$ then the factor $(\log |\alpha|)^{\frac{1}{2}}$ in (1.26) and (1.28) can be dropped. Moreover, the power $|\alpha|^{-\frac{1}{2}}$ can be also improved by assuming enough regularity of f . For example, if $\mathcal{Q}_0 f \in W_0^{1,2}(\Omega)^2$ in addition, then $|\alpha|^{-\frac{1}{2}}$ is replaced by $|\alpha|^{-\frac{3}{4}}$, though we do not go into the detail on this point. The new ingredient of the proof of Theorems 1.2 and 1.3 is stated in Proposition 3.2 and consists in refined estimates for the linearized problem (S_α) . The nonlinear problem is handled exactly in the same manner as in the proof of Theorem 1.1. For (S_α) we observe that in polar coordinates the angular mode n of the streamfunction satisfies the ODE in $r \in (1, \infty)$

$$\left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{n^2}{r^2} + i\alpha n \left(1 - \frac{1}{r^2}\right) \right) \left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{n^2}{r^2} \right) \psi_n = 0, \quad (1.29)$$

with the boundary condition $\psi_n(1) = \frac{d\psi_n}{dr}(1) = 0$ when $|n| \geq 1$. The thickness of the boundary layer originating from the fast rotation is determined by the balance between $\frac{d^2}{dr^2}$ and $i\alpha n(1 - \frac{1}{r^2}) \approx 2i\alpha n(r-1)$ near $r = 1$ as long as the dissipation $-\frac{n^2}{r^2} \approx -n^2$ is moderate. This implies that the thickness is $|2\alpha n|^{-\frac{1}{3}}$. Then the regime of n where the dissipation is relatively moderate is estimated from the condition $n^2 \ll \frac{d^2}{dr^2} \approx O(|\alpha n|^{\frac{2}{3}})$, which leads to $|n| \ll O(|\alpha|^{\frac{1}{2}})$. From this observation we employ the boundary layer analysis when the angular frequency n satisfies $1 \leq |n| \ll O(|\alpha|^{\frac{1}{2}})$, while we just apply Proposition 3.1 in the regime $|n| \geq O(|\alpha|^{\frac{1}{2}})$ where the boundary layer due to the fast rotation is no longer present. In the regime $1 \leq |n| \ll O(|\alpha|^{\frac{1}{2}})$ we first consider (S_α) with $f = \mathcal{P}_n f$ but under the slip boundary condition $v_{r,n} = \text{rot } v_n = 0$ on $\partial\Omega$. The estimate of this *slip* solution is obtained by the energy method for the vorticity equations thanks to the boundary condition $\text{rot } v_n = 0$ on $\partial\Omega$. The key point is that the term $U^\perp \text{rot}$ in the velocity equation becomes $U \cdot \nabla$ in the vorticity equation which is antisymmetric because of $\text{div } U = 0$, and thus, it is easy to apply the energy method for the vorticity under the slip boundary condition. The *noslip* solution is then obtained by correcting the boundary condition. To this end we construct the boundary layer solution called the fast mode. The leading profile of the boundary layer is given by a suitable integral of the Airy function. In order to recover the noslip boundary condition, the fact that $\int_0^\infty \text{Ai}(s) ds \neq 0$ is crucial and it plays the role of a nondegeneracy condition in our construction of the solution. This construction gives a formula as in (1.27) for the solution to (S_α) . Compared with Proposition 3.1, which is based only on an energy computation for the velocity field, the estimate of the n mode v_n is drastically improved for $1 \leq |n| \ll O(|\alpha|^{\frac{1}{2}})$ thanks to the boundary layer analysis. On the other hand, in the regime $|n| \geq O(|\alpha|^{\frac{1}{2}})$, Proposition 3.1 for the *noslip* solution already gives the same decay estimates as in Proposition 3.4 for the *slip* solution, as expected.

This paper is organized as follows. In Section 2 we recall some basic facts on operators and vector fields in polar coordinates. In Section 3 the linearized problem (S_α) is studied. This section is the core of the paper. In Subsection 3.1 we prove the linear estimates which are valid for all $\alpha \in \mathbb{R} \setminus \{0\}$. These are summarized in Proposition 3.1. Subsection 3.2 is devoted to the linear analysis for the case $|\alpha| \gg 1$, and the main result of this section is Proposition 3.2. The nonlinear problem is discussed in Section 4. Some basics on the Airy function and the proof of the interpolation inequality (1.19) are given in the appendix.

2 Preliminaries

In this preliminary section we state basic results on some differential operators and vector fields in polar coordinates.

2.1 Operators in polar coordinates

The following formulas will be used frequently:

$$\operatorname{div} v = \partial_1 v_1 + \partial_2 v_2 = \frac{1}{r} \partial_r (r v_r) + \frac{1}{r} \partial_\theta v_\theta, \quad (2.1)$$

$$\operatorname{rot} v = \partial_1 v_2 - \partial_2 v_1 = \frac{1}{r} \partial_r (r v_\theta) - \frac{1}{r} \partial_\theta v_r, \quad (2.2)$$

$$|\nabla v|^2 = |\partial_r v_r|^2 + |\partial_r v_\theta|^2 + \frac{1}{r^2} (|\partial_\theta v_r - v_\theta|^2 + |v_r + \partial_\theta v_\theta|^2), \quad (2.3)$$

and

$$\begin{aligned} -\Delta v &= \left(-\partial_r \left(\frac{1}{r} \partial_r (r v_r) \right) - \frac{1}{r^2} \partial_\theta^2 v_r + \frac{2}{r^2} \partial_\theta v_\theta \right) \mathbf{e}_r \\ &\quad + \left(-\partial_r \left(\frac{1}{r} \partial_r (r v_\theta) \right) - \frac{1}{r^2} \partial_\theta^2 v_\theta - \frac{2}{r^2} \partial_\theta v_r \right) \mathbf{e}_\theta, \end{aligned} \quad (2.4)$$

$$\mathbf{e}_r \cdot \nabla v = (\partial_r v_r) \mathbf{e}_r + (\partial_r v_\theta) \mathbf{e}_\theta,$$

$$\mathbf{e}_\theta \cdot \nabla v = \frac{\partial_\theta v_r - v_\theta}{r} \mathbf{e}_r + \frac{\partial_\theta v_\theta + v_r}{r} \mathbf{e}_\theta.$$

In particular, we have

$$\begin{aligned} x^\perp \cdot \nabla v - v^\perp &= |x| (\mathbf{e}_\theta \cdot \nabla v) - (v_r \mathbf{e}_r^\perp + v_\theta \mathbf{e}_\theta^\perp) \\ &= (\partial_\theta v_r - v_\theta) \mathbf{e}_r + (\partial_\theta v_\theta + v_r) \mathbf{e}_\theta - (v_r \mathbf{e}_r^\perp + v_\theta \mathbf{e}_\theta^\perp) \\ &= \partial_\theta v_r \mathbf{e}_r + \partial_\theta v_\theta \mathbf{e}_\theta. \end{aligned} \quad (2.5)$$

From (2.3) and the definition of \mathcal{P}_n in (1.4) it follows that for $n \in \mathbb{N} \cup \{0\}$ and for v in $W^{1,2}(\Omega)^2$,

$$\begin{aligned} \|\nabla v\|_{L^2(\Omega)}^2 &= \sum_{n \in \mathbb{Z}} \|\nabla \mathcal{P}_n v\|_{L^2(\Omega)}^2, \\ |\nabla \mathcal{P}_n v|^2 &= |\partial_r v_{r,n}|^2 + \frac{1+n^2}{r^2} |v_{r,n}|^2 \\ &\quad + |\partial_r v_{\theta,n}|^2 + \frac{1+n^2}{r^2} |v_{\theta,n}|^2 - \frac{4n}{r^2} \Im(v_{\theta,n} \overline{v_{r,n}}). \end{aligned}$$

In particular, we have

$$|\nabla \mathcal{P}_n v|^2 \geq |\partial_r v_{r,n}|^2 + \frac{(|n|-1)^2}{r^2} |v_{r,n}|^2 + |\partial_r v_{\theta,n}|^2 + \frac{(|n|-1)^2}{r^2} |v_{\theta,n}|^2, \quad (2.6)$$

and thus, from the definition of \mathcal{Q}_m in (1.5),

$$\|\nabla \mathcal{Q}_m v\|_{L^2(\Omega)}^2 \geq \|\partial_r (\mathcal{Q}_m v)_r\|_{L^2(\Omega)}^2 + \|\partial_r (\mathcal{Q}_m v)_\theta\|_{L^2(\Omega)}^2 + m^2 \left\| \frac{v}{|x|} \right\|_{L^2(\Omega)}^2.$$

2.2 The Biot-Savart law in polar coordinates

For a given scalar field ω in Ω , the streamfunction ψ is formally defined as the solution to the Poisson equation: $-\Delta\psi = \omega$ in Ω , $\psi = 0$ on $\partial\Omega$. For $n \in \mathbb{Z}$ and $\omega \in L^2(\Omega)$ we set

$$\begin{aligned}\mathcal{P}_n\omega &:= \frac{1}{2\pi} \int_0^{2\pi} \omega(r \cos s, r \sin s) e^{-ins} ds e^{in\theta}, \\ \omega_n &:= (\mathcal{P}_n\omega) e^{-in\theta}.\end{aligned}\tag{2.7}$$

By using the Laplace operator in polar coordinates, the Poisson equation for the Fourier mode n is given by

$$H_n\psi_n := -\psi_n'' - \frac{1}{r}\psi_n' + \frac{n^2}{r^2}\psi_n = \omega_n, \quad r > 1, \quad \psi_n(1) = 0.\tag{2.8}$$

Let $|n| \geq 1$. Then the solution $\psi_n = \psi_n[\omega_n]$ to the ordinary differential equation (2.8) decaying at spatial infinity is formally given as

$$\begin{aligned}\psi_n[\omega_n](r) &= \frac{1}{2|n|} \left(-\frac{d_n[\omega_n]}{r^{|n|}} + \frac{1}{r^{|n|}} \int_1^r s^{1+|n|}\omega_n(s) ds + r^{|n|} \int_r^\infty s^{1-|n|}\omega_n(s) ds \right), \\ d_n[\omega_n] &:= \int_1^\infty s^{1-|n|}\omega_n(s) ds.\end{aligned}$$

The Biot-Savart law $V_n[\omega_n]$ is then written as

$$\begin{aligned}V_n[\omega_n] &:= V_{r,n}[\omega_n] e^{in\theta} \mathbf{e}_r + V_{\theta,n}[\omega_n] e^{in\theta} \mathbf{e}_\theta, \\ V_{r,n}[\omega_n] &:= \frac{in}{r}\psi_n[\omega_n], \quad V_{\theta,n}[\omega_n] := -\frac{d}{dr}\psi_n[\omega_n].\end{aligned}\tag{2.9}$$

The velocity $V_n[\omega_n]$ is well defined at least when $r^{1-|n|}\omega_n \in L^1((1, \infty))$, and it is straightforward to see that

$$\begin{aligned}\operatorname{div} V_n[\omega_n] &= 0, \quad \operatorname{rot} V_n[\omega_n] = \omega_n e^{in\theta} \quad \text{in } \Omega, \\ \mathbf{e}_r \cdot V_n[\omega_n] &= 0 \quad \text{on } \partial\Omega.\end{aligned}\tag{2.10}$$

The condition $r^{1-|n|}\omega_n \in L^1((1, \infty))$ is automatically satisfied when $\omega \in L^2(\Omega)$ and $|n| \geq 2$. When $|n| = 1$ the integral in the definition of $\psi_n[\omega_n]$ does not always converge absolutely for general $\omega \in L^2(\Omega)$. However, it is well-defined if $\omega = \operatorname{rot} u$ for some $u \in W^{1,2}(\Omega)^2$, for one can apply the integration by parts that ensures the convergence of $\lim_{N \rightarrow \infty} \int_r^N \omega_n dr$ even when $|n| = 1$. As a result, we can check that for any solenoidal vector field v in $L^2_\sigma(\Omega) \cap W^{1,2}(\Omega)^2$, the n mode $v_n = \mathcal{P}_n v$ is expressed in terms of its vorticity ω_n by the formula (2.9) when $|n| \geq 1$.

3 Analysis of the linearized system

The linearized system around αU for $(\widetilde{\text{NS}}_\alpha)$ is

$$\begin{cases} -\Delta v - \alpha(x^\perp \cdot \nabla v - v^\perp) + \nabla q + \alpha U^\perp \operatorname{rot} v = f, & x \in \Omega, \\ \operatorname{div} v = 0, & x \in \Omega, \\ v = 0, & x \in \partial\Omega. \end{cases}\tag{S_\alpha}$$

In this section we study (S_α) for $\alpha \in \mathbb{R} \setminus \{0\}$.

3.1 General estimate

In this subsection we establish estimates on solutions to (S_α) that are valid for all $\alpha \neq 0$. For convenience we set for scalar functions $g, h : [1, \infty) \rightarrow \mathbb{C}$,

$$\langle g, h \rangle_{L^2(\Omega)} := 2\pi \int_1^\infty g(r) \overline{h(r)} r \, dr, \quad \|g\|_{L^2(\Omega)}^2 := 2\pi \int_1^\infty |g(r)|^2 r \, dr.$$

Before going into details let us give a remark on the verification of the energy argument. Let us assume that $f \in L^2(\Omega)^2$ and $v \in W_0^{1,2}(\Omega)^2 \cap W_{loc}^{2,2}(\overline{\Omega})^2$ is a solution to (S_α) for some $q \in W_{loc}^{1,2}(\overline{\Omega})$. We have to be careful when applying the energy argument due to the presence of the term $x^\perp \cdot \nabla v$ in the first equation of (S_α) , for this term involves a linearly growing coefficient, and therefore it is not clear whether the inner product $\langle x^\perp \cdot \nabla v, v \rangle_{L^2(\Omega)}$ makes sense or not. A similar difficulty appears in taking the inner product $\langle \nabla q, v \rangle_{L^2(\Omega)}$, since we are assuming only $q \in W_{loc}^{1,2}(\overline{\Omega})$. The most convenient way to overcome this difficulty is to consider the equation for $v_n := \mathcal{P}_n v$, which is identified with $(v_{r,n}, v_{\theta,n})$. Note that $v_{r,n}, v_{\theta,n} \in W_0^{1,2}((1, \infty); r \, dr) \cap W_{loc}^{2,2}([1, \infty))$ if $v \in W_0^{1,2}(\Omega)^2 \cap W_{loc}^{2,2}(\overline{\Omega})^2$. Let us denote by $\omega_n = \omega_n(r)$ the n mode of the vorticity of v in the polar coordinates, i.e., $\omega_n(r) = (\text{rot } \mathcal{P}_n v) e^{-in\theta}$. Similarly, we set $q_n = q_n(r) = (\mathcal{P}_n q) e^{-in\theta}$, where the projection \mathcal{P}_n for the scalar q is defined as

$$\mathcal{P}_n q := \frac{1}{2\pi} \int_0^{2\pi} q(r \cos s, r \sin s) e^{-ins} \, ds e^{in\theta}.$$

Then, from (2.4) and (2.5), $v_{r,n}$ and $v_{\theta,n}$ obey the following equations:

$$-\partial_r \left(\frac{1}{r} \partial_r (r v_{r,n}) \right) + \frac{n^2}{r^2} v_{r,n} + \frac{2in}{r^2} v_{\theta,n} - i\alpha n v_{r,n} - \alpha \frac{\omega_n}{r} + \partial_r q_n = f_{r,n}, \quad (3.1)$$

$$-\partial_r \left(\frac{1}{r} \partial_r (r v_{\theta,n}) \right) + \frac{n^2}{r^2} v_{\theta,n} - \frac{2in}{r^2} v_{r,n} - i\alpha n v_{\theta,n} + in q_n = f_{\theta,n}, \quad (3.2)$$

together with the divergence free condition $\partial_r (r v_{r,n}) + in v_{\theta,n} = 0$ and the noslip boundary condition $v_{r,n}(1) = v_{\theta,n}(1) = 0$. Then the key observation is that the factor $-i\alpha n$ is now regarded as a resolvent parameter, and by setting $\lambda := -i\alpha n$, the above system is equivalent to

$$(\lambda - \Delta) v_n + \nabla \mathcal{P}_n q + \alpha U^\perp \text{rot } v_n = f_n, \quad x \in \Omega, \quad (3.3)$$

with $\text{div } v_n = 0$ and $v_n|_{\partial\Omega} = 0$, where $f_n = \mathcal{P}_n f \in \mathcal{P}_n L^2(\Omega)^2$. Indeed, system (3.3) in polar coordinates is exactly (3.1) and (3.2). The key point is that there is no term involving a linearly growing coefficient in (3.3), and therefore we can apply the standard regularity theory of the Stokes resolvent system with a resolvent parameter λ . Let us assume that $n \neq 0$. Then $\lambda \neq 0$ since we are assuming that $\alpha \neq 0$. If $v_n \in L_\sigma^2(\Omega) \cap W_0^{1,2}(\Omega)^2 \cap W_{loc}^{2,2}(\overline{\Omega})^2$ and $\mathcal{P}_n q \in W_{loc}^{1,2}(\overline{\Omega})$ is a solution to (3.3), then $(v_n, \mathcal{P}_n q)$ is a weak solution to the Stokes system with source term $f_n - \alpha U^\perp \text{rot } v_n$ which clearly belongs to $L^2(\Omega)^2$, and thus, the regularity theory of the Stokes system implies that $v_n \in W^{2,2}(\Omega)^2$ and $\nabla \mathcal{P}_n q \in L^2(\Omega)^2$. In this way we can recover the summability of $\nabla^2 v_n, \nabla \mathcal{P}_n q \in L^2(\Omega)^2$. Then, by going back to the system (3.1) and (3.2), we also find that $q_n \in L^2((1, \infty); r \, dr)$ from (3.2), for all the other terms in (3.2) belong to $L^2([1, \infty); r \, dr)$. As a summary, for any solution $v \in$

$L^2_\sigma(\Omega) \cap W_0^{1,2}(\Omega)^2 \cap W_{loc}^{2,2}(\overline{\Omega})^2$ and $q \in W_{loc}^{1,2}(\overline{\Omega})$ to (S_α) , we can rigorously verify the energy computation for the system (3.1)-(3.2) in each n mode $(v_{r,n}, v_{\theta,n})$ with $n \neq 0$. The estimate for the 0 mode is handled in a different way from the energy method, and is discussed in Subsection 3.1.1 below.

Our main result in this subsection is stated as follows. Let us recall that the projection \mathcal{Q}_0 is defined as $\mathcal{Q}_0 := \sum_{n \neq 0} \mathcal{P}_n$.

Proposition 3.1 *Let $\alpha \in \mathbb{R} \setminus \{0\}$.*

(i) *For any external force $f_0 \in \mathcal{P}_0 L^1(\Omega)^2$ the system (S_α) admits a unique solution $(v_0, \nabla q)$ with $v_0 \in \mathcal{P}_0 L^\infty(\Omega)^2 \cap W_0^{1,\infty}(\Omega)^2$, $\nabla^2 v_0 \in L^1_{loc}(\overline{\Omega})^{2 \times 2}$, $q \in W_{loc}^{1,1}(\overline{\Omega})$. Moreover, $v_0 = v_{\theta,0} \mathbf{e}_\theta$ and*

$$\|v_0\|_{L^\infty(\Omega)} + \||x|\nabla v_0\|_{L^\infty(\Omega)} \leq C \|f_{\theta,0}\|_{L^1(\Omega)}. \quad (3.4)$$

Here $f_{\theta,0} := f_0 \cdot \mathbf{e}_\theta$ and C is independent of α .

(ii) *For any external force $f \in \mathcal{Q}_0 L^2(\Omega)^2$ the system (S_α) admits a unique solution $(v, \nabla q)$ with $v \in \mathcal{Q}_0 L^2_\sigma(\Omega) \cap W_0^{1,2}(\Omega)^2 \cap W_{loc}^{2,2}(\overline{\Omega})^2$ and $q \in W_{loc}^{1,2}(\overline{\Omega})$. Moreover, $v_n = \mathcal{P}_n v$ satisfies the following estimates: if $1 \leq |n| < 1 + \sqrt{2|\alpha|}$ then*

$$\|v_n\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \left(\frac{1}{|n|} + \frac{1}{|\alpha|^{\frac{1}{2}}} \right) \|f_n\|_{L^2(\Omega)}, \quad (3.5)$$

$$\left\| \frac{\sqrt{|x|^2 - 1}}{|x|} v_n \right\|_{L^2(\Omega)} \leq \frac{C}{|\alpha n|^{\frac{1}{2}} |\alpha|^{\frac{1}{4}}} \left(\frac{1}{|n|} + \frac{1}{|\alpha|^{\frac{1}{2}}} \right)^{\frac{1}{2}} \|f_n\|_{L^2(\Omega)}, \quad (3.6)$$

$$\|v_n\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{4}}} \left(\frac{1}{|n|} + \frac{1}{|\alpha|^{\frac{1}{2}}} \right) \|f_n\|_{L^2(\Omega)}, \quad (3.7)$$

$$\|\nabla v_n\|_{L^2(\Omega)} \leq C \left(\frac{1}{|n|} + \frac{1}{|\alpha|^{\frac{1}{2}}} \right) \|f_n\|_{L^2(\Omega)}, \quad (3.8)$$

while if $|n| \geq 1 + \sqrt{2|\alpha|}$ then

$$\|v_n\|_{L^2(\Omega)} \leq \frac{C}{|n|} \left(\frac{1}{|n|} + \frac{1}{|\alpha|} \right) \|f_n\|_{L^2(\Omega)}, \quad (3.9)$$

$$\left\| \frac{\sqrt{|x|^2 - 1}}{|x|} v_n \right\|_{L^2(\Omega)} \leq \frac{C}{|\alpha n|^{\frac{1}{2}} |n|^{\frac{1}{2}}} \left(\frac{1}{|n|} + \frac{1}{|\alpha|} \right)^{\frac{1}{2}} \|f_n\|_{L^2(\Omega)}, \quad (3.10)$$

$$\|v_n\|_{L^\infty(\Omega)} \leq \frac{C}{|n|^{\frac{3}{4}}} \left(\frac{1}{|n|} + \frac{1}{|\alpha|} \right)^{\frac{3}{4}} \|f_n\|_{L^2(\Omega)}, \quad (3.11)$$

$$\|\nabla v_n\|_{L^2(\Omega)} \leq \frac{C}{|n|^{\frac{1}{2}}} \left(\frac{1}{|n|} + \frac{1}{|\alpha|} \right)^{\frac{1}{2}} \|f_n\|_{L^2(\Omega)}. \quad (3.12)$$

Finally if $|\alpha| \gg 1$ and $|n| = O(|\alpha|^{\frac{1}{2}})$ then

$$\|v_n\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|} \|f_n\|_{L^2(\Omega)}, \quad (3.13)$$

$$\|v_n\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{3}{4}}} \|f_n\|_{L^2(\Omega)}, \quad (3.14)$$

$$\|\nabla v_n\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|f_n\|_{L^2(\Omega)}. \quad (3.15)$$

Here $f_n := \mathcal{P}_n f$ and C is independent of n and α .

3.1.1 Structure and estimate of the 0 mode

Firstly we observe that if v_0 satisfies $v_0 \in \mathcal{P}_0 L^\infty(\Omega)^2 \cap W_0^{1,\infty}(\Omega)^2$ then the divergence-free condition in polar coordinates (2.1) implies that

$$\frac{d(rv_{r,0})}{dr} = 0,$$

and thus, $v_{r,0} = \frac{C}{r}$ with some constant C . Then the noslip boundary condition leads to $C = 0$, and therefore, $v_{r,0} = 0$. So it suffices to consider the angular part $v_{\theta,0}$. From (2.5) we have

$$x^\perp \cdot \nabla v_0^\perp - v_0^\perp = 0,$$

and we also note that the term $U^\perp \text{rot } v_0$ in (S_α) with $U^\perp = -\frac{x}{|x|^2}$ is always written in a gradient form, so can be absorbed in a pressure term.

Collecting these remarks, we see that any solution $(v_0, \nabla q)$ to (S_α) with $f_0 \in \mathcal{P}_0 L^1(\Omega)^2$ satisfying $v_0 \in \mathcal{P}_0 L^\infty(\Omega)^2 \cap W_0^{1,\infty}(\Omega)^2$, $\nabla^2 v_0 \in L_{loc}^1(\overline{\Omega})^{2 \times 2}$, $q \in W_{loc}^{1,1}(\overline{\Omega})$ must be written as $v_0 = v_{0,\theta} e_\theta$, where $v_{\theta,0} = v_{\theta,0}(r)$ obeys from (3.2) the ordinary differential equation

$$-\frac{d^2 v_{\theta,0}}{dr^2} - \frac{1}{r} \frac{dv_{\theta,0}}{dr} + \frac{v_{\theta,0}}{r^2} = f_{\theta,0}, \quad r > 1, \quad v_{\theta,0}(1) = 0. \quad (3.16)$$

The bounded solution to (3.16) is written as

$$v_{\theta,0}(r) = \frac{1}{2} \left(-\frac{1}{r} \int_1^\infty f_{\theta,0} ds + \frac{1}{r} \int_1^r s^2 f_{\theta,0} ds + r \int_r^\infty f_{\theta,0} ds \right). \quad (3.17)$$

We note that

$$\|f_{\theta,0}\|_{L^1(\Omega)} = 2\pi \int_1^\infty |f_{\theta,0}| s ds.$$

Thus we see from (3.17) that

$$\|v_{\theta,0}\|_{L^\infty(\Omega)} + \left\| r \frac{dv_{\theta,0}}{dr} \right\|_{L^\infty(\Omega)} \leq C \|f_{\theta,0}\|_{L^1(\Omega)},$$

which implies (3.4).

3.1.2 A priori estimate of the n mode with $|n| \geq 1$

Let v denote a solution to (S_α) satisfying $v \in \mathcal{Q}L_\sigma^2(\Omega) \cap W_0^{1,2}(\Omega)^2 \cap W_{loc}^{2,2}(\overline{\Omega})^2$ for $f \in \mathcal{Q}_0 L^2(\Omega)^2$ with some $q \in W_{loc}^{1,2}(\overline{\Omega})$. Then, as we have already seen in the beginning of Section 3.1, for each $n \neq 0$, the n mode $v_n = \mathcal{P}_n v$ belongs in addition to $W^{2,2}(\Omega)^2$ and we also have that $\mathcal{P}_n q$ belongs to $W^{1,2}(\Omega)$. Hence the energy computation below for $(v_{r,n}, v_{\theta,n})$ to the system (3.1)-(3.2) is rigorously verified. With this important remark in mind we multiply both sides of (3.1) by $r\bar{v}_{r,n}$ and of (3.2) by $r\bar{v}_{\theta,n}$ and integrate over $[1, \infty)$, which results in the following identities:

$$\|\nabla v_n\|_{L^2(\Omega)}^2 = -\alpha \Re \langle U^\perp \text{rot } v_n, v_n \rangle_{L^2(\Omega)} + \Re \langle f_n, v_n \rangle_{L^2(\Omega)}, \quad (3.18)$$

$$-\alpha n (\|v_{r,n}\|_{L^2(\Omega)}^2 + \|v_{\theta,n}\|_{L^2(\Omega)}^2) + \alpha \Im \langle U^\perp \text{rot } v_n, v_n \rangle_{L^2(\Omega)} = \Im \langle f_n, v_n \rangle_{L^2(\Omega)}. \quad (3.19)$$

Note that $U^\perp(x) = -\frac{x}{|x|^2}$ thus we see from $\text{rot } v_n = -\Delta \psi_n$ by definition of the stream-function ψ_n for the n mode with $n \neq 0$,

$$\begin{aligned} \Re \langle U^\perp \text{rot } v_n, v_n \rangle_{L^2(\Omega)} &= \Re \langle \frac{1}{|x|} \Delta \psi_n, v_{r,n} \rangle_{L^2(\Omega)} \\ &= 2\pi n \Im \int_1^\infty (\partial_r^2 \psi_n + \frac{1}{r} \partial_r \psi_n - \frac{n^2}{r^2} \psi_n) \overline{\psi_n} \frac{dr}{r} \\ &= 4\pi n \Im \int_1^\infty \frac{1}{r} \partial_r \psi_n \overline{\psi_n} \frac{dr}{r}. \end{aligned}$$

This gives the bound

$$\left| \Re \langle U^\perp \text{rot } v_n, v_n \rangle_{L^2(\Omega)} \right| \leq 2 \left\| \frac{v_{r,n}}{r} \right\|_{L^2(\Omega)} \left\| \frac{v_{\theta,n}}{r} \right\|_{L^2(\Omega)}. \quad (3.20)$$

Therefore, (3.18) and (3.20) with the lower bound (2.6) imply

$$\begin{aligned} \|\partial_r v_{r,n}\|_{L^2(\Omega)}^2 + \|\partial_r v_{\theta,n}\|_{L^2(\Omega)}^2 + ((|n| - 1)^2 - |\alpha|) \left(\left\| \frac{v_{r,n}}{r} \right\|_{L^2(\Omega)}^2 + \left\| \frac{v_{\theta,n}}{r} \right\|_{L^2(\Omega)}^2 \right) \\ \leq |\Re \langle f_n, v_n \rangle_{L^2(\Omega)}|. \end{aligned} \quad (3.21)$$

Next we study identity (3.19). We see that

$$\begin{aligned} \Im \langle U^\perp \text{rot } v_n, v_n \rangle_{L^2(\Omega)} &= \Im \langle \frac{1}{|x|} \Delta \psi_n, v_{r,n} \rangle_{L^2(\Omega)} \\ &= -2\pi n \Re \int_1^\infty (\partial_r^2 \psi_n + \frac{1}{r} \partial_r \psi_n - \frac{n^2}{r^2} \psi_n) \overline{\psi_n} \frac{dr}{r}. \end{aligned}$$

Integrations by parts yield

$$\begin{aligned} &\Re \int_1^\infty (\partial_r^2 \psi_n + \frac{1}{r} \partial_r \psi_n - \frac{n^2}{r^2} \psi_n) \overline{\psi_n} \frac{dr}{r} \\ &= - \int_1^\infty |\partial_r \psi_n|^2 \frac{dr}{r} + 2\Re \int_1^\infty \frac{1}{r} \partial_r \psi_n \overline{\psi_n} \frac{dr}{r} - \int_1^\infty \frac{n^2 |\psi_n|^2}{r^2} \frac{dr}{r} \\ &= - \int_1^\infty |\partial_r \psi_n|^2 \frac{dr}{r} - (n^2 - 2) \int_1^\infty \frac{|\psi_n|^2}{r^2} \frac{dr}{r}. \end{aligned}$$

Hence,

$$\Im \langle U^\perp \text{rot } v_n, v_n \rangle_{L^2(\Omega)} = n \left(\left\| \frac{v_{\theta,n}}{r} \right\|_{L^2(\Omega)}^2 + \left(1 - \frac{2}{n^2}\right) \left\| \frac{v_{r,n}}{r} \right\|_{L^2(\Omega)}^2 \right). \quad (3.22)$$

Hence, (3.19) and (3.22) give

$$\begin{aligned} \alpha n \left(\left\| v_{r,n} \right\|_{L^2(\Omega)}^2 - \left(1 - \frac{2}{n^2}\right) \left\| \frac{v_{r,n}}{r} \right\|_{L^2(\Omega)}^2 + \|v_{\theta,n}\|_{L^2(\Omega)}^2 - \left\| \frac{v_{\theta,n}}{r} \right\|_{L^2(\Omega)}^2 \right) \\ = -\Im \langle f_n, v_n \rangle_{L^2(\Omega)}, \end{aligned} \quad (3.23)$$

which in particular leads to

$$\left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)}^2 + \left\| \frac{\sqrt{r^2-1}}{r} v_{\theta,n} \right\|_{L^2(\Omega)}^2 \leq \frac{1}{|\alpha n|} |\Im \langle f_n, v_n \rangle_{L^2(\Omega)}|, \quad (3.24)$$

$$\left\| \frac{v_{r,n}}{r} \right\|_{L^2(\Omega)}^2 \leq \left| \frac{n}{2\alpha} \right| |\Im \langle f_n, v_n \rangle_{L^2(\Omega)}|. \quad (3.25)$$

Then, from (3.23) and (3.25) we gather

$$\begin{aligned} \|v_{r,n}\|_{L^2(\Omega)}^2 &\leq \left\| \frac{v_{r,n}}{r} \right\|_{L^2(\Omega)}^2 + \frac{1}{|\alpha n|} |\Im \langle f_n, v_n \rangle_{L^2(\Omega)}| \\ &\leq \left(\left| \frac{n}{2\alpha} \right| + \frac{1}{|\alpha n|} \right) |\Im \langle f_n, v_n \rangle_{L^2(\Omega)}| \\ &\leq \left| \frac{3n}{2\alpha} \right| |\Im \langle f_n, v_n \rangle_{L^2(\Omega)}|, \quad |n| \geq 3, \end{aligned} \quad (3.26)$$

while for $|n| = 1, 2$, a slightly finer estimate is available from (3.23):

$$\|v_{r,n}\|_{L^2(\Omega)}^2 \leq \frac{1}{|\alpha n|} |\Im \langle f_n, v_n \rangle_{L^2(\Omega)}|, \quad |n| = 1, 2. \quad (3.27)$$

To obtain the estimate of $\|v_n\|_{L^2(\Omega)}$ we first observe that the following interpolation inequality holds for any scalar function $g \in W^{1,2}((1, \infty); r \, dr)$:

$$\|g\|_{L^2(\Omega)} \leq C \|\partial_r g\|_{L^2(\Omega)}^{\frac{1}{3}} \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}^{\frac{2}{3}} + C \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}. \quad (3.28)$$

See Appendix B for the proof of (3.28). From (3.28) and (3.24) we have

$$\begin{aligned} |\alpha| (\|v_{r,n}\|_{L^2(\Omega)}^2 + \|v_{\theta,n}\|_{L^2(\Omega)}^2) &\leq C |\alpha|^{\frac{3}{2}} \left(\left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)}^2 + \left\| \frac{\sqrt{r^2-1}}{r} v_{\theta,n} \right\|_{L^2(\Omega)}^2 \right) \\ &\quad + \frac{1}{4} (\|\partial_r v_{r,n}\|_{L^2(\Omega)}^2 + \|\partial_r v_{\theta,n}\|_{L^2(\Omega)}^2) \\ &\quad + C |\alpha| \left(\left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)}^2 + \left\| \frac{\sqrt{r^2-1}}{r} v_{\theta,n} \right\|_{L^2(\Omega)}^2 \right) \\ &\leq C \left(\frac{|\alpha|^{\frac{1}{2}}}{|n|} + \frac{1}{|n|} \right) \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \\ &\quad + \frac{1}{4} (\|\partial_r v_{r,n}\|_{L^2(\Omega)}^2 + \|\partial_r v_{\theta,n}\|_{L^2(\Omega)}^2), \end{aligned}$$

and thus,

$$\begin{aligned} |\alpha| (\|v_{r,n}\|_{L^2(\Omega)}^2 + \|v_{\theta,n}\|_{L^2(\Omega)}^2) \\ \leq \frac{C}{n^2} \left(1 + \frac{1}{|\alpha|} \right) \|f_n\|_{L^2(\Omega)}^2 + \frac{1}{2} (\|\partial_r v_{r,n}\|_{L^2(\Omega)}^2 + \|\partial_r v_{\theta,n}\|_{L^2(\Omega)}^2). \end{aligned} \quad (3.29)$$

Hence (3.21) and (3.29) imply

$$\|\partial_r v_{r,n}\|_{L^2(\Omega)}^2 + \|\partial_r v_{\theta,n}\|_{L^2(\Omega)}^2 \leq \frac{C}{n^2} \left(1 + \frac{1}{|\alpha|} \right) \|f_n\|_{L^2(\Omega)}^2 + 2 |\Re \langle f_n, v_n \rangle_{L^2(\Omega)}|. \quad (3.30)$$

Then (3.29) and (3.30) yield

$$\|v_{r,n}\|_{L^2(\Omega)}^2 + \|v_{\theta,n}\|_{L^2(\Omega)}^2 \leq \frac{C}{|\alpha| n^2} \left(1 + \frac{1}{|\alpha|} \right) \|f_n\|_{L^2(\Omega)}^2 + \frac{1}{|\alpha|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)},$$

that is,

$$\|v_n\|_{L^2(\Omega)}^2 \leq C \left(\frac{1}{|\alpha|n^2} + \frac{1}{|\alpha|^2} \right) \|f_n\|_{L^2(\Omega)}^2. \quad (3.31)$$

This proves (3.5) and (3.13). Note that the factor $\frac{1}{|\alpha|n^2}$ dominates $\frac{1}{\alpha^2}$ in the regime $|n| \leq O(|\alpha|^{\frac{1}{2}})$ and $|\alpha| \geq 1$. Once we have proved (3.31) the following estimates are immediately obtained from (3.24) and (3.30):

$$\left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)}^2 + \left\| \frac{\sqrt{r^2-1}}{r} v_{\theta,n} \right\|_{L^2(\Omega)}^2 \leq \frac{C}{|\alpha n|} \left(\frac{1}{|\alpha|^{\frac{1}{2}}|n|} + \frac{1}{|\alpha|} \right) \|f_n\|_{L^2(\Omega)}^2, \quad (3.32)$$

$$\|\partial_r v_{r,n}\|_{L^2(\Omega)}^2 + \|\partial_r v_{\theta,n}\|_{L^2(\Omega)}^2 \leq C \left(\frac{1}{n^2} + \frac{1}{|\alpha|} \right) \|f_n\|_{L^2(\Omega)}^2. \quad (3.33)$$

The constant C in (3.31), (3.32), and (3.33) is independent of $|n| \geq 1$ and α . Inequality in (3.32) proves (3.6). Moreover, (3.31) and (3.21) prove (3.8) and (3.15) since

$$|\alpha| \left\| \frac{v_n}{|x|} \right\|_{L^2(\Omega)}^2 \leq |\alpha| \|v_n\|_{L^2(\Omega)}^2.$$

Finally (3.7) and (3.14) are obtained from the interpolation inequality for functions in the space $W_0^{1,2}((1, \infty); dr)$.

3.1.3 A priori estimate of the n mode with $|n| \gg O(\sqrt{1+|\alpha|})$

If $|n| \geq 1 + \sqrt{2|\alpha|}$ then (3.21) yields

$$\begin{aligned} \|\partial_r v_{r,n}\|_{L^2(\Omega)}^2 + \|\partial_r v_{\theta,n}\|_{L^2(\Omega)}^2 + \frac{n^2}{8} \left(\left\| \frac{v_{r,n}}{r} \right\|_{L^2(\Omega)}^2 + \left\| \frac{v_{\theta,n}}{r} \right\|_{L^2(\Omega)}^2 \right) \\ \leq |\Re \langle f_n, v_n \rangle_{L^2(\Omega)}|. \end{aligned} \quad (3.34)$$

Then (3.23) and (3.34) give

$$\begin{aligned} \|v_n\|_{L^2(\Omega)}^2 &\leq \left\| \frac{v_n}{|x|} \right\|_{L^2(\Omega)}^2 + \frac{1}{|\alpha n|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \\ &\leq \left(\frac{8}{n^2} + \frac{1}{|\alpha n|} \right) \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)}, \end{aligned}$$

which shows

$$\|v_n\|_{L^2(\Omega)}^2 \leq \left(\frac{8}{n^2} + \frac{1}{|\alpha n|} \right)^2 \|f_n\|_{L^2(\Omega)}^2. \quad (3.35)$$

Note that (3.35) is better than (3.31) in the regime $|n| \gg 1 + \sqrt{2|\alpha|}$, while both are of the same order in the regime $|n| = O(\sqrt{1+|\alpha|})$. The estimates (3.34) and (3.35) yield

$$\|\nabla v_n\|_{L^2(\Omega)}^2 \leq C \left(\frac{1}{n^2} + \frac{1}{|\alpha n|} \right) \|f_n\|_{L^2(\Omega)}^2, \quad (3.36)$$

while (3.24) and (3.35) lead to

$$\left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)}^2 + \left\| \frac{\sqrt{r^2-1}}{r} v_{\theta,n} \right\|_{L^2(\Omega)}^2 \leq \frac{1}{|\alpha n|} \left(\frac{8}{n^2} + \frac{1}{|\alpha n|} \right) \|f_n\|_{L^2(\Omega)}^2. \quad (3.37)$$

Again, (3.36) and (3.37) are better than (3.33) and (3.32) in the regime $|n| \gg 1 + \sqrt{2|\alpha|}$. The estimates (3.35), (3.36), and (3.37) show (3.9), (3.12), and (3.10). Then (3.11) follows by interpolation using (3.9) and (3.12)

3.1.4 Proof of Proposition 3.1

The statement (i) of Proposition 3.1 is proved in Subsection 3.1.1. In particular, we have that $v_0 = v_{\theta,0} \mathbf{e}_\theta$ and $v_{\theta,0}$ is given by the formula (3.17). It remains to prove (ii) of Proposition 3.1.

(Estimates and Uniqueness) We have already proved the a priori estimates of solutions in Subsections 3.1.2 and 3.1.3, which give (3.5)-(3.12). The uniqueness of solutions directly follows from these a priori estimates.

(Existence) By considering the Helmholtz-Leray projection which is bounded in L^2 and invariant under the action of \mathcal{P}_n , we may assume that f_n belongs to $\mathcal{Q}_0 L_\sigma^2(\Omega)$, rather than $\mathcal{Q}_0 L^2(\Omega)^2$. To show the existence of solutions we consider the operator

$$\mathbb{A}_\alpha := \mathbb{P}\Delta - \alpha \mathbb{P}U^\perp \text{rot}$$

in the L^2 framework, where $\mathbb{P} : L^2(\Omega)^2 \rightarrow L_\sigma^2(\Omega)$ is the Helmholtz-Leray projection and Δ is the Dirichlet Laplacian in $L^2(\Omega)$. The operator $\mathbb{P}\Delta$ is thus the standard Stokes operator in $L_\sigma^2(\Omega)$. Let us consider the operator \mathbb{A}_α in the invariant space $\mathcal{P}_n L_\sigma^2(\Omega)$, $n \neq 0$. Note that the spectrum of the Stokes operator $\mathbb{P}\Delta$ in $\mathcal{P}_n L_\sigma^2(\Omega)$ is included in the half real line $\overline{\mathbb{R}}_- = \{\lambda \leq 0\}$, while the operator $\mathbb{P}U^\perp \text{rot}$ is relatively compact with respect to $\mathbb{P}\Delta$ in $\mathcal{P}_n L_\sigma^2(\Omega)$, for U^\perp is smooth and decays at infinity. Thus, the difference between the spectrum of \mathbb{A}_α and the one of $\mathbb{P}\Delta$ consists only of discrete eigenvalues with finite multiplicities. Then the scalar $\lambda := -i\alpha n$ must belong to the resolvent set of \mathbb{A}_α in $\mathcal{P}_n L_\sigma^2(\Omega)$, otherwise $-i\alpha n$ is an eigenvalue of \mathbb{A}_α in $\mathcal{P}_n L_\sigma^2(\Omega)$ but this cannot be true due to the a priori estimates on solutions of (3.3) with $\lambda = -i\alpha n$ which we have shown above. Hence, when $\lambda = -i\alpha n$, for any $f_n = \mathcal{P}_n f \in \mathcal{P}_n L_\sigma^2(\Omega)$ there exists a unique solution v_n to (3.3) belonging to the space $\mathcal{P}_n L_\sigma^2(\Omega) \cap W_0^{1,2}(\Omega)^2 \cap W^{2,2}(\Omega)^2$ with a suitable pressure field belonging to $W_{loc}^{1,2}(\overline{\Omega})$. Then $v = \sum_{n \neq 0} v_n$ belongs to $\mathcal{Q}_0 L_\sigma^2(\Omega) \cap W_0^{1,2}(\Omega)^2 \cap W^{2,2}(\Omega)^2$ and solves (S_α) by construction, for (3.3) with $\lambda = -i\alpha n$ is equivalent to (3.1) and (3.2) for each $n \neq 0$. The proof of (ii) of Proposition 3.1 is complete. \square

3.2 Analysis in the fast rotation case $|\alpha| \gg 1$

In this subsection we focus on the behavior of solutions to (S_α) in the case $|\alpha| \gg 1$. Let us define the parameter

$$\beta := (-2i\alpha n)^{\frac{1}{3}} = \begin{cases} (2|\alpha n|)^{\frac{1}{3}} c_- & \text{if } \alpha n > 0, \\ (2|\alpha n|)^{\frac{1}{3}} c_+ & \text{if } \alpha n < 0, \end{cases} \quad (3.38)$$

where

$$c_\pm := \frac{\sqrt{3} \pm i}{2}. \quad (3.39)$$

Our goal is to prove the following structure result on solutions to (S_α) .

Proposition 3.2 *There is a constant $\kappa \in (0, 1)$ such that as long as $1 \leq |n| \leq \kappa |\alpha|^{\frac{1}{2}}$ and $|\beta|$ large enough (implying $|\alpha|$ large enough), the n mode of the velocity field v_n solving (S_α) may be decomposed into four parts*

$$v_n = v_n^{\text{slip}} + v_n^{\text{slow}} + v_{n,\text{BL}} + \tilde{v}_n,$$

where

- the term v_n^{slip} satisfies the system (mS_α) and the estimates (3.46)-(3.48) of Proposition 3.4.
- the stream function of v_n^{slow} is given by

$$\psi_n^{\text{slow}}(r) = a_n r^{-|n|},$$

where

$$|a_n| \leq C|\alpha n|^{-\frac{5}{6}} \|f_n\|_{L^2}.$$

- the boundary layer term $v_{n,\text{BL}}$ is given by

$$v_{n,\text{BL},r}(r) = \frac{inb_n}{r} G_{n,\alpha}(|\beta|(r-1)), \quad v_{n,\text{BL},\theta} = -|\beta|b_n G'_{n,\alpha}(|\beta|(r-1))$$

with $G_{n,\alpha}$ a smooth function, decaying exponentially at infinity, uniformly in n and α , and where

$$|b_n| \leq C|\alpha n|^{-\frac{5}{6}} \|f_n\|_{L^2}.$$

- the term \tilde{v}_n is a remainder term in the sense that

$$\begin{aligned} \|\tilde{v}_n\|_{L^2(\Omega)} &\leq C|\alpha n|^{-1} \|f_n\|_{L^2}, \quad \|\tilde{v}_n\|_{L^\infty(\Omega)} \leq C|\alpha n|^{-\frac{5}{6}} \|f_n\|_{L^2}, \\ \text{and} \quad \|\nabla \tilde{v}_n\|_{L^2(\Omega)} &\leq C|\alpha n|^{-\frac{2}{3}} \|f_n\|_{L^2}. \end{aligned} \quad (3.40)$$

In particular, the following estimates hold for v_n :

$$\begin{aligned} \|v_n\|_{L^2(\Omega)} &\leq C|\alpha n|^{-\frac{2}{3}} \|f_n\|_{L^2}, \quad \|v_n\|_{L^\infty(\Omega)} \leq C|\alpha n|^{-\frac{1}{2}} \|f_n\|_{L^2}, \\ \text{and} \quad \|\nabla v_n\|_{L^2(\Omega)} &\leq C|\alpha n|^{-\frac{1}{3}} \|f_n\|_{L^2}. \end{aligned} \quad (3.41)$$

The constant C is independent of n , α , and f_n .

Remark 3.3 (1) The velocity fields v_n^{slip} , v_n^{slow} and $v_{n,\text{BL}}$ have the same decay order in the spaces L^2 , L^∞ and H^1 , although their structure is different. In particular contrary to the other velocity fields, the term $v_{n,\text{BL}}$ is negligible away from the boundary due to its rapid decay. The key point is that the boundary layer analysis enables us to improve the decay order in the low and middle frequencies $|n| \ll O(|\alpha|^{\frac{1}{2}})$, compared with the results of Proposition 3.1 which are based only on energy computations. Indeed, it is not clear whether (3.41) can be shown only from energy computations without using the boundary layer analysis.

(2) The vorticity of v_n^{slow} vanishes in Ω , that is, v_n^{slow} is irrotational in Ω . The function $G_{n,\alpha}$ and its derivatives of finite order are uniformly bounded in n and α which satisfy $|\alpha| \geq 1$ and $1 \leq |n| \leq |\alpha|^{\frac{1}{2}}$. The uniform decay estimate of $G_{n,\alpha}(\rho)$ for $\rho \gg 1$ is stated in (3.80) below. The smallness of κ in Proposition 3.2 is required only in obtaining some lower bound for the quantity $\left| |\beta|G'_{n,\alpha}(0) + |n|G_{n,\alpha}(0) \right|$ which is essential to construct the solution satisfying the noslip boundary condition and possessing the structure stated in Proposition 3.2.

Proof of Proposition 3.2: We first consider a linearized problem similar to (S_α) but with a different boundary condition, such that the vorticity vanishes on the boundary:

$$\begin{cases} -\Delta v - \alpha(x^\perp \cdot \nabla v - v^\perp) + \nabla q + \alpha U^\perp \text{rot } v = f, & x \in \Omega, \\ \text{div } v = 0, & x \in \Omega, \\ v_r = \text{rot } v = 0, & x \in \partial\Omega. \end{cases} \quad (\text{mS}_\alpha)$$

Note that the vorticity field ω then satisfies the following equations

$$\begin{cases} -\Delta \omega - \alpha x^\perp \cdot \nabla \omega + \alpha U \cdot \nabla \omega = \text{rot } f, & x \in \Omega, \\ \omega|_{\partial\Omega} = 0. \end{cases} \quad (3.42)$$

Let us prove the following proposition.

Proposition 3.4 *For any $\alpha \in \mathbb{R}$ with $|\alpha| \geq 1$ and external force $f \in \mathcal{Q}_0 L^2(\Omega)^2$ the system (mS_α) admits a unique solution $(v, \nabla q)$ with $v \in \mathcal{Q}_0 L^2_\sigma(\Omega) \cap W^{1,2}(\Omega)^2 \cap W^{2,2}_{loc}(\overline{\Omega})$ and $q \in W^{1,2}_{loc}(\overline{\Omega})$. Moreover, $\omega = \text{rot } v \in \mathcal{Q}_0 W_0^{1,2}(\Omega)$ satisfies for $n \neq 0$,*

$$\|\mathcal{P}_n \omega\|_{L^2(\Omega)} \leq \frac{C}{|\alpha n|^{\frac{1}{3}}} \|f_n\|_{L^2(\Omega)}, \quad (3.43)$$

$$\left\| \frac{\sqrt{|x|^2 - 1}}{|x|} \mathcal{P}_n \omega \right\|_{L^2(\Omega)} \leq \frac{1}{|\alpha n|^{\frac{1}{2}}} \|f_n\|_{L^2(\Omega)}, \quad (3.44)$$

$$\|\nabla \mathcal{P}_n \omega\|_{L^2(\Omega)} \leq \|f_n\|_{L^2(\Omega)}, \quad (3.45)$$

while $v_n = \mathcal{P}_n v$ satisfies for $1 \leq |n| \leq 2|\alpha|^{\frac{1}{2}}$,

$$\|v_n\|_{L^2(\Omega)} \leq \frac{C}{|\alpha n|^{\frac{2}{3}}} \|f_n\|_{L^2(\Omega)}, \quad (3.46)$$

$$\|v_n\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha n|^{\frac{1}{2}}} \|f_n\|_{L^2(\Omega)}, \quad (3.47)$$

$$\left\| \frac{\sqrt{|x|^2 - 1}}{|x|} v_n \right\|_{L^2(\Omega)} \leq \frac{C}{|\alpha n|^{\frac{5}{6}}} \|f_n\|_{L^2(\Omega)}. \quad (3.48)$$

Here $f_n := \mathcal{P}_n f$ and C is independent of n and α .

Remark 3.5 Note that

$$\|\nabla v_n\|_{L^2(\Omega)} \leq C \left(\|\mathcal{P}_n \omega\|_{L^2(\Omega)} + \left\| \frac{v_n}{|x|} \right\|_{L^2(\Omega)} \right) \leq \frac{C}{|\alpha n|^{\frac{1}{3}}} \|f_n\|_{L^2(\Omega)}. \quad (3.49)$$

The decay order in each estimate of (3.49) and (3.46)-(3.48) is better than the order in (3.5)-(3.8) for the solution subject to the noslip boundary condition. This faster decay is shown below by an energy estimate thanks to the slip boundary condition in (mS_α) , which reduces the magnitude of the boundary layer arising from the fast rotation in low and middle frequencies.

Proof of Proposition 3.4: For simplicity we set $\omega_n = \omega_n(r) := (\mathcal{P}_n \omega) e^{-in\theta}$.

(A priori estimates) We first show the a priori estimates stated in (3.43)-(3.48). As in the proof of Proposition 3.1, for each n mode the energy computation for v_n based on the integration by parts is verified for any solution $(v, \nabla q)$ to (mS_α) such that $v \in \mathcal{Q}_0 L_\sigma^2(\Omega) \cap W^{1,2}(\Omega)^2 \cap W_{loc}^{2,2}(\bar{\Omega})$ and $q \in W_{loc}^{1,2}(\bar{\Omega})$ (see the argument at the beginning of Subsection 3.1). We also note that the n mode of the vorticity $\omega_n \in W_0^{1,2}((1, \infty); r dr)$ satisfies the following ordinary differential equations on $(1, \infty)$ in the weak sense:

$$-\left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{n^2}{r^2} + i\alpha n \left(1 - \frac{1}{r^2}\right)\right) \omega_n = \frac{1}{r} \frac{d}{dr} (r f_{\theta,n}) - \frac{in}{r} f_{r,n}, \quad (3.50)$$

with the Dirichlet boundary condition $\omega_n(1) = 0$. We can multiply both sides by $r\bar{\omega}_n$ and integrate over $[1, \infty)$, which gives the following identities:

$$\int_1^\infty \left(|\partial_r \omega_n|^2 + n^2 \frac{|\omega_n|^2}{r^2} \right) r dr = -\Re \left(\int_1^\infty f_{\theta,n} \frac{d\bar{\omega}_n}{dr} r dr + in \int_1^\infty f_{r,n} \bar{\omega}_n dr \right), \quad (3.51)$$

$$\alpha n \int_1^\infty \left(1 - \frac{1}{r^2}\right) |\omega_n|^2 r dr = \Im \left(\int_1^\infty f_{\theta,n} \frac{d\bar{\omega}_n}{dr} r dr + in \int_1^\infty f_{r,n} \bar{\omega}_n dr \right). \quad (3.52)$$

The first identity (3.51) gives the bound

$$\|\nabla \mathcal{P}_n \omega\|_{L^2(\Omega)} \leq \|f_n\|_{L^2(\Omega)},$$

and then the second identity (3.52) yields

$$\left\| \frac{\sqrt{|x|^2 - 1}}{|x|} \mathcal{P}_n \omega \right\|_{L^2(\Omega)}^2 \leq \frac{1}{|\alpha n|} \|f_n\|_{L^2(\Omega)} \|\nabla \mathcal{P}_n \omega\|_{L^2(\Omega)} \leq \frac{1}{|\alpha n|} \|f_n\|_{L^2(\Omega)}^2.$$

Thus (3.44) and (3.45) hold. The estimate (3.43) follows from (3.44) and (3.45) by the interpolation inequality (3.28) and $|\alpha| \geq 1$. The estimates on the velocity v_n are obtained from the estimates for the streamfunction ψ_n which is the solution to the Poisson equation (2.8).

Let us first estimate $\frac{d^2 \psi_n}{dr^2}$ and $\frac{\psi_n}{r^2}$. We write $\psi'_n = \frac{d\psi_n}{dr}$ and $\psi''_n = \frac{d^2 \psi_n}{dr^2}$ for simplicity. We observe that

$$\int_1^\infty \frac{1}{r} \psi'_n \bar{\psi}''_n r dr = \int_1^\infty \left(\frac{\psi_n}{r} \right)' \bar{\psi}''_n r dr + \int_1^\infty \frac{\psi_n}{r^2} \bar{\psi}''_n r dr.$$

A similar computation yields

$$-\Re \int_1^\infty \frac{\psi_n}{r^2} \bar{\psi}''_n r dr = \Re \int_1^\infty \left(\frac{\psi_n}{r} \right)' \bar{\psi}'_n r dr = \int_1^\infty \left| \left(\frac{\psi_n}{r} \right)' \right|^2 r dr$$

and

$$\Re \int_1^\infty \frac{1}{r} \psi'_n \frac{\bar{\psi}_n}{r^2} r dr = \int_1^\infty \left| \frac{\psi_n}{r^2} \right|^2 r dr.$$

Hence we have from (2.8), by multiplying both sides by $r\bar{\psi}''_n$ and by integrating over $[1, \infty)$,

$$\int_1^\infty |\psi''_n|^2 r dr + \Re \int_1^\infty \left(\frac{\psi_n}{r} \right)' \bar{\psi}''_n r dr + (n^2 - 1) \int_1^\infty \left| \left(\frac{\psi_n}{r} \right)' \right|^2 r dr = -\Re \int_1^\infty \omega_n \bar{\psi}''_n r dr,$$

and similarly, by multiplying by $\frac{\bar{\psi}_n}{r}$ in (2.8) and integrating over $[1, \infty)$,

$$\int_1^\infty \left| \left(\frac{\psi_n}{r} \right)' \right|^2 r dr + (n^2 - 1) \int_1^\infty \left| \frac{\psi_n}{r^2} \right|^2 r dr = \Re \int_1^\infty \omega_n \frac{\bar{\psi}_n}{r^2} r dr.$$

Combining these two identities gives the following bounds: when $|n| \geq 2$,

$$\|\psi_n''\|_{L^2(\Omega)} + \left\| \left(\frac{n\psi_n}{r} \right)' \right\|_{L^2(\Omega)} + n^2 \left\| \frac{\psi_n}{r^2} \right\|_{L^2(\Omega)} \leq C \|\omega_n\|_{L^2(\Omega)}, \quad (3.53)$$

while when $|n| = 1$,

$$\begin{aligned} \left\| \left(\frac{\psi_n}{r} \right)' \right\|_{L^2(\Omega)}^2 &\leq \|\omega_n\|_{L^2(\Omega)} \left\| \frac{\psi_n}{r^2} \right\|_{L^2(\Omega)}, \\ \|\psi_n''\|_{L^2(\Omega)}^2 &\leq C \|\omega_n\|_{L^2(\Omega)} \left(\|\omega_n\|_{L^2(\Omega)} + \left\| \frac{\psi_n}{r^2} \right\|_{L^2(\Omega)} \right), \end{aligned} \quad (3.54)$$

where C is independent of n . Next we observe that the identity (3.23) holds even under the slip boundary condition, and thus, (3.24) and (3.25) are valid also for solutions to (mS_α) . Hence, by recalling the relation $v_{r,n} = \frac{in}{r}\psi_n$ and using the interpolation inequality (3.28), we obtain for $|n| \geq 2$, thanks to (3.53) and (3.24),

$$\begin{aligned} \|v_{r,n}\|_{L^2(\Omega)} &\leq C \|\partial_r v_{r,n}\|_{L^2(\Omega)}^{\frac{1}{3}} \left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)}^{\frac{2}{3}} + C \left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)} \\ &\leq C \|\omega_n\|_{L^2(\Omega)}^{\frac{1}{3}} \left(\frac{1}{|\alpha n|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{1}{3}} + C \left(\frac{1}{|\alpha n|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{1}{2}}, \end{aligned}$$

while for $|n| = 1$ we use (3.54) and also (3.25), which give

$$\begin{aligned} \|v_{r,n}\|_{L^2(\Omega)} &\leq C \|\partial_r v_{r,n}\|_{L^2(\Omega)}^{\frac{1}{3}} \left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)}^{\frac{2}{3}} + C \left\| \frac{\sqrt{r^2-1}}{r} v_{r,n} \right\|_{L^2(\Omega)} \\ &\leq C \|\omega_n\|_{L^2(\Omega)}^{\frac{1}{6}} \left(\frac{1}{|\alpha|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{5}{12}} + C \left(\frac{1}{|\alpha|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{1}{2}}. \end{aligned}$$

Similarly, we have from $v_{\theta,n} = -\psi_n'$, (3.28), (3.53), and (3.24), for $|n| \geq 2$,

$$\|v_{\theta,n}\|_{L^2(\Omega)} \leq C \|\omega_n\|_{L^2(\Omega)}^{\frac{1}{3}} \left(\frac{1}{|\alpha n|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{1}{3}} + C \left(\frac{1}{|\alpha n|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{1}{2}},$$

on the other hand, for $|n| = 1$, by applying (3.28), (3.54), (3.24) and (3.25),

$$\begin{aligned} \|v_{\theta,n}\|_{L^2(\Omega)} &\leq C \|\omega_n\|_{L^2(\Omega)}^{\frac{1}{3}} \left(\frac{1}{|\alpha|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{1}{3}} \\ &\quad + C \|\omega_n\|_{L^2(\Omega)}^{\frac{1}{6}} \left(\frac{1}{|\alpha|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{5}{12}} + C \left(\frac{1}{|\alpha|} \|f_n\|_{L^2(\Omega)} \|v_n\|_{L^2(\Omega)} \right)^{\frac{1}{2}}. \end{aligned}$$

Hence we have from (3.43) for the estimate of $\|\omega_n\|_{L^2(\Omega)}$ and $|\alpha| \geq 1$,

$$\|v_n\|_{L^2(\Omega)} \leq \frac{C}{|\alpha n|^{\frac{2}{3}}} \|f_n\|_{L^2(\Omega)}, \quad |n| \geq 1. \quad (3.55)$$

The estimate (3.48) follows from (3.24) and (3.55). As for the L^∞ estimate in the case $|n| \geq 2$, we have from the interpolation inequality and (3.53),

$$\begin{aligned} \|v_{r,n}\|_{L^\infty(\Omega)}^2 &\leq C \|\partial_r v_{r,n}\|_{L^2(\Omega)} \|v_{r,n}\|_{L^2(\Omega)} \leq C \|\omega_n\|_{L^2(\Omega)} \|v_{r,n}\|_{L^2(\Omega)} \\ &\leq \frac{C}{|\alpha n|} \|f_n\|_{L^2(\Omega)}^2, \end{aligned}$$

and similarly,

$$\|v_{\theta,n}\|_{L^\infty(\Omega)}^2 \leq \frac{C}{|\alpha n|} \|f_n\|_{L^2(\Omega)}^2.$$

The case $|n| = 1$ is handled in the same manner, and in this case we use (3.54) instead of (3.53) to estimate $\|\partial_r v_{r,n}\|_{L^2(\Omega)}$ and $\|\partial_r v_{\theta,n}\|_{L^2(\Omega)}$. This modification does not produce any change in the final estimate

$$\|v_n\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|f_n\|_{L^2(\Omega)}.$$

The details are omitted here. Thus (3.43)-(3.48) hold.

(Existence and uniqueness) The uniqueness follows from the a priori estimates. The existence is shown by the same argument as in the proof of Proposition 3.1 in Subsection 3.1.4 (the proof for the statement (ii)), so we omit the details here. The proof of Proposition 3.4 is complete. \square

Let us return to the proof of Proposition 3.2. We denote by v^{slip} the solution to (mS_α) with the external force f in Proposition 3.4. Note that v^{slip} satisfies the desired estimate (3.41), while v^{slip} is not necessarily subject to the noslip boundary condition. Hence, starting from the perfect-slip solution v^{slip} , our next task is to recover the noslip boundary condition by the boundary layer analysis. To this end we consider the following equations related to the stream function:

$$\begin{cases} \left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{n^2}{r^2} + i\alpha n \left(1 - \frac{1}{r^2}\right) \right) \left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{n^2}{r^2} \right) \varphi_n = 0, & r > 1, \\ |\varphi_n(1)| \leq 1, & \lim_{r \rightarrow \infty} \varphi_n = 0. \end{cases} \quad (3.56)$$

Indeed, the first equation in (3.56) is nothing but the equation for the n mode of the stream function in our problem (in polar coordinates), while here we do not need to impose the exact boundary value on the boundary $r = 1$, and the key point is that there exists a solution to (3.56) which possess a boundary layer structure due to $|\alpha n| \gg 1$. We shall decompose it into a boundary layer part and a remainder:

$$\varphi_n = \varphi_{n,\text{BL}} + \tilde{\varphi}_n, \quad (3.57)$$

where $\varphi_{n,\text{BL}}$ is a function on the boundary layer variable $|\beta|(r-1)$ and $\tilde{\varphi}_n$ vanishes at $r = 1$ and is smaller than $\varphi_{n,\text{BL}}$ up to its derivatives. The precise construction of $\varphi_{n,\text{BL}}$ and $\tilde{\varphi}_n$ will be stated later.

Once φ_n is constructed as in (3.57), the n mode of the noslip solution v_n to (3.56), or equivalently its stream function ψ_n , is obtained by the following argument. Noticing that (3.56) has an explicit solution given by $r \mapsto r^{-|n|}$ (corresponding to a vorticity-free solution), we look for the stream function ψ_n under the following form:

$$\psi_n(r) = a_n r^{-|n|} + b_n \varphi_n(r) + \psi_n^{\text{slip}}(r) \quad (3.58)$$

where ψ_n^{slip} is the n mode of the streamfunction for v_n^{slip} , namely,

$$\psi_n^{\text{slip}} = \frac{r}{in} v_{r,n}^{\text{slip}}, \quad n \neq 0, \quad (3.59)$$

and the coefficients a_n and b_n are determined from the prescription that ψ_n and ψ'_n vanish at the boundary $r = 1$:

$$\begin{aligned} a_n + b_n \varphi_n(1) &= 0 \\ -|n|a_n + b_n \frac{d\varphi_n}{dr}(1) &= -\frac{d\psi_n^{\text{slip}}}{dr}(1) \quad (= v_{\theta,n}^{\text{slip}}(1)). \end{aligned}$$

In other words there holds

$$b_n \left(\frac{d\varphi_n}{dr}(1) + |n|\varphi_n(1) \right) = v_{\theta,n}^{\text{slip}}(1), \quad a_n = -b_n \varphi_n(1). \quad (3.60)$$

The key point is that for $|n| \leq \kappa|\alpha|^{\frac{1}{2}}$ for some small enough κ , the term $\frac{d\varphi_n}{dr}(1) + |n|\varphi_n(1)$ will be shown to be nonzero and actually large, due to a specific boundary layer structure of φ_n . Combining (3.57) with (3.58) gives the formula

$$\psi_n(r) = \psi_n^{\text{slip}}(r) + a_n r^{-|n|} + b_n \varphi_{n,\text{BL}}(r) + b_n \tilde{\varphi}_n(r).$$

Thus, with the notation of Proposition 3.2, the remainder velocity \tilde{v}_n is given in terms of the stream function $\tilde{\psi}_n$ defined as

$$\tilde{\psi}_n(r) := b_n \tilde{\varphi}_n(r).$$

We now focus on the construction of φ_n and its associate velocity field. To estimate a possible boundary layer thickness we observe from the first equation of (3.56) that there is a natural scale balance between $-\frac{d^2}{dr^2}$ and $-i\alpha n(1 - \frac{1}{r^2}) \approx -2i\alpha n(r - 1)$ near the boundary $r = 1$, which formally implies that the thickness of the boundary layer is $|2\alpha n|^{-\frac{1}{3}} = |\beta|^{-1}$ where β is defined in (3.38). Before stating the result leading to the construction of the boundary layer term, let us recall the notation introduced in (2.8):

$$H_n := -\frac{d^2}{dr^2} - \frac{1}{r} \frac{d}{dr} + \frac{n^2}{r^2}, \quad (3.61)$$

and let us denote

$$A_n := H_n - i\alpha n \left(1 - \frac{1}{r^2} \right) \quad (3.62)$$

so that the first equation in (3.56) translates into $A_n H_n \varphi_n = 0$. The next proposition is the construction of $\varphi_{n,\text{BL}}$ in (3.57), which describes the leading part of the boundary layer.

Proposition 3.6 *There exist $\kappa \in (0, 1)$ and $C > 0$ such that the following statement holds. If $|n| \leq \kappa|\alpha|^{\frac{1}{2}}$ and if $|\beta| = (2|\alpha n|)^{\frac{1}{3}}$ is large enough, then there exist smooth functions $\varphi_{n,\text{BL}}$ and $g_{n,\text{BL}}$ on $[1, \infty)$ such that*

$$A_n H_n \varphi_{n,\text{BL}} = \frac{1}{r} \frac{d}{dr} (r g_{n,\text{BL}}), \quad \text{with} \quad \|g_{n,\text{BL}}\|_{L^2(\Omega)} \leq C|\beta|^{\frac{3}{2}}, |\varphi_{n,\text{BL}}(1)| \leq 1, \quad (3.63)$$

and

- there holds

$$\left| \frac{d\varphi_{n,\text{BL}}}{dr}(1) + |n|\varphi_{n,\text{BL}}(1) \right| \geq \frac{\kappa}{2} |\beta|, \quad (3.64)$$

- there is a smooth function $G_{n,\alpha}$ decaying exponentially at infinity uniformly in n and α such that

$$\varphi_{n,\text{BL}}(r) = G_{n,\alpha}(|\beta|(r-1)), \quad r \geq 1. \quad (3.65)$$

More precisely, $G_{n,\alpha}$ in (3.65) satisfies the estimate (3.80) stated below. Let us postpone the proof of Proposition 3.6 and conclude the proof of Proposition 3.2. We construct a couple $(\varphi_{n,\text{BL}}, g_{n,\text{BL}})$ as in Proposition 3.6, which produces a boundary layer vector field

$$v_{r,n,\text{BL}}(r) := \frac{inb_n}{r} G_{n,\alpha}(|\beta|(r-1)), \quad v_{\theta,n,\text{BL}}(r) := -|\beta|b_n G'_{n,\alpha}(|\beta|(r-1)). \quad (3.66)$$

Next we fix $\varphi_n(1) = \varphi_{n,\text{BL}}(1)$, so that the remainder $\tilde{\varphi}_n$ as in (3.57) vanishes at the boundary. Moreover, from the requirement $A_n H_n \varphi_n = 0$ and thanks to (3.63), $\tilde{\varphi}_n$ is obtained as the solution to

$$A_n H_n \tilde{\varphi}_n = -\frac{1}{r} \frac{d}{dr}(r g_{n,\text{BL}}) \quad r > 1, \quad (H_n \tilde{\varphi}_n)(1) = \tilde{\varphi}_n(1) = 0.$$

More precisely, the construction of $\tilde{\varphi}_n$ is as follows: we first construct $\tilde{\omega}_n(r)e^{in\theta}$ as the solution to (3.42) with right-hand side $-\text{rot}(g_{n,\text{BL}}e^{in\theta}\mathbf{e}_\theta)$. Then $\tilde{\varphi}_n$ is obtained as the solution to $H_n \tilde{\varphi}_n = \tilde{\omega}_n$ under the boundary condition $\tilde{\varphi}_n(1) = 0$. Let us denote by \tilde{w}_n the velocity field whose stream function is $\tilde{\varphi}_n$. Then the estimates of \tilde{w}_n and $\tilde{\omega}_n$ follow from Proposition 3.4 and the fact that $\|g_{n,\text{BL}}\|_{L^2(\Omega)} \leq C|\beta|^{\frac{3}{2}}$ as stated in (3.63). In particular, (3.47) in Proposition 3.4 produces from $|\beta| = |2\alpha n|^{\frac{1}{3}}$ that

$$\left| \frac{d\tilde{\varphi}_n}{dr}(1) \right| \leq \|\tilde{w}_{\theta,n}\|_{L^\infty(\Omega)} \lesssim |\beta|^{-\frac{3}{2}} \|g_{n,\text{BL}}\|_{L^2(\Omega)} \leq C.$$

Together with (3.57) and (3.64), we find that

$$\begin{aligned} \left| \frac{d\varphi_n}{dr}(1) + |n|\varphi_n(1) \right| &\geq \left| \frac{d\varphi_{n,\text{BL}}}{dr}(1) + |n|\varphi_{n,\text{BL}}(1) \right| - \left| \frac{d\tilde{\varphi}_n}{dr}(1) \right| \\ &\geq \frac{\kappa}{2} |\beta| - C \\ &\geq \frac{\kappa}{4} |\beta| \end{aligned}$$

for $|\beta| \gg \kappa^{-1}$. Let us estimate the coefficients a_n and b_n , which are defined in (3.60). Since

$$|v_{\theta,n}^{\text{slip}}(1)| \leq \|v_{\theta,n}^{\text{slip}}\|_{L^\infty(\Omega)} \lesssim |\alpha n|^{-\frac{1}{2}} \|f_n\|_{L^2(\Omega)}$$

thanks to Proposition 3.4 we infer that the parameter b_n satisfies

$$|b_n| = \left| \frac{v_{\theta,n}^{\text{slip}}(1)}{\frac{d\varphi_n}{dr}(1) + |n|\varphi_n(1)} \right| \lesssim |\alpha n|^{-\frac{5}{6}} \|f_n\|_{L^2(\Omega)}$$

while since $|\varphi_n(1)| \leq 1$,

$$|a_n| = |b_n \varphi_n(1)| \lesssim |\alpha n|^{-\frac{5}{6}} \|f_n\|_{L^2(\Omega)}.$$

Thus, the estimates of the boundary layer velocity $v_{n,\text{BL}}$ easily follow from its definition in (3.66). The estimates of the remainder velocity \tilde{v}_n follow from Proposition 3.4 and the

estimates of b_n since $\tilde{\psi}_n = b_n \tilde{\varphi}_n$ that implies $\tilde{v}_n = b_n \tilde{w}_n$. This concludes the proof of Proposition 3.2. \square

Proof of Proposition 3.6: Without loss of generality we assume from now on that $\alpha > 0$. As already mentioned, we formally estimate the thickness of the boundary layer to be of the order $|\beta| = |2\alpha n|^{\frac{1}{3}}$. One important remark here is the size of $\frac{n^2}{r^2}$ in the operators H_n and A_n defined in (3.61) and (3.62). Recall that we are interested in the regime $|n| \leq O(\alpha^{\frac{1}{2}})$. If $|n| = O(\alpha^{\frac{1}{2}})$ then we observe that $|\beta| = |2\alpha n|^{\frac{1}{3}} = O((\alpha^{\frac{3}{2}})^{\frac{1}{3}}) = O(|n|)$, and thus, the term $\frac{n^2}{r^2}$ has the same size near the boundary as ∂_r^2 and $\alpha n(r-1)$. Hence, in the construction of the boundary layer we also need to take into account the term $\frac{n^2}{r^2}$, for this term is no longer small in the regime $|n| = O(\alpha^{\frac{1}{2}})$. With this remark in mind let us rewrite H_n and A_n in more convenient forms: we define

$$\tilde{H}_n = -\frac{d^2}{dr^2} + n^2 \quad \text{and} \quad \tilde{A}_n = \tilde{H}_n - 2i\alpha n(r-1) = -\frac{d^2}{dr^2} + \beta^3 \left(r-1 + \frac{in}{2\alpha} \right),$$

so that

$$H_n = \tilde{H}_n - \frac{1}{r} \frac{d}{dr} + n^2 \left(\frac{1}{r^2} - 1 \right)$$

and by using $1 - \frac{1}{r^2} = 2(r-1) + (r-1)^2 \frac{1+2r}{r^2}$,

$$\begin{aligned} A_n &= H_n - 2i\alpha n(r-1) - i\alpha n(r-1)^2 \frac{1+2r}{r^2} \\ &= \tilde{H}_n - 2i\alpha n(r-1) - \frac{1}{r} \frac{d}{dr} + n^2 \left(\frac{1}{r^2} - 1 \right) - i\alpha n(r-1)^2 \frac{1+2r}{r^2} \\ &= \tilde{A}_n - \frac{1}{r} \frac{d}{dr} + n^2 \left(\frac{1}{r^2} - 1 \right) - 2i\alpha n(r-1)^2 \frac{1+2r}{r^2}. \end{aligned}$$

That is, \tilde{H}_n and \tilde{A}_n have the leading size of H_n and A_n , respectively, for the boundary layer functions. Then we write $A_n H_n$ as

$$A_n H_n = \tilde{A}_n \tilde{H}_n + (A_n - \tilde{A}_n) H_n + \tilde{A}_n (H_n - \tilde{H}_n), \quad (3.67)$$

and we claim that

$$(A_n - \tilde{A}_n) H_n + \tilde{A}_n (H_n - \tilde{H}_n) = \text{rot}_n R_n \quad (3.68)$$

with a suitable operator R_n , where rot_n is defined in polar coordinates with the n mode for the angular variable. Note that the leading operator $\tilde{A}_n \tilde{H}_n$, when applied to a boundary layer function of the form $h(|\beta|(r-1))$, formally has size $O(|\beta|^4)$, the term $(A_n - \tilde{A}_n) H_n + \tilde{A}_n (H_n - \tilde{H}_n)$ has size $O(|\beta|^3)$, and then R_n is of size $O(|\beta|^2)$.

Let us look for the boundary layer $\varphi_{n,\text{BL}}$ as a solution to

$$\tilde{A}_n \tilde{H}_n \varphi_{n,\text{BL}} = 0. \quad (3.69)$$

Since \tilde{H}_n is easily inverted for $|n| \geq 1$, we start by considering the homogeneous problem $\tilde{A}_n \phi = 0$. By its very definition the operator \tilde{A}_n is nothing but the Airy operator with a complex coefficient. Hence we introduce the Airy function $\text{Ai}(z)$ which is a solution to

$$\frac{d^2 \text{Ai}}{dz^2} - z \text{Ai} = 0$$

in \mathbb{C} ; for details, see Appendix A. Then we define

$$\tilde{G}_{n,\alpha}(\rho) := \text{Ai}\left(c_-(\rho + \frac{in|\beta|}{2\alpha})\right), \quad \rho > 0, \quad (3.70)$$

which satisfies from $c_-^3 = -i$,

$$\left(\frac{d^2}{d\rho^2} + i\left(\rho + \frac{in}{2\alpha}|\beta|\right)\right)\tilde{G}_{n,\alpha} = 0, \quad \rho > 0. \quad (3.71)$$

Next we set

$$G_{n,\alpha}(\rho) := -\int_{\rho}^{\infty} e^{-\frac{|n|}{|\beta|}(\rho-\tau)} \int_{\tau}^{\infty} e^{-\frac{|n|}{|\beta|}(\sigma-\tau)} \tilde{G}_{n,\alpha}(\sigma) d\sigma d\tau, \quad (3.72)$$

which satisfies

$$-\frac{d^2 G_{n,\alpha}}{d\rho^2} + \frac{n^2}{|\beta|^2} G_{n,\alpha} = \tilde{G}_{n,\alpha}, \quad \rho > 0. \quad (3.73)$$

Finally we define

$$C_{0,n,\alpha} = \begin{cases} \frac{1}{G_{n,\alpha}(0)} & \text{if } |G_{n,\alpha}(0)| \geq 1, \\ 1 & \text{otherwise} \end{cases} \quad (3.74)$$

and we set

$$\varphi_{n,\text{BL}}(r) := C_{0,n,\alpha} G_{n,\alpha}(|\beta|(r-1)), \quad (3.75)$$

which satisfies from $-i|\beta|^3 = \beta^3$,

$$\tilde{A}_n \tilde{H}_n \varphi_{n,\text{BL}} = 0, \quad r > 1,$$

as desired. Notice also that

$$|\varphi_{n,\text{BL}}(1)| \leq 1.$$

The key quantity is

$$\begin{aligned} \frac{d\varphi_{n,\text{BL}}}{dr}\Big|_{r=1} &= C_{0,n,\alpha} |\beta| \frac{dG_{n,\alpha}}{d\rho}\Big|_{\rho=0} \\ &= C_{0,n,\alpha} |\beta| \left(\int_0^{\infty} e^{-\frac{|n|}{|\beta|}\sigma} \tilde{G}_{n,\alpha}(\sigma) d\sigma - \frac{|n|}{|\beta|} G_{n,\alpha}(0) \right) \\ &= C_{0,n,\alpha} |\beta| \left(\int_0^{\infty} e^{-\frac{|n|}{|\beta|}\sigma} \text{Ai}\left(c_-(\sigma + \frac{in|\beta|}{2\alpha})\right) d\sigma - \frac{|n|}{|\beta|} G_{n,\alpha}(0) \right) \\ &= C_{0,n,\alpha} |\beta| \left(\frac{1}{c_-} \int_0^{\infty} e^{-\lambda s} \text{Ai}\left(s + \frac{in|\beta|c_-}{2\alpha}\right) ds - \frac{|n|}{|\beta|} G_{n,\alpha}(0) \right), \end{aligned}$$

with

$$\lambda = \lambda_{n,\beta} := \frac{|n|}{|\beta|c_-} = \frac{|n|c_+}{|\beta|}.$$

We find from $\beta^3 = |\beta|^3 c_-^3$ that

$$\frac{in|\beta|c_-}{2\alpha} = \frac{n^2|\beta|c_-}{-2i\alpha n} = \frac{n^2|\beta|c_-}{\beta^3} = \frac{n^2|\beta|}{|\beta|^3 c_-^2} = \frac{n^2 c_+^2}{|\beta|^2} = \lambda^2 \quad (3.76)$$

so

$$\frac{d\varphi_{n,\text{BL}}}{dr}\Big|_{r=1} = C_{0,n,\alpha}|\beta|\left(\frac{1}{c_-}\int_0^\infty e^{-\lambda s}\text{Ai}(s+\lambda^2)ds - \frac{|n|}{|\beta|}G_{n,\alpha}(0)\right). \quad (3.77)$$

Note that $|\lambda|^2 = 4^{-1}|n|^{\frac{4}{3}}\alpha^{-\frac{2}{3}}$ is small when $|n| \ll \alpha^{\frac{1}{2}}$. The proof of the following lemma is postponed to Appendix A.

Lemma 3.7 *There holds*

$$\tilde{C}_0 := \inf\left\{|C_{0,n,\alpha}| \mid \alpha \geq 1, 1 \leq |n| \leq \alpha^{\frac{1}{2}}\right\} > 0, \quad (3.78)$$

and there is a constant $\varepsilon \in (0, 1)$ such that defining

$$\Sigma_\varepsilon := \left\{\mu \in \mathbb{C} \mid \arg \mu = \frac{\pi}{6}, 0 \leq |\mu| \leq \varepsilon\right\},$$

then

$$\tilde{\kappa}_\varepsilon := \inf_{\mu \in \Sigma_\varepsilon} \left| \int_0^\infty e^{-\mu s} \text{Ai}(s + \mu^2) ds \right| > 0. \quad (3.79)$$

Moreover, the function $G_{n,\alpha}$ defined in (3.72) satisfies for R large enough

$$\sup_{\alpha \geq 1} \sup_{1 \leq |n| \leq \alpha^{1/2}} \sup_{\rho \geq R} e^\rho \left| \frac{d^k G_{n,\alpha}}{d\rho^k}(\rho) \right| < \infty, \quad k = 0, 1, 2, 3. \quad (3.80)$$

Now let us return to the proof of Proposition 3.6. We define $\varphi_{n,\text{BL}}$ as in (3.75). Note that thanks to (3.77), (3.79) and (3.78) there holds as long as $\frac{|n|}{|\beta|} = |\lambda| \leq \varepsilon$,

$$\begin{aligned} \left| \frac{d\varphi_{n,\text{BL}}}{dr}(1) \right| &\geq \tilde{C}_0 |\beta| \left(\tilde{\kappa}_\varepsilon - \frac{|n|}{|\beta|} |G_{n,\alpha}(0)| \right) \\ &\geq \tilde{C}_0 |\beta| \left(\tilde{\kappa}_\varepsilon - \frac{\varepsilon}{\tilde{C}_0} \right). \end{aligned}$$

Here we have also used (3.74). Hence

$$\left| \frac{d\varphi_{n,\text{BL}}}{dr}(1) \right| \geq \frac{\tilde{C}_0 \tilde{\kappa}_\varepsilon}{2} |\beta| \quad (3.81)$$

as long as $2\varepsilon \leq \tilde{C}_0 \tilde{\kappa}_\varepsilon$, which is possible since $\tilde{\kappa}_\varepsilon$ is nonincreasing in $\varepsilon > 0$. This proves (3.64) since

$$\begin{aligned} \left| \frac{d\varphi_{n,\text{BL}}}{dr}(1) + |n|\varphi_{n,\text{BL}}(1) \right| &\geq \left| \frac{d\varphi_{n,\text{BL}}}{dr}(1) \right| - |n||\varphi_{n,\text{BL}}(1)| \\ &\geq \frac{\tilde{C}_0 \tilde{\kappa}_\varepsilon}{2} |\beta| - |n| \\ &\geq \frac{\check{\kappa}}{2} |\beta| \end{aligned}$$

with $\check{\kappa} := \tilde{C}_0 \tilde{\kappa}_\varepsilon / 2$, and the last inequality holds as long as $|n| \leq \min(\varepsilon, \check{\kappa}/2) |\beta|$. It then suffices to choose $\kappa \leq \min(\sqrt{2}\varepsilon^{\frac{3}{2}}, \check{\kappa}^{\frac{3}{2}}/2)$ which ensures that

$$|n| \leq \kappa \alpha^{\frac{1}{2}} \Rightarrow |n| \leq \min\left(\varepsilon, \frac{\check{\kappa}}{2}\right) |\beta|.$$

The result (3.65) is an obvious consequence of the previous construction so it remains to prove that (3.63) is satisfied for a suitable $g_{n,\text{BL}}$. Let us recall (3.67). Then we define $g_{n,\text{BL}}$ as

$$g_{n,\text{BL}}(r) := -\frac{1}{r} \int_r^\infty s \left(\tilde{A}_n(H_n - \tilde{H}_n) + (A_n - \tilde{A}_n)H_n \right) \varphi_{n,\text{BL}} \, ds, \quad (3.82)$$

which then satisfies

$$\left((A_n - \tilde{A}_n)H_n + \tilde{A}_n(H_n - \tilde{H}_n) \right) \varphi_{n,\text{BL}} = \frac{1}{r} \frac{d}{dr} (r g_{n,\text{BL}}).$$

Let us consider the estimate of $g_{n,\text{BL}}$. We compute the highest order terms in $\tilde{A}_n(H_n - \tilde{H}_n)$ and $\tilde{A}_n(H_n - \tilde{H}_n)$, which are of order $O(|\beta|^3)$ when $n = O(\alpha^{\frac{1}{2}})$:

$$\begin{aligned} \frac{1}{|\beta|^3} \tilde{A}_n(H_n - \tilde{H}_n) \varphi_{n,\text{BL}} &= \frac{1}{r} G_{n,\alpha}''' + \frac{n^2 r^2 - 1}{|\beta| r^2} G_{n,\alpha}'' - \frac{n^2}{r |\beta|^2} G_{n,\alpha}' - \frac{n^4}{|\beta|^3 r^2} (r^2 - 1) G_{n,\alpha} \\ &\quad + \frac{2i\alpha n}{|\beta|^2 r} (r-1) G_{n,\alpha}' + \frac{2i\alpha n^3}{|\beta|^3 r^2} (r-1)^2 (r+1) G_{n,\alpha} + \text{l.o.t} \end{aligned}$$

where all the $G_{n,\alpha}$ are computed at $|\beta|(r-1)$ and the lower order terms are to be understood in terms of $|\beta|$ for $n = O(\alpha^{\frac{1}{2}})$. It is clear from this formula that as long as $n^2 \lesssim \alpha$ then there is a function $R_{n,\alpha,I}(r, \rho)$, uniformly bounded in r and exponentially decaying at infinity in ρ , such that

$$\tilde{A}_n(H_n - \tilde{H}_n) \varphi_{n,\text{BL}} = |\beta|^3 R_{n,\alpha,I}(r, |\beta|(r-1)).$$

Similarly one has

$$\begin{aligned} \frac{1}{|\beta|^3} (A_n - \tilde{A}_n) H_n \varphi_{n,\text{BL}} &= \frac{1}{r} G_{n,\alpha}''' - \frac{n^2}{|\beta| r^2} (1-r^2) G_{n,\alpha}'' + \frac{2i\alpha n}{|\beta| r^2} (r-1)^2 (1+2r) G_{n,\alpha}' \\ &\quad - \frac{n^2}{r |\beta|^2} G_{n,\alpha}' + \frac{2i\alpha n^3}{|\beta|^2 r^2} (r-1)^2 (1+2r) G_{n,\alpha} + \text{l.o.t} \end{aligned}$$

so again

$$(A_n - \tilde{A}_n) H_n \varphi_{n,\text{BL}} = |\beta|^3 R_{n,\alpha,II}(r, |\beta|(r-1))$$

with some $R_{n,\alpha,II}(r, \rho)$ which is uniformly bounded in r and exponentially decaying as $\rho \rightarrow \infty$. This implies that $g_{n,\text{BL}}$ defined in (3.82) satisfies

$$\|g_{n,\text{BL}}\|_{L^2(\Omega)} \lesssim |\beta|^{\frac{3}{2}}.$$

The result (3.63) follows. Proposition 3.6 is proved. \square

4 The nonlinear problem

In this section we construct the solution to the nonlinear problem $(\widetilde{\text{NS}}_\alpha)$

$$\begin{cases} -\Delta v - \alpha(x^\perp \cdot \nabla v - v^\perp) + \nabla q + \alpha U^\perp \text{rot } v = -v^\perp \text{rot } v + f, & x \in \Omega, \\ \text{div } v = 0, & x \in \Omega, \\ v = 0, & x \in \partial\Omega, \end{cases} \quad (\widetilde{\text{NS}}_\alpha)$$

in the class $v = (\mathcal{P}_0 v, \mathcal{Q}_0 v) \in X = \mathcal{P}_0 W_0^{1,\infty}(\Omega)^2 \times \mathcal{Q}_0 W_0^{1,2}(\Omega)^2$ with a suitable pressure $q \in W_{loc}^{1,1}(\overline{\Omega})$. As already noted in the introduction, for the solvability of $(\widetilde{\text{NS}}_\alpha)$, the key observation is the decomposition of the nonlinear term $v^\perp \text{rot } v$: we have for $v \in X$,

$$\begin{aligned} v^\perp \text{rot } v &= (\mathcal{P}_0 v)^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{P}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{P}_0 v)^\perp \text{rot } \mathcal{P}_0 v \\ &= \mathcal{Q}_0 \left((\mathcal{P}_0 v)^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{P}_0 v \right) + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{P}_0 v)^\perp \text{rot } \mathcal{P}_0 v. \end{aligned}$$

Here we have used $\mathcal{P}_0 \left((\mathcal{P}_0 v)^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{P}_0 v \right) = 0$. Furthermore, since the last term on the right-hand side can be written in a gradient form, the problem $(\widetilde{\text{NS}}_\alpha)$ is in fact reduced to the next system

$$\begin{cases} -\Delta v - \alpha(x^\perp \cdot \nabla v - v^\perp) + \nabla \tilde{q} + \alpha U^\perp \text{rot } v = G(v) + f, & x \in \Omega, \\ \text{div } v = 0, & x \in \Omega, \\ v = 0, & x \in \partial\Omega. \end{cases} \quad (\widehat{\text{NS}}_\alpha)$$

Here we have set

$$G(v) := -\mathcal{Q}_0 \left((\mathcal{P}_0 v)^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{P}_0 v \right) - (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{Q}_0 v, \quad (4.1)$$

and $\tilde{q} := q + Q$, where $Q = Q(|x|)$ is a radial function satisfying $\nabla Q = -(\mathcal{P}_0 v)^\perp \text{rot } \mathcal{P}_0 v$.

Our aim is to prove the existence and uniqueness of solutions (v, q) for $(\widehat{\text{NS}}_\alpha)$ in a suitable subset of X , under some conditions on the external force $f = (\mathcal{P}_0 f, \mathcal{Q}_0 f)$ in $Y = \mathcal{P}_0 L^1(\Omega)^2 \times \mathcal{Q}_0 L^2(\Omega)^2$. The proofs in Subsections 4.2 and 4.3 rely on the standard Banach fixed point argument, where the estimate of the nonlinearity $G(v)$ in the space Y is important. Thanks to the identity

$$\mathcal{P}_0 G(v) = -\mathcal{P}_0 \left((\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{Q}_0 v \right),$$

we see that $\mathcal{P}_0 G(v)$ belongs to $L^1(\Omega)^2$, which is the same summability as the space Y . In order to control the L^2 -norm of $\mathcal{Q}_0 G(v)$ in the iteration scheme, we introduce the closed subspace X_0 of X equipped with the norm $\|\cdot\|_{X_0}$:

$$\begin{aligned} X_0 &:= \left\{ v \in X \mid \text{div } \mathcal{P}_0 v = \text{div } \mathcal{Q}_0 v = 0 \text{ in } \Omega, \right. \\ &\quad \left. \|v\|_{X_0} := \|\mathcal{P}_0 v\|_{L^\infty(\Omega)} + \|\nabla \mathcal{P}_0 v\|_{L^\infty(\Omega)} \right. \\ &\quad \left. + \|\mathcal{Q}_0 v\|_{L^2(\Omega)} + \|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} + \sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} < \infty \right\}. \end{aligned} \quad (4.2)$$

Indeed, we can easily establish an a priori estimate for

$$\mathcal{Q}_0 G(v) = -\mathcal{Q}_0 \left((\mathcal{P}_0 v)^\perp \text{rot } \mathcal{Q}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{P}_0 v + (\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{Q}_0 v \right)$$

in $L^2(\Omega)^2$ for $v \in X_0$; see (4.21) below for the estimate of $\|\mathcal{Q}_0 \left((\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{Q}_0 v \right)\|_{L^2(\Omega)}$.

After proving Theorem 1.1 and 1.2, we revisit Theorem 1.2 in order to study the qualitative behavior of solutions. By fixing the external force $f \in Y$, we consider the fast rotation limit $|\alpha| \rightarrow \infty$ for the solution $(v, q) = (v^{(\alpha)}, q^{(\alpha)})$ to $(\widetilde{\text{NS}}_\alpha)$. The results are summarized in Theorem 1.3, which will be proved in Subsection 4.4.

4.1 Useful estimates on (S_α)

Before we give the proofs of Theorems 1.1 and 1.2, let us restate the main estimates for the linearized problem (S_α) for general α , which will be used throughout this section, as well as some specific estimates corresponding to $|\alpha| \gg 1$.

4.1.1 The case of general α

Let $\alpha \in \mathbb{R} \setminus \{0\}$ and $f \in Y$, and let $(v, q) \in X \times W_{loc}^{1,1}(\overline{\Omega})$ be the unique solution to (S_α) given in Proposition 3.1. For notational simplicity, we denote by v^{low} and v^{high} , respectively, the low and high frequency parts of $\mathcal{Q}_0 v$:

$$v^{\text{low}} := \sum_{1 \leq |n| < 1 + \sqrt{2|\alpha|}} \mathcal{P}_n v, \quad v^{\text{high}} := \sum_{|n| \geq 1 + \sqrt{2|\alpha|}} \mathcal{P}_n v. \quad (4.3)$$

From the estimates (3.5) and (3.9) in Proposition 3.1, we have

$$\|v^{\text{low}}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \left(1 + \frac{1}{|\alpha|^{\frac{1}{2}}}\right) \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (4.4)$$

$$\|v^{\text{high}}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (4.5)$$

and from (3.8) and (3.12) in the same proposition, we have the estimates for the derivatives.

$$\|\nabla v^{\text{low}}\|_{L^2(\Omega)} \leq C \left(1 + \frac{1}{|\alpha|^{\frac{1}{2}}}\right) \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (4.6)$$

$$\|\nabla v^{\text{high}}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (4.7)$$

For the fixed point argument in the space X_0 , the estimate of $\sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)}$ is also needed. In the case $0 < |\alpha| < 1$, (3.7) and (3.11) in Proposition 3.1, and the Hölder inequality for sequences lead to

$$\begin{aligned} \sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} &\leq \frac{C}{|\alpha|^{\frac{3}{4}}} \sum_{|n| \geq 1} \frac{1}{|n|^{\frac{3}{4}}} \|\mathcal{P}_n f\|_{L^2(\Omega)} \\ &\leq \frac{C}{|\alpha|^{\frac{3}{4}}} \left(\sum_{|n| \geq 1} \frac{1}{|n|^{\frac{3}{2}}} \right)^{\frac{1}{2}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \end{aligned}$$

which implies

$$\sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{3}{4}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (4.8)$$

In the case $|\alpha| \geq 1$ we have from (3.7),

$$\sum_{1 \leq |n| < 1 + \sqrt{2|\alpha|}} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{4}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (4.9)$$

and from (3.11) with the Hölder inequality,

$$\sum_{|n| \geq 1 + \sqrt{2|\alpha|}} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (4.10)$$

4.1.2 The case of large $|\alpha|$

Now let us consider the special case when $|\alpha| \gg 1$. Thanks to the boundary layer analysis in Proposition 3.2, we have a better estimate for the linearized problem (S_α) in terms of decay in the parameter $|\alpha n|$. Given the parameter κ defined in Proposition 3.2 we assume that $|\alpha|$ is large enough, as required in Proposition 3.2, and let us truncate frequencies, similarly to (4.3), as follows:

$$v^{\text{low},\kappa} := \sum_{1 \leq |n| \leq \kappa |\alpha|^{\frac{1}{2}}} \mathcal{P}_n v, \quad v^{\text{high},\kappa} := \sum_{|n| > \kappa |\alpha|^{\frac{1}{2}}} \mathcal{P}_n v. \quad (4.11)$$

Then thanks to (3.41) we know that

$$\|v^{\text{low},\kappa}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{2}{3}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (4.12)$$

$$\|\nabla v^{\text{low},\kappa}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{3}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (4.13)$$

and

$$\begin{aligned} \sum_{1 \leq |n| \leq \kappa |\alpha|^{\frac{1}{2}}} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} &\leq \frac{C}{|\alpha|^{\frac{1}{2}}} \left(\sum_{1 \leq |n| \leq \kappa |\alpha|^{\frac{1}{2}}} \frac{1}{|n|} \right)^{\frac{1}{2}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \\ &\leq \frac{C(\log |\alpha|)^{\frac{1}{2}}}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \end{aligned} \quad (4.14)$$

Note that the constant C depends on κ , which is fixed. One can see that the decay in terms of $|\alpha|$ compared with (4.4), (4.6) and (4.9) are improved.

Similarly thanks to (3.9), (3.12), (3.13) and (3.15) there holds

$$\|v^{\text{high},\kappa}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \quad (4.15)$$

$$\|\nabla v^{\text{high},\kappa}\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (4.16)$$

Finally from (3.11), (3.14) and the Hölder inequality, we derive

$$\sum_{|n| \geq \kappa |\alpha|^{\frac{1}{2}}} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (4.17)$$

4.2 Proof for general α

In this subsection we prove Theorem 1.1, by means of a fixed point argument. The solutions to $(\widehat{\text{NS}}_\alpha)$ will be found in the closed convex set $\mathcal{B}_{\vec{\delta},\epsilon}$ of X_0 defined as follows:

$$\begin{aligned} \mathcal{B}_{\vec{\delta},\epsilon} &:= \mathcal{B}_{(\delta_1, \delta_2, \delta_3, \delta_4), \epsilon} \\ &:= \left\{ h \in X_0 \mid \|\mathcal{P}_0 h\|_{L^\infty(\Omega)} + \|\nabla \mathcal{P}_0 h\|_{L^\infty(\Omega)} \leq \epsilon |\alpha|^{\delta_1}, \right. \\ &\quad \left. \|\mathcal{Q}_0 h\|_{L^2(\Omega)} \leq \epsilon |\alpha|^{\delta_2}, \quad \|\nabla \mathcal{Q}_0 h\|_{L^2(\Omega)} \leq \epsilon |\alpha|^{\delta_3}, \quad \sum_{|n| \geq 1} \|\mathcal{P}_n h\|_{L^\infty(\Omega)} \leq \epsilon |\alpha|^{\delta_4} \right\}, \end{aligned}$$

where we have set $\vec{\delta} := (\delta_1, \delta_2, \delta_3, \delta_4)$, and the numbers $\delta_1, \dots, \delta_4$ and the positive number ϵ will be chosen later. For any $h \in \mathcal{B}_{\vec{\delta}, \epsilon}$, let (v_h, q_h) be the unique solution constructed in Proposition 3.1 to the linear system

$$\begin{cases} -\Delta v_h - \alpha(x^\perp \cdot \nabla v_h - v_h^\perp) + \nabla q_h + \alpha U^\perp \text{rot } v_h = G(h) + f, & x \in \Omega, \\ \text{div } v_h = 0, & x \in \Omega, \\ v_h = 0, & x \in \partial\Omega, \end{cases} \quad (4.18)$$

where the function G is defined in (4.1).

Let us start with the estimate of $G(h)$ in the space Y . The first two terms in the right-hand side of (4.1) with v replaced by h can be estimated as

$$\begin{aligned} & \|\mathcal{Q}_0((\mathcal{P}_0 h)^\perp \text{rot } \mathcal{Q}_0 h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0((\mathcal{Q}_0 h)^\perp \text{rot } \mathcal{P}_0 h)\|_{L^2(\Omega)} \\ &= \left(\sum_{|n| \geq 1} \|(\mathcal{P}_0 h)^\perp \text{rot } \mathcal{P}_n h\|_{L^2(\Omega)}^2 \right)^{\frac{1}{2}} + \left(\sum_{|n| \geq 1} \|(\mathcal{P}_n h)^\perp \text{rot } \mathcal{P}_0 h\|_{L^2(\Omega)}^2 \right)^{\frac{1}{2}} \\ &\leq \|\mathcal{P}_0 h\|_{L^\infty(\Omega)} \|\nabla \mathcal{Q}_0 h\|_{L^2(\Omega)} + \|\nabla \mathcal{P}_0 h\|_{L^\infty(\Omega)} \|\mathcal{Q}_0 h\|_{L^2(\Omega)}. \end{aligned} \quad (4.19)$$

For the last term in the right-hand side of (4.1) with v replaced by h , we observe that

$$\mathcal{P}_n((\mathcal{Q}_0 h)^\perp \text{rot } \mathcal{Q}_0 h) = \sum_{k \in \mathbb{Z} \setminus \{0, n\}} (\mathcal{P}_k h)^\perp \text{rot } \mathcal{P}_{n-k} h.$$

Then, applying the Hölder inequality we have

$$\begin{aligned} \|\mathcal{P}_0((\mathcal{Q}_0 h)^\perp \text{rot } \mathcal{Q}_0 h)\|_{L^1(\Omega)} &\leq \sum_{k \in \mathbb{Z} \setminus \{0\}} \|\mathcal{P}_k h\|_{L^2(\Omega)} \|\mathcal{P}_{-k} \nabla h\|_{L^2(\Omega)} \\ &\leq \|\mathcal{Q}_0 h\|_{L^2(\Omega)} \|\nabla \mathcal{Q}_0 h\|_{L^2(\Omega)}, \end{aligned} \quad (4.20)$$

and the Young inequality for sequences implies that

$$\begin{aligned} \|\mathcal{Q}_0((\mathcal{Q}_0 h)^\perp \text{rot } \mathcal{Q}_0 h)\|_{L^2(\Omega)} &= \left(\sum_{|n| \geq 1} \|\mathcal{P}_n((\mathcal{Q}_0 h)^\perp \text{rot } \mathcal{Q}_0 h)\|_{L^2(\Omega)}^2 \right)^{\frac{1}{2}} \\ &\leq \left(\sum_{|n| \geq 1} \left(\sum_{k \in \mathbb{Z} \setminus \{0, n\}} \|\mathcal{P}_k h\|_{L^\infty(\Omega)} \|\nabla \mathcal{P}_{n-k} h\|_{L^2(\Omega)} \right)^2 \right)^{\frac{1}{2}} \\ &\leq \left(\sum_{|n| \geq 1} \|\mathcal{P}_n h\|_{L^\infty(\Omega)} \right) \|\nabla \mathcal{Q}_0 h\|_{L^2(\Omega)}. \end{aligned} \quad (4.21)$$

Next we estimate the difference $G(h^{(1)}) - G(h^{(2)})$ for $h^{(1)}, h^{(2)} \in X$. Setting $\mathbf{h} = (h^{(1)}, h^{(2)})$ for simplicity, we define the function $H(\mathbf{h})$ on $\mathcal{B}_{\vec{\delta}, \epsilon} \times \mathcal{B}_{\vec{\delta}, \epsilon}$ as

$$\begin{aligned} H(\mathbf{h}) &:= G(h^{(1)}) - G(h^{(2)}) \\ &= \mathcal{Q}_0 \left((\mathcal{P}_0 h^{(1)} - \mathcal{P}_0 h^{(2)})^\perp \text{rot } \mathcal{Q}_0 h^{(1)} + (\mathcal{P}_0 h^{(2)})^\perp \text{rot } (\mathcal{Q}_0 h^{(1)} - \mathcal{Q}_0 h^{(2)}) \right. \\ &\quad \left. + (\mathcal{Q}_0 h^{(1)} - \mathcal{Q}_0 h^{(2)})^\perp \text{rot } \mathcal{P}_0 h^{(1)} + (\mathcal{Q}_0 h^{(2)})^\perp \text{rot } (\mathcal{P}_0 h^{(1)} - \mathcal{P}_0 h^{(2)}) \right) \\ &\quad + (\mathcal{Q}_0 h^{(1)} - \mathcal{Q}_0 h^{(2)})^\perp \text{rot } \mathcal{Q}_0 h^{(1)} + (\mathcal{Q}_0 h^{(2)})^\perp \text{rot } (\mathcal{Q}_0 h^{(1)} - \mathcal{Q}_0 h^{(2)}). \end{aligned} \quad (4.22)$$

The following estimates on $H(\mathbf{h})$ are obtained in the same way as (4.19)-(4.21):

$$\|\mathcal{P}_0 H(\mathbf{h})\|_{L^1(\Omega)} \leq (\|\nabla \mathcal{Q}_0 h^{(1)}\|_{L^2(\Omega)} + \|\mathcal{Q}_0 h^{(2)}\|_{L^2(\Omega)}) \|h^{(1)} - h^{(2)}\|_{X_0}, \quad (4.23)$$

$$\begin{aligned} \|\mathcal{Q}_0 H(\mathbf{h})\|_{L^2(\Omega)} &\leq \left(\|\nabla \mathcal{Q}_0 h^{(1)}\|_{L^2(\Omega)} + \|\mathcal{Q}_0 h^{(2)}\|_{L^2(\Omega)} + \|\nabla \mathcal{P}_0 h^{(1)}\|_{L^\infty(\Omega)} \right. \\ &\quad \left. + \|\mathcal{P}_0 h^{(2)}\|_{L^\infty(\Omega)} + \sum_{|n| \geq 1} \|\mathcal{P}_n h^{(2)}\|_{L^\infty(\Omega)} \right) \|h^{(1)} - h^{(2)}\|_{X_0}. \end{aligned} \quad (4.24)$$

From (4.19)-(4.21), (4.23)-(4.24), and the definition of $\mathcal{B}_{\vec{\delta}, \epsilon}$, we obtain the following estimates on $G(h)$ and $H(\mathbf{h})$ in Y .

$$\|\mathcal{P}_0 G(h)\|_{L^1(\Omega)} \leq \epsilon^2 |\alpha|^{\delta_2 + \delta_3}, \quad (4.25)$$

$$\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} \leq \epsilon^2 (|\alpha|^{\delta_1 + \delta_2} + |\alpha|^{\delta_1 + \delta_3} + |\alpha|^{\delta_3 + \delta_4}), \quad (4.26)$$

$$\|\mathcal{P}_0 H(\mathbf{h})\|_{L^1(\Omega)} \leq \epsilon (|\alpha|^{\delta_2} + |\alpha|^{\delta_3}) \|h^{(1)} - h^{(2)}\|_{X_0}, \quad (4.27)$$

$$\|\mathcal{Q}_0 H(\mathbf{h})\|_{L^2(\Omega)} \leq \epsilon (|\alpha|^{\delta_1} + |\alpha|^{\delta_2} + |\alpha|^{\delta_3} + |\alpha|^{\delta_4}) \|h^{(1)} - h^{(2)}\|_{X_0}. \quad (4.28)$$

Now let us define the mapping $\Phi : \mathcal{B}_{\vec{\delta}, \epsilon} \rightarrow X_0$ by setting $\Phi[h] := v_h$, where v_h is the unique solution to (4.18). Our aim is to show that

(i) Φ is Lipschitz continuous on $\mathcal{B}_{\vec{\delta}, \epsilon}$ in the topology of X_0 . Namely, there exists $\tau \in (0, 1)$ such that $\|\Phi[h^{(1)}] - \Phi[h^{(2)}]\|_{X_0} \leq \tau \|h^{(1)} - h^{(2)}\|_{X_0}$ for any $h^{(1)}, h^{(2)} \in \mathcal{B}_{\vec{\delta}, \epsilon}$,

(ii) Φ is a mapping from $\mathcal{B}_{\vec{\delta}, \epsilon}$ into $\mathcal{B}_{\vec{\delta}, \epsilon}$, if the pair $(\vec{\delta}, \epsilon)$ and the external force $f = (\mathcal{P}_0 f, \mathcal{Q}_0 f) \in Y$ satisfy a suitable condition.

For convenience in the following proof, let K_1 and K_2 denote the largest constant C (larger than 1 without loss of generality) appearing in (3.4), (4.4)-(4.8), and (3.4), (4.4)-(4.7), (4.9)-(4.10), respectively.

We first show (i). For any $\mathbf{h} = (h^{(1)}, h^{(2)}) \in \mathcal{B}_{\vec{\delta}, \epsilon} \times \mathcal{B}_{\vec{\delta}, \epsilon}$, we observe that the differences $u_{\mathbf{h}} := \Phi[h^{(1)}] - \Phi[h^{(2)}]$ and $p_{\mathbf{h}} := q_{h^{(1)}} - q_{h^{(2)}}$ solve the following system:

$$\begin{cases} -\Delta u_{\mathbf{h}} - \alpha(x^\perp \cdot \nabla u_{\mathbf{h}} - u_{\mathbf{h}}^\perp) + \nabla p_{\mathbf{h}} + \alpha U^\perp \text{rot } u_{\mathbf{h}} = H(\mathbf{h}), & x \in \Omega, \\ \text{div } u_{\mathbf{h}} = 0, & x \in \Omega, \\ u_{\mathbf{h}} = 0, & x \in \partial\Omega, \end{cases}$$

where $H(\mathbf{h})$ is defined in (4.22). We consider the case $0 < |\alpha| < 1$. Then for any $h^{(1)}, h^{(2)}$ in X_0 , by (3.4) and (4.4)-(4.8) combined with (4.27)-(4.28), we see that

$$\begin{aligned} &\|\Phi[h^{(1)}] - \Phi[h^{(2)}]\|_{X_0} \\ &= \|\mathcal{P}_0 u_{\mathbf{h}}\|_{L^\infty(\Omega)} + \|\nabla \mathcal{P}_0 u_{\mathbf{h}}\|_{L^\infty(\Omega)} + \|\mathcal{Q}_0 u_{\mathbf{h}}\|_{L^2(\Omega)} + \|\nabla \mathcal{Q}_0 u_{\mathbf{h}}\|_{L^2(\Omega)} + \sum_{|n| \geq 1} \|\mathcal{P}_n u_{\mathbf{h}}\|_{L^\infty(\Omega)} \\ &\leq K_1 \|\mathcal{P}_0 H(\mathbf{h})\|_{L^1(\Omega)} + \frac{7K_1}{|\alpha|} \|\mathcal{Q}_0 H(\mathbf{h})\|_{L^2(\Omega)} \\ &\leq 8K_1 \epsilon (|\alpha|^{\delta_1 - 1} + |\alpha|^{\delta_2 - 1} + |\alpha|^{\delta_3 - 1} + |\alpha|^{\delta_4 - 1}) \|h^{(1)} - h^{(2)}\|_{X_0}. \end{aligned}$$

Hence, if we choose the pair $(\vec{\delta}, \epsilon)$ to satisfy

$$\delta_j \geq 1 \quad \text{for } j = 1 \dots 4, \quad 0 < \epsilon < \frac{1}{32K_1}, \quad \text{when } 0 < |\alpha| < 1, \quad (4.29)$$

then the mapping Φ is Lipschitz continuous on the set $\mathcal{B}_{\vec{\delta}, \epsilon}$. For the case $|\alpha| \geq 1$, from the estimates (3.4), (4.4)-(4.7), and (4.9)-(4.10) combined with (4.27)-(4.28), we have

$$\begin{aligned} \|\Phi[h^{(1)}] - \Phi[h^{(2)}]\|_{X_0} &\leq K_2 \|\mathcal{P}_0 H(\mathbf{h})\|_{L^1(\Omega)} + 8K_2 \|\mathcal{Q}_0 H(\mathbf{h})\|_{L^2(\Omega)} \\ &\leq 9K_1 \epsilon (|\alpha|^{\delta_1} + |\alpha|^{\delta_2} + |\alpha|^{\delta_3} + |\alpha|^{\delta_4}) \|h^{(1)} - h^{(2)}\|_{X_0}. \end{aligned} \quad (4.30)$$

Then we obtain the next condition

$$\delta_j \leq 0 \quad \text{for } j = 1 \dots 4, \quad 0 < \epsilon < \frac{1}{36K_2}, \quad \text{when } |\alpha| \geq 1, \quad (4.31)$$

for the Lipschitz continuity of Φ on $\mathcal{B}_{\vec{\delta}, \epsilon}$. We have shown (i) provided the pair $(\vec{\delta}, \epsilon)$ satisfies the conditions (4.29) or (4.31).

Next we prove (ii). In the case $0 < |\alpha| < 1$, the estimates (3.4) and (4.25) imply

$$\begin{aligned} \|\mathcal{P}_0 \Phi[h]\|_{L^\infty(\Omega)} + \|\nabla \mathcal{P}_0 \Phi[h]\|_{L^\infty(\Omega)} &\leq K_1 (\|(\mathcal{P}_0 G(h))_\theta\|_{L^1(\Omega)} + \|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)}) \\ &\leq K_1 (\epsilon^2 |\alpha|^{\delta_2 + \delta_3} + \|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)}), \end{aligned} \quad (4.32)$$

for any $h \in X_0$. From (4.4)-(4.5), (4.6)-(4.7), and (4.26) we have,

$$\begin{aligned} \|\mathcal{Q}_0 \Phi[h]\|_{L^2(\Omega)} &\leq \frac{3K_1}{|\alpha|} (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 3K_1 \left(\epsilon^2 (|\alpha|^{\delta_1 + \delta_2 - 1} + |\alpha|^{\delta_1 + \delta_3 - 1} + |\alpha|^{\delta_3 + \delta_4 - 1}) + |\alpha|^{-1} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right), \\ \|\nabla \mathcal{Q}_0 \Phi[h]\|_{L^2(\Omega)} &\leq \frac{3K_1}{|\alpha|^{\frac{1}{2}}} (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 3K_1 \left(\epsilon^2 (|\alpha|^{\delta_1 + \delta_2 - \frac{1}{2}} + |\alpha|^{\delta_1 + \delta_3 - \frac{1}{2}} + |\alpha|^{\delta_3 + \delta_4 - \frac{1}{2}}) + |\alpha|^{-\frac{1}{2}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right), \end{aligned}$$

and

$$\begin{aligned} \sum_{|n| \geq 1} \|\mathcal{P}_n \Phi[h]\|_{L^\infty(\Omega)} &\leq \frac{K_1}{|\alpha|^{\frac{3}{4}}} (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq K_1 \left(\epsilon^2 (|\alpha|^{\delta_1 + \delta_2 - \frac{3}{4}} + |\alpha|^{\delta_1 + \delta_3 - \frac{3}{4}} + |\alpha|^{\delta_3 + \delta_4 - \frac{3}{4}}) + |\alpha|^{-\frac{3}{4}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right). \end{aligned}$$

Hence, recalling the condition (4.29), if we choose the pair $(\vec{\delta}, \epsilon)$ and $f \in Y$ to satisfy

$$\delta_1 = \delta_2 = \delta_3 = \delta_4 = 1, \quad 0 < \epsilon < \frac{1}{32K_1}, \quad \text{when } 0 < |\alpha| < 1, \quad (4.33)$$

and

$$\|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} \leq \frac{\epsilon}{2K_1} |\alpha|, \quad \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \leq \frac{\epsilon}{6K_1} |\alpha|^2, \quad \text{when } 0 < |\alpha| < 1, \quad (4.34)$$

then Φ defines a mapping from $\mathcal{B}_{\vec{\delta}, \epsilon}$ into itself.

In the case $|\alpha| \geq 1$, we have (4.32) with K_1 replaced by K_2 , and moreover, from (4.4)-(4.5), (4.6)-(4.7), and (4.9)-(4.10), along with (4.26), we have

$$\begin{aligned} \|\mathcal{Q}_0 \Phi[h]\|_{L^2(\Omega)} &\leq \frac{3K_2}{|\alpha|^{\frac{1}{2}}} (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 3K_2 \left(\epsilon^2 (|\alpha|^{\delta_1 + \delta_2 - \frac{1}{2}} + |\alpha|^{\delta_1 + \delta_3 - \frac{1}{2}} + |\alpha|^{\delta_3 + \delta_4 - \frac{1}{2}}) + |\alpha|^{-\frac{1}{2}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right), \end{aligned} \quad (4.35)$$

$$\begin{aligned} \|\nabla \mathcal{Q}_0 \Phi[h]\|_{L^2(\Omega)} &\leq 3K_2 (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 3K_2 \left(\epsilon^2 (|\alpha|^{\delta_1 + \delta_2} + |\alpha|^{\delta_1 + \delta_3} + |\alpha|^{\delta_3 + \delta_4}) + \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right), \end{aligned} \quad (4.36)$$

and

$$\begin{aligned} \sum_{|n| \geq 1} \|\mathcal{P}_n \Phi[h]\|_{L^\infty(\Omega)} &\leq \frac{2K_2}{|\alpha|^{\frac{1}{4}}} (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 2K_2 \left(\epsilon^2 (|\alpha|^{\delta_1 + \delta_2 - \frac{1}{4}} + |\alpha|^{\delta_1 + \delta_3 - \frac{1}{4}} + |\alpha|^{\delta_3 + \delta_4 - \frac{1}{4}}) + |\alpha|^{-\frac{1}{4}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right). \end{aligned} \quad (4.37)$$

Then recalling (4.31), if the pair $(\vec{\delta}, \epsilon)$ and $f \in Y$ satisfy

$$\delta_1 = \delta_2 = \delta_3 = \delta_4 = 0, \quad 0 < \epsilon < \frac{1}{36K_2}, \quad \text{when } |\alpha| \geq 1, \quad (4.38)$$

and

$$\|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} \leq \frac{\epsilon}{2K_2}, \quad \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \leq \frac{\epsilon}{6K_2}, \quad \text{when } |\alpha| \geq 1, \quad (4.39)$$

then we see that Φ defines a mapping $\Phi : \mathcal{B}_{\vec{\delta}, \epsilon} \rightarrow \mathcal{B}_{\vec{\delta}, \epsilon}$.

Now we have shown that the mapping Φ defines a contraction on $\mathcal{B}_{\vec{\delta}, \epsilon}$ under the conditions (4.33) and (4.34) when $0 < |\alpha| < 1$, and under (4.38) and (4.39) when $|\alpha| \geq 1$. Then there is a unique fixed point of Φ in the closed convex set $\mathcal{B}_{\vec{\delta}, \epsilon}$. Hence we finally obtain a unique solution to the nonlinear problem $(\widehat{\text{NS}}_\alpha)$ in $\mathcal{B}_{\vec{\delta}, \epsilon}$.

The estimates in Theorem 1.1 are obtained as follows. We consider only the case $0 < |\alpha| < 1$, in particular the estimates (1.9)-(1.12), since the case $|\alpha| \geq 1$ can be handled similarly. Let v denote the unique fixed point of Φ in $\mathcal{B}_{\vec{\delta}, \epsilon}$. Applying the linear estimates (4.4) and (4.5) to $(\widehat{\text{NS}}_\alpha)$ along with (4.19)-(4.21) we see that

$$\begin{aligned} \|\mathcal{Q}_0 v\|_{L^2(\Omega)} &\leq \frac{3K_1}{|\alpha|} (\|\mathcal{Q}_0 G(v)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq \frac{3K_1}{|\alpha|} \left(\|\mathcal{P}_0 v\|_{L^\infty(\Omega)} \|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} + \|\nabla \mathcal{P}_0 v\|_{L^\infty(\Omega)} \|\mathcal{Q}_0 v\|_{L^2(\Omega)} \right. \\ &\quad \left. + \left(\sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} \right) \|\mathcal{Q}_0 v\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right) \\ &\leq 3K_1 \epsilon (\|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} + 2\|\mathcal{Q}_0 v\|_{L^2(\Omega)}) + \frac{3K_1}{|\alpha|} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \end{aligned}$$

In the last inequality we have applied the bounds derived from the assumption $v \in \mathcal{B}_{\vec{\delta}, \epsilon}$ with $\delta_j = 1$ for all j . Since $0 < 6K_1 \epsilon < \frac{1}{5}$ under the choice of ϵ in (4.33), we obtain

$$\|\mathcal{Q}_0 v\|_{L^2(\Omega)} \leq \frac{15}{4} K_1 \epsilon \|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} + \frac{15}{4} \frac{K_1}{|\alpha|} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \quad (4.40)$$

On the other hand, if we apply (4.6) and (4.7) to $(\widehat{\text{NS}}_\alpha)$, then we have by the same argument

$$\begin{aligned} \|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} &\leq \frac{3K_1}{|\alpha|^{\frac{1}{2}}} (\|\mathcal{Q}_0 G(v)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 6K_1 \epsilon |\alpha|^{\frac{1}{2}} \|\mathcal{Q}_0 v\|_{L^2(\Omega)} + 3K_1 \epsilon |\alpha|^{\frac{1}{2}} \|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} + \frac{3K_1}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}. \end{aligned}$$

Inserting (4.40) in the above inequality, and by the smallness of $K_1\epsilon$ again, we see that

$$\|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} \leq \frac{C}{|\alpha|^{\frac{1}{2}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)},$$

which implies the estimate (1.12). We can obtain (1.10) by (4.40) combined with (1.12) with the condition $0 < |\alpha| < 1$. For the estimate (1.11), by (4.8), (4.19) and (4.21), and the condition $v \in \mathcal{B}_{\vec{\delta}, \epsilon}$ with $\delta_j = 1$ for all j we have

$$\begin{aligned} \sum_{|n| \geq 1} \|\mathcal{P}_n v\|_{L^\infty(\Omega)} &\leq \frac{K_1}{|\alpha|^{\frac{3}{4}}} (\|\mathcal{Q}_0 G(v)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq K_1 \epsilon |\alpha|^{\frac{1}{4}} (\|\nabla \mathcal{Q}_0 v\|_{L^2(\Omega)} + 2\|\mathcal{Q}_0 v\|_{L^2(\Omega)}) + \frac{K_1}{|\alpha|^{\frac{3}{4}}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)}, \end{aligned}$$

which combined with (1.10) and (1.12) leads to (1.11). The estimate (1.9) follows from (3.4) and the nonlinear estimate (4.20) with h replaced by v . This completes the proof of Theorem 1.1. \square

4.3 Proof for large $|\alpha|$

In this subsection we prove Theorem 1.2. We adopt the same notation as in the proof of Theorem 1.1 of the previous subsection. Let K_3 denote the largest constant of C (larger than 1 without loss of generality) appearing in (3.4) and (4.12)-(4.17) for convenience.

We show that the mapping Φ is a contraction on the set $\mathcal{B}_{\vec{\delta}, \epsilon}$. For the Lipschitz continuity of Φ on $\mathcal{B}_{\vec{\delta}, \epsilon}$, we observe that the estimate (4.30) is improved in terms of the decay in $|\alpha|$ into

$$\begin{aligned} \|\Phi[h^{(1)}] - \Phi[h^{(2)}]\|_{X_0} &\leq K_3 \|(\mathcal{P}_0 H(\mathbf{h}))_\theta\|_{L^1(\Omega)} + \frac{6K_3}{|\alpha|^{\frac{1}{3}}} \|\mathcal{Q}_0 H(\mathbf{h})\|_{L^2(\Omega)} \\ &\leq 7K_3 \epsilon (|\alpha|^{\delta_1 - \frac{1}{3}} + |\alpha|^{\delta_2} + |\alpha|^{\delta_3} + |\alpha|^{\delta_4 - \frac{1}{3}}) \|h^{(1)} - h^{(2)}\|_{X_0}. \end{aligned}$$

Hence, if we choose the pair $(\vec{\delta}, \epsilon)$ to satisfy

$$\delta_1 \leq \frac{1}{3}, \quad \delta_2 \leq 0, \quad \delta_3 \leq 0, \quad \delta_4 \leq \frac{1}{3}, \quad 0 < \epsilon < \frac{1}{28K_3}, \quad (4.41)$$

then the mapping Φ is Lipschitz continuous on $\mathcal{B}_{\vec{\delta}, \epsilon}$.

Next we check that Φ is a mapping from $\mathcal{B}_{\vec{\delta}, \epsilon}$ into $\mathcal{B}_{\vec{\delta}, \epsilon}$. We have (4.32) with K_1 replaced by K_3 , and we see that (4.35)-(4.37) are improved in terms of decay in $|\alpha|$ respectively to

$$\begin{aligned} \|\mathcal{Q}_0 \Phi[h]\|_{L^2(\Omega)} &\leq \frac{2K_3}{|\alpha|^{\frac{2}{3}}} (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 2K_3 \left(\epsilon^2 (|\alpha|^{\delta_1 + \delta_2 - \frac{2}{3}} + |\alpha|^{\delta_1 + \delta_3 - \frac{2}{3}} + |\alpha|^{\delta_3 + \delta_4 - \frac{2}{3}}) + |\alpha|^{-\frac{2}{3}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right), \\ \|\nabla \mathcal{Q}_0 \Phi[h]\|_{L^2(\Omega)} &\leq \frac{2K_3}{|\alpha|^{\frac{1}{3}}} (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 2K_3 \left(\epsilon^2 (|\alpha|^{\delta_1 + \delta_2 - \frac{1}{3}} + |\alpha|^{\delta_1 + \delta_3 - \frac{1}{3}} + |\alpha|^{\delta_3 + \delta_4 - \frac{1}{3}}) + |\alpha|^{-\frac{1}{3}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right), \end{aligned}$$

and

$$\begin{aligned} \sum_{|n| \geq 1} \|\mathcal{P}_n \Phi[h]\|_{L^\infty(\Omega)} &\leq \frac{2K_3(\log |\alpha|)^{\frac{1}{2}}}{|\alpha|^{\frac{1}{2}}} (\|\mathcal{Q}_0 G(h)\|_{L^2(\Omega)} + \|\mathcal{Q}_0 f\|_{L^2(\Omega)}) \\ &\leq 2K_3 \left(\epsilon^2 (\log |\alpha|)^{\frac{1}{2}} (|\alpha|^{\delta_1 + \delta_2 - \frac{1}{2}} + |\alpha|^{\delta_1 + \delta_3 - \frac{1}{2}} + |\alpha|^{\delta_3 + \delta_4 - \frac{1}{2}}) \right. \\ &\quad \left. + (\log |\alpha|)^{\frac{1}{2}} |\alpha|^{-\frac{1}{2}} \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \right). \end{aligned}$$

Then, under the condition (4.41), if we choose the pair $(\vec{\delta}, \epsilon)$ and $f \in Y$ to satisfy

$$\delta_1 = \delta_4 = \frac{1}{3}, \quad \delta_2 = \delta_3 = 0, \quad 0 < \epsilon < \frac{1}{28K_3}, \quad (4.42)$$

and

$$\|(\mathcal{P}_0 f)_\theta\|_{L^1(\Omega)} \leq \frac{\epsilon}{2K_3} |\alpha|^{\frac{1}{3}}, \quad \|\mathcal{Q}_0 f\|_{L^2(\Omega)} \leq \frac{\epsilon}{6K_2} |\alpha|^{\frac{1}{3}}, \quad (4.43)$$

then Φ defines a mapping $\Phi : \mathcal{B}_{\vec{\delta}, \epsilon} \rightarrow \mathcal{B}_{\vec{\delta}, \epsilon}$. Now we have shown that Φ is a contraction on $\mathcal{B}_{\vec{\delta}, \epsilon}$ if we assume the conditions (4.42) and (4.43). Hence there exists a unique fixed point v of Φ in $\mathcal{B}_{\vec{\delta}, \epsilon}$. The estimates (1.22)-(1.25) can be obtained in the same way as in the proof of Theorem 1.1. This completes the proof of Theorem 1.2. \square

4.4 Proof of Theorem 1.3

This subsection is devoted to the proof of Theorem 1.3. For a given $f \in Y$, let us take $\alpha \in \mathbb{R}$ large enough to satisfy both the condition (1.21) in Theorem 1.2 and the assumption in Proposition 3.2. Then, from Theorem 1.2 we see that there exists a solution $(v^{(\alpha)}, q^{(\alpha)})$ in $X \times W_{loc}^{1,1}(\overline{\Omega})$ to $(\widetilde{\text{NS}}_\alpha)$ satisfying the estimates (1.22)-(1.25). Hence, the proof will be completed as soon as we show all the estimates in Theorem 1.3 and the decomposition (1.27) of $v^{(\alpha)}$. Note that $v^{(\alpha)}$ also solves the system $(\widehat{\text{NS}}_\alpha)$, which is introduced in the beginning of this section, with a suitable new pressure $\tilde{q}^{(\alpha)} \in W_{loc}^{1,1}(\overline{\Omega})$.

We deduce the estimate (1.26) from the triangle inequality

$$\begin{aligned} \|v^{(\alpha)} - v_0^{\text{linear}}\|_{L^\infty(\Omega)} &\leq \|\mathcal{Q}_0 v^{(\alpha)}\|_{L^\infty(\Omega)} + \|\mathcal{P}_0 v^{(\alpha)} - v_0^{\text{linear}}\|_{L^\infty(\Omega)} \\ &\leq \sum_{|n| \geq 1} \|\mathcal{P}_n v^{(\alpha)}\|_{L^\infty(\Omega)} + \|\mathcal{P}_0 v^{(\alpha)} - v_0^{\text{linear}}\|_{L^\infty(\Omega)}. \end{aligned} \quad (4.44)$$

Since $v := \mathcal{P}_0 v^{(\alpha)} - v_0^{\text{linear}} \in W_0^{1,\infty}(\Omega)^2$ is a solution to the next system

$$\begin{cases} -\Delta v - \alpha(x^\perp \cdot \nabla v - v^\perp) + \nabla q + \alpha U^\perp \text{rot } v = -\mathcal{P}_0 \left((\mathcal{Q}_0 v)^\perp \text{rot } \mathcal{Q}_0 v \right), & x \in \Omega, \\ \text{div } v = 0, & x \in \Omega, \\ v = 0, & x \in \partial\Omega, \end{cases}$$

with some pressure $q \in W_{loc}^{1,1}(\overline{\Omega})$, we have from (3.4) and (4.20) with h replaced by $v^{(\alpha)}$,

$$\|\mathcal{P}_0 v^{(\alpha)} - v_0^{\text{linear}}\|_{L^\infty(\Omega)} \leq C \|\mathcal{Q}_0 v^{(\alpha)}\|_{L^2(\Omega)} \|\nabla \mathcal{Q}_0 v^{(\alpha)}\|_{L^2(\Omega)}. \quad (4.45)$$

Then, by (4.44)-(4.45) along with (1.23)-(1.25) we obtain the estimate (1.26).

The decomposition (1.27) and the related estimates follow from the results in Proposition 3.2. Indeed, we know that $(v^{(\alpha)}, \tilde{q}^{(\alpha)})$ solves $(\widetilde{\text{NS}}_\alpha)$, and we have the next estimate of $\|\mathcal{P}_n G(v^{(\alpha)})\|_{L^2(\Omega)}$, which is uniform both in $|n| \geq 1$ and $|\alpha|$, combined with estimates (1.22)-(1.25):

$$\begin{aligned} \|\mathcal{P}_n G(v^{(\alpha)})\|_{L^2(\Omega)} &\leq \|\mathcal{Q}_0 G(v^{(\alpha)})\|_{L^2(\Omega)} \\ &\leq \|\mathcal{Q}_0((\mathcal{P}_0 v^{(\alpha)})^\perp \text{rot } \mathcal{Q}_0 v^{(\alpha)})\|_{L^2(\Omega)} + \|\mathcal{Q}_0((\mathcal{Q}_0 v^{(\alpha)})^\perp \text{rot } \mathcal{P}_0 v^{(\alpha)})\|_{L^2(\Omega)} \\ &\quad + \|\mathcal{Q}_0((\mathcal{Q}_0 v^{(\alpha)})^\perp \text{rot } \mathcal{Q}_0 v^{(\alpha)})\|_{L^2(\Omega)} \\ &\leq \|\mathcal{P}_0 v^{(\alpha)}\|_{L^\infty(\Omega)} \|\nabla \mathcal{Q}_0 v^{(\alpha)}\|_{L^2(\Omega)} + \|\nabla \mathcal{P}_0 v^{(\alpha)}\|_{L^\infty(\Omega)} \|\mathcal{Q}_0 v^{(\alpha)}\|_{L^2(\Omega)} \\ &\quad + \left(\sum_{|n| \geq 1} \|\mathcal{P}_n v^{(\alpha)}\|_{L^\infty(\Omega)} \right) \|\nabla \mathcal{Q}_0 v^{(\alpha)}\|_{L^2(\Omega)}, \end{aligned}$$

where the nonlinear estimates (4.19) and (4.21) with h replaced by $v^{(\alpha)}$ are applied. The proof of Theorem 1.3 is complete. \square

A Solving the boundary layer equation: proof of Lemma 3.7

In this section we prove Lemma 3.7. All the results concerning the Airy function can be found for instance in [1], Chapter 10. Let us first recall that a solution to the Airy equation

$$\frac{d^2 f(\rho)}{d\rho^2} - \rho f(\rho) = 0 \quad \text{in } \mathbb{R}$$

is given by the Airy function Ai , which can be extended as an entire analytic function on \mathbb{C} satisfying

$$\frac{d^2 f(z)}{dz^2} - z f(z) = 0 \quad \text{in } \mathbb{C}. \quad (\text{A.1})$$

It is the inverse Fourier transform of

$$\xi \mapsto \exp\left(\frac{i\xi^3}{3}\right)$$

and satisfies

$$\text{Ai}(0) = \frac{1}{3^{\frac{2}{3}} \Gamma\left(\frac{2}{3}\right)}, \quad \text{Ai}'(0) = -\frac{1}{3^{\frac{1}{3}} \Gamma\left(\frac{1}{3}\right)},$$

where Γ is the Gamma function. Moreover

$$\text{Ai}(z) \sim_{|z| \rightarrow \infty} z^{-\frac{1}{4}} \exp\left(-\frac{2}{3} z^{\frac{3}{2}}\right), \quad |\arg z| < \pi - \epsilon, \quad \epsilon > 0. \quad (\text{A.2})$$

The results (3.78) and (3.80) are easy consequences of (A.2). Indeed we can write

$$|C_{0,n,\alpha}|^{-1} = \left| \int_0^\infty e^{\frac{2|n|}{|\beta|}t} \int_t^\infty e^{-\frac{|n|}{|\beta|}s} \text{Ai}(c_- s + \lambda^2) ds dt \right|,$$

where $\lambda = \frac{|n|c_+}{|\beta|}$ with $|\beta| = (2|\alpha n|)^{\frac{1}{3}}$ and $c_\pm = \frac{\sqrt{3} \pm i}{2}$ (hence $\frac{|n|\beta|c_-}{2\alpha} = \lambda^2$), and therefore $|C_{0,n,\alpha}|^{-1}$ is bounded uniformly in the set $\{(n, \alpha) \in \mathbb{Z} \times \mathbb{R} \mid |\alpha| \geq 1, 1 \leq |n| \leq |\alpha|^{\frac{1}{2}}\}$

thanks to (A.2). The result (3.80) is obtained in the same way. As to (3.79), it is known (see for instance [1] page 449) that

$$\int_0^\infty \text{Ai}(s) \, ds = \frac{1}{3},$$

so the result follows by continuity of the map $\mu \mapsto \int_0^\infty e^{-\mu s} \text{Ai}(s + \mu^2) \, ds$. This concludes the proof of Lemma 3.7. \square

B Proof of the interpolation inequality (3.28)

We may assume that $g \in W^{1,2}((1, \infty); r \, dr)$ is nontrivial. Let $\delta \in (0, 1]$ be a fixed number which will be determined later. Then we have

$$\begin{aligned} \|g\|_{L^2(\Omega)}^2 &= 2\pi \int_1^\infty |g|^2 r \, dr \\ &\leq C \int_1^{1+\delta} r \, dr \|g\|_{L^\infty((1, \infty))}^2 + \frac{C}{\delta} \int_{1+\delta}^2 \frac{r^2-1}{r^2} |g|^2 r \, dr + C \int_2^\infty \frac{r^2-1}{r^2} |g|^2 r \, dr \\ &\leq C\delta \|g\|_{L^\infty((1, \infty))}^2 + \frac{C}{\delta} \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}^2 + C \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}^2. \end{aligned} \quad (\text{B.1})$$

Let us take

$$\delta = \frac{\left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}}{\|g\|_{L^\infty((1, \infty))} + \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}}.$$

Then

$$\|g\|_{L^2(\Omega)}^2 \leq C \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)} \|g\|_{L^\infty((1, \infty))} + C \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}^2. \quad (\text{B.2})$$

The estimate (B.2) combined with the standard interpolation inequality

$$\|g\|_{L^\infty((1, \infty))} \leq C \|\partial_r g\|_{L^2((1, \infty))}^{\frac{1}{2}} \|g\|_{L^2((1, \infty))}^{\frac{1}{2}} \leq C \|\partial_r g\|_{L^2(\Omega)}^{\frac{1}{2}} \|g\|_{L^2(\Omega)}^{\frac{1}{2}}$$

yields

$$\|g\|_{L^2(\Omega)}^2 \leq C \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}^{\frac{4}{3}} \|\partial_r g\|_{L^2(\Omega)}^{\frac{2}{3}} + C \left\| \frac{\sqrt{r^2-1}}{r} g \right\|_{L^2(\Omega)}^2.$$

The proof of (3.28) is complete.

Acknowledgement

The third author is partially supported by JSPS Program for Advancing Strategic International Networks to Accelerate the Circulation of Talented Researchers, 'Development of Concentrated Mathematical Center Linking to Wisdom of the Next Generation', which is organized by Mathematical Institute of Tohoku University. The second and third authors are grateful to Université Paris Diderot for their kind hospitality during their stay in the spring semester of 2017.

References

- [1] Abramowitz, M. and Stegun, I., *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, **55**, 1964, xiv+1046 pp.
- [2] Borchers, W., Zur Stabilität und Faktorisierungsmethode für die Navier-Stokes Gleichungen inkompressibler viskoser Flüssigkeiten, *Habilitationsschrift, Universität Paderborn* (1992).
- [3] Farwig, R., Galdi, G. P. , and Kyed, M., Asymptotic structure of a Leray solution to the Navier-Stokes flow around a rotating body, *Pacific J. Math.*, **253** (2011), 367–382.
- [4] Farwig, R. and Hishida, T., Stationary Navier-Stokes flow around a rotating obstacle, *Funkcial. Ekvac.*, **50** (2007), 371–403.
- [5] Farwig, R. and Hishida, T., Asymptotic profile of steady Stokes flow around a rotating obstacle, *Manuscripta Math.*, **136** (2011), 315–338.
- [6] Farwig, R. and Hishida, T., Leading term at infinity of steady Navier-Stokes flow around a rotating obstacle, *Math. Nachr.*, **284** (2011), 2065–2077.
- [7] Farwig, R. and Neustupa, J., On the spectrum of a Stokes-type operator arising from flow around a rotating body, *Manuscripta Math.*, **122** (2007), 419–437.
- [8] Galdi, G. P., Steady flow of a Navier-Stokes fluid around a rotating obstacle, *J. Elast.*, **71** (2003), 1–31.
- [9] Galdi, G. P., An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Steady-State Problems. Second Edition, Springer, (2011).
- [10] Galdi, G. P. and Silvestre, A. L., Strong solutions to the Navier-Stokes equations around a rotating obstacle, *Arch. Ration. Mech. Anal.*, **176** (2005), 331–350.
- [11] Geissert, M., Heck, H., and Hieber, M., L^p -theory of the Navier-Stokes flow in the exterior of a moving or rotating obstacle, *J. Reine Angew. Math.*, **596** (2006), 45–62.
- [12] Higaki, M., Maekawa, Y., and Nakahara, Y., On stationary Navier-Stokes flows around a rotating obstacle in two-dimensions, preprint (2017), arXiv:1701.01215v1.
- [13] Higaki, M., Maekawa, Y., and Nakahara, Y., On the two-dimensional steady Navier-Stokes equations related to flows around a rotating obstacle, accepted for publication in RIMS Kokyuroku Bessatsu “Workshop on the Boltzmann Equation, Microlocal Analysis and Related Topics”.
- [14] Hillairet, M., Wittwer, P., On the existence of solutions to the planar exterior Navier-Stokes system. *J. Differential Equations.* **255** (2013), 2996–3019.
- [15] Hishida, T., An existence theorem for the Navier-Stokes flow in the exterior of a rotating obstacle, *Arch. Rational Mech. Anal.*, **150** (1999), 307–348.
- [16] Hishida, T., L^q estimates of weak solutions to the stationary Stokes equations around a rotating body, *J. Math. Soc. Japan.*, **58** (2006), 743–767.

- [17] Hishida, T., Asymptotic structure of steady Stokes flow around a rotating obstacle in two dimensions, *Mathematical Fluid Dynamics, Present and Future*, Springer Proceedings in Mathematics & Statistics, **183** (2016), 95–137.
- [18] Hishida, T. and Shibata, Y., L^p - L^q estimate of the Stokes operator and Navier-Stokes flows in the exterior of a rotating obstacle, *Arch. Ration. Mech. Anal.*, **193** (2009), 339–421.
- [19] Maekawa, Y., On stability of steady circular flows in a two-dimensional exterior disk. *Archive for Rational Mechanics and Analysis*, 225 (2017), 287–374.
- [20] Silvestre, A. L., On the existence of steady flows of a Navier-Stokes liquid around a moving rigid body, *Math. Meth. Appl. Sci.*, **27** (2004), 1399–1409.