

Root architecture and anchorage: what we learned from maritime pine on sandy soils in the Landes forest

<u>Frédéric Danjon</u>¹, Pauline Défossez², Ming Yang², Clément Saint Cast¹², Antoine Danquechin Dorval¹, Céline Meredieu¹ ¹INRA & Université de Bordeaux UMR1202 BIOGECO, Cestas ² ISPA, Villenave d'Ornon, France

Anchorage is a key factor in wind damage by stem leaning/toppling/uprooting



Experiments to determine anchorage parameters

-> <u>Winching</u> = maximum turning moment



-> inventory of storm damage

-> Empirical relationships

But winching a large amount of trees can be tedious and winching is not predictive

What we would need is quality of anchorage **as a function of factors** like species, genotype, nursery conditions (cutting/seed), installation techniques (planting/sowing), soil profile, soil water content, soil preparation

-> explanatory approach:

Root architecture & material properties assessment

<u>Anchorage strength</u> is given by:

Winching / storm damage / stem straightness / biomechanical models

Thanks to explanatory approaches, we may provide anchorage parameters for a range of factors with **extrapolations** at a given age and from age to age soil can be a highly heterogeneous medium + variable
water content -> high variability in root systems

- all is **hidden in the soil,** entangled in roots from other plants.

- tree anchorage failure is a **discrete** phenomenon:



Root architecture and anchorage in trees: what's known ?



Root architecture and anchorage in trees: toppling

Crook, Ennos, Goodmann, Mickovski et al.: deeper tap-rooted species (*Mallotus*, *Larix*, *Pinus*)

taproot: 80% anchorage

sinkers predominant

Toppling



Mallotus, Crook et al. 1997

3-9 laterals, 3.2 cm diameter

very simple root system or simplified description of root architecture

The way we choose to study relationship between architecture and anchorage







Mechanical uprooting

In-depth phenotyping of complex root architectures





In-depth phenotyping of complex root architecture



In-depth phenotyping of complex root architecture







Deep Phenotyping of Coarse Root Architecture in *R. pseudoacacia* Reveals That Tree Root System Plasticity Is Confined within Its Architectural Model

Frédéric Danjon^{1,2}*, Hayfa Khuder³, Alexia Stokes⁴

1 INRA, UMR1202 BIOGECO, Cestas, France, 2 Université de Bordeaux, UMR1202 BIOGECO, Cestas, France, 3 Université de Bordeaux, UMR 5295, Institut de Mécanique et d'Ingénierie - Bordeaux (I2M), Département Génie Civil et Environnemental (GCE), Bordeaux, France, 4 INRA, UMR AMAP (Botanique et bioinformatique de l'architecture des plantes), Montpellier, France

Abstract

This study aims at assessing the influence of slope angle and multi-directional flexing and their interaction on the root architecture of *Robinia pseudoacacia* seedlings, with a particular focus on architectural model and trait plasticity. 36 trees

Pinus Pinaster, podzols: large trees failing by uprooting





Very large acclimation: Selective reinforcement as a function of constraint experienced (tension/flexion)

branched windward roots beyond ZRT

few big roots leeward ZRT

chuck





Large acclimation to soil depth/soil profile



-> soil profile is determined by the micro-topography

<u>ridge</u>



-> large plasticity of root system as a response to available soil depth

Danjon et al., 2005

Pinus Pinaster, architectural scheme





Klaus storm (2009) 48 trees, including 12 toppled 17 cm DBH

Estimation of roots lost during uprooting = yellow - (Danjon et al. 2013)



Danquechin Dorval PhD (2015) // Danjon Dorval Meredieu submitted



Computing mechanical characteristics from morphological data



relative weighted maximum tensile load (solid line - kPa.m-1)



Danquechin Dorval PhD (2015) // Danjon Dorval Meredieu submitted

4-years-old trees / 10 trees per genetic unit



Dominance of drought avoidance characteristics

Unclear results for artificial selection

Turf plug planting: 300 three-years-old trees.

no clustering, clumping of shallow roots, generally good regeneration of taproot.



retrospective approach : what happened at plantation ?

Danquechin Dorval PhD (2015)

Prospective approach: portion of crop trees in stands



half the plots had less than 50% crop trees

large variation in perturbations of architectural model: could not been attributed to sylvicultural practices.

In a sandy soil: influence of soil preparation on root architecture

Strip ploughing leaving a ditch + mapping shallow roots/sinker branching point in situ: roots follow soil surface







Conclusion: potential contribution of explanatory approaches

We have <u>static data</u>, we could quantify <u>factors</u> effect on root architecture

<u>Growth model</u> for interpolation/extrapolation/summarizing knowledge

<u>Anchorage</u> assessed with the biomechanical model (Yang et al. 2014)

Input in wind risk models:

e.g. planting technique: could not be linked to anchorage strength

low stem straightness: higher probability of anchorage failure

