



Genetic diversity and inoculum exchange of *Botrytis cinerea* between tomato greenhouses in northern Algeria

Ahmed Adjebli, Christel Leyronas, Kamel Aissat, Philippe C. Nicot

► To cite this version:

Ahmed Adjebli, Christel Leyronas, Kamel Aissat, Philippe C. Nicot. Genetic diversity and inoculum exchange of *Botrytis cinerea* between tomato greenhouses in northern Algeria. 17. International Botrytis Symposium, Oct 2016, Santa Cruz, Chile. , 2016, Abstract book XVII International Botrytis symposium, 23 - 28 October 2016, Santa cruz - Chile. hal-01594881

HAL Id: hal-01594881

<https://hal.science/hal-01594881>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

P.3.4: GENETIC DIVERSITY AND INOCULUM EXCHANGE OF *Botrytis cinerea* BETWEEN TOMATO GREENHOUSES IN NORTHERN ALGERIA

Ahmed Adjebli^{1*}, Christel Leyronas², Kamel Aissat¹ and Philippe C. Nicot²

¹Laboratoire d'écologie Microbienne, Faculté des Sciences de la Nature et de la Vie, Université Abderrahmane Mira, Bejaia 06000, Algérie;

²INRA, UR407 Pathologie, France.

ahmed.adjebli@univ-bejaia.dz

To estimate the genetic diversity for a better understanding of the spread of *Botrytis cinerea*, we genotyped with nine microsatellite markers 174 isolates collected from four greenhouses during three growing seasons in the region of Bejaia. Four of these isolates were detected as *Botrytis pseudocinerea* according to the allele size at locus Bc6. For all other isolates further studied, all loci were polymorphic, with the mean number of alleles per locus ranging from 2.77 to 5.22. Considerable genetic variability was detected in all subpopulations ($D^* > 0.87$; $H_{nb} > 0.40$). Based on the standardized index of association analysis, significant but low levels of clonality occurred, not excluding the possibility of recombination ($r_D = 0.07$, $P < 0.001$). A total of 109 haplotypes were characterized among the isolates, few of which were shared between subpopulations. This, together with moderate genetic differentiation among subpopulations according to the geographical origin ($0.080 < F_{ST} < 0.167$), suggested a low level of inoculum exchange among greenhouses and little carry-over of inoculum from one sampling season to the next. The importance of genetic structure of *B. cinerea* populations is discussed and should be taken into consideration for the management of grey mould.



ABSTRACT BOOK

***XVII INTERNATIONAL
BOTRYTIS
SYMPOSIUM***

***23 - 28 OCTOBER, 2016,
SANTA CRUZ • CHILE***

