



HAL
open science

Alternative vector control methods to manage the Zika virus outbreak: more haste, less speed

Jérémy Bouyer, Fabrice Chandre, Jérémie Gilles, Thierry Baldet

► To cite this version:

Jérémy Bouyer, Fabrice Chandre, Jérémie Gilles, Thierry Baldet. Alternative vector control methods to manage the Zika virus outbreak: more haste, less speed. *The Lancet global health*, 2016, 4 (6), pp.E364-E364. 10.1016/S2214-109X(16)00082-6 . hal-01594951

HAL Id: hal-01594951

<https://hal.science/hal-01594951>

Submitted on 26 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Alternative vector control methods to manage the Zika virus outbreak: more haste, less speed

As stated by Laith Yakob and Thomas Walker,¹ the need for efficient and novel mosquito control methods in the context of the Zika virus outbreak is undisputed. However, technologies such as Release of Insects carrying Dominant Lethal genes (RIDL) are not flawless. Transgenic *Aedes aegypti* larvae die if they are not fed with a diet supplemented with an antibiotic (tetracycline). However, deprivation of the mosquitoes from the antibiotic results in 5% survival of transgenic larvae,² resulting in the uncontrolled diffusion of the transgene in large-scale deployment in the field. Moreover, tetracycline is widespread in surface waters in countries where the Zika virus is rapidly progressing, including Brazil.³

By contrast, the Sterile Insect Technique (SIT) is safe and is undergoing field validation against *A. aegypti* and *Aedes albopictus*, but has received little attention to date. SIT does not raise public opposition or require regulatory approvals and no intellectual property issues are involved. Most importantly, the released sterilised insects cannot become established in nature. We believe that the SIT could be further boosted by use of sterile males as conveyors of biocides to their wild counterparts.⁴ A European project, titled Revolutionizing insect control, was launched in 2016 to study the dispersion of densovirus (a species-specific natural entomopathogenic virus)⁵ from sterile males to wild females and then to their larval habitats as a result of their skipping oviposition behaviour (figure). This strategy might enlarge the effect of the SIT, offering unprecedented opportunities for the control of Zika virus and other mosquito-borne diseases.




Figure: Principle of the boosted sterile insect technique using a densovirus

(A) Sterile males of *Aedes aegypti* or *Aedes albopictus* are coated with crystallised densovirus (AeDENV)5 and released by air. (B) Part of the virus is transmitted to the females during mating and the females that are induced sterile carry it to their breeding sites. (C) The virus contaminates larvae in the breeding sites and multiplies with a long-term persistence. (D) Some larvae survive and emerge as contaminated adults that subsequently deliver the larvicide to neighbouring larval habitats.

Although transgenic mosquitoes might be exciting to the media, more biosafe alternatives should not be neglected.

We declare no competing interests. This project has been selected for funding from the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement No 682387—REVOLINC). This Comment reflects only the authors' views and the Agency is not responsible for any use that may be made of the information it contains.

Copyright © Bouyer et al. Open Access article distributed under the terms of CC BY-NC-ND.

**Jérémy Bouyer, Fabrice Chandre, Jérémie Gilles, Thierry Baldet*
bouyer@cirad.fr

Unité Mixte de Recherche Contrôle des Maladies Animales Exotiques et Emergentes, Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), 34398 Montpellier, France (JB, TB); Unité Mixte de Recherche Maladies infectieuses et vecteurs: écologie, génétique, évolution et contrôle, Institut de Recherche pour le Développement, Montpellier, France (FC); and Insect Pest Control Laboratory, Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, Vienna, Austria (JG)

1 Yakob L, Walker T. Zika virus outbreak in the Americas: the need for novel mosquito control methods. *Lancet Glob Health* 2016; **4**: e148–49.

- 2 Curtis Z, Matzen K, Oviedo MN, et al. Assessment of the impact of potential tetracycline exposure on the phenotype of *Aedes aegypti* OX513A: implications for field use. *PLoS Negl Trop Dis* 2015; **9**: e0003999.
- 3 Locatelli MAF, Sodr  FF, Jardim WF. Determination of antibiotics in Brazilian surface waters using liquid chromatography-electrospray tandem mass spectrometry. *Arch Environ Contam Toxicol* 2011; **60**: 385–93.
- 4 Bouyer J, Lefran ois T. Boosting the sterile insect technique to control mosquitoes. *Trends Parasitol* 2014; **30**: 271–73.
- 5 Carlson J, Suchman E, Buchatsky L. Densoviruses for control and genetic manipulation of mosquitoes. *Adv Virus Res* 2006; **68**: 361–92.