Identification of dietary modulators of cognitive function in ageing using metabolomics discovery

Dorrain Low^{1*}, Sophie Lefèvre-Arbogast^{3*}, Melanie Petera², Pierre Micheau¹, Delphine Centeno², Stephanie Durand², Cécile Proust-Lima³, Boris Hejblum³, Mireia Urpi-Sarda⁴, Raul Gonzalez-Dominguez⁴, Cristina Andres-Lacueva⁴, Sandrine Thuret⁵, Cécilia Samieri^{3**}, Claudine Manach^{1**}

¹Human Nutrition Unit, UMR 1019, INRA Auvergne-Rhône-Alpes, Clermont Ferrand, France

dorrain.low-yanwen@inra.fr

With a global rise in ageing population and age-associated diseases, understanding how diet modifies cognitive ageing represents key revenues for prevention. Epidemiology suggests inverse associations between specific dietary patterns (i.e. Mediterranean diet) and cognitive decline and neurodegenerative diseases [1-3]- which raises an important question: which dietary bioactives (i.e. metabolites derived from plant foods) are capable of modulating neuronal ageing?

In this discovery (D-CogPlast⁶) study, we aim to identify a combination of diet-derived metabolites associated with cognitive decline using untargeted metabolomics. We leveraged the Three-City (3C) study (a large French cohort of elderly people) and compared the metabolic profiles of 209 individuals who later developed cognitive decline over 13 years against 209 controls with preserved cognition. Subjects were matched for age at baseline, gender and level of education. Serum samples (collected at the beginning of the cohort study when all participants were cognitively healthy) were profiled using high-resolution UHPLC-QToF (Bruker Impact II) and raw UHPLC-MS data were processed using Galaxy (WorkFlow4Metabolomics.org). Validated PLS-DA clearly distinguished between case and control populations. To account for the matched case-control study design and for potential confounders (i.e. season of blood sampling, body mass index, and number of medications consumed), sparse conditional logistic regression (with bootstrapped re-sampling) was adapted. 17 ions were representative of a serum metabolomic profile associated with cognition. These ions and their clusters whose intensities were significantly elevated in each of the population groups were annotated using online and in-house databases, literature search and commercial standards. Tandem MS/MS fragmentation is presently in progress for further validation of these ions' identification.

Next, the robustness of this set of ions will be validated in a separate cohort of 400 subjects. The ability of these ions predictive of cognitive decline to modulate brain plasticity and neuronal integrity will be further investigated in an *in vitro* parabiosis assay and finally in a proof-of-principle dietary intervention mouse model. It is expected that these identified and validated biomarkers will lead to dietary intervention and recommendations for cognitive decline prevention.

⁶The D-CogPlast study is funded by JPI-HDHL and AgreenSkills+ Young incoming fellowship.

References

- 1. Hardman, R.J., et al., Adherence to a Mediterranean-style diet and effects on cognition in adults: a qualitative evaluation and systemic review of longitudinal and prospective trials. Frontiers in Nutrition, 2016. 3(22): p. 1-13.
- Pelletier, A., et al., Mediterranean diet and preserved brain structural connectivity in older subjects. Alzheimer's & Dementia, 2015. 11(9): p. 1023-1031.
- 3. Singh, B., et al., Association of mediterranean diet with mild cognitive impairment and Alzheimer's disease: a systematic review and meta-analysis. Journal of Alzheimer's Disease, 2014. 39(2): p. 271-282.

²Platforme d'Exploration du Métabolisme, INRA Auvergne-Rhône-Alpes, Saint-Genès-Champanelle, France

³Univ. Bordeaux, Inserm, Bordeaux Population Health Research Center, UMR 1219, F-33000 Bordeaux, France

⁴Nutrition and Food Science department, XaRTA, INSA, Pharmacy and Food Sciences faculty, University of Barcelona, Barcelona, Spain

⁵Department of Basic and Clinical Neuroscience, Maurice Wohl Neuroscience Institute, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

^{*}Both first authors share equal contributions

^{**} Both authors share co-last authorship