



HAL
open science

Functional traits of indigenous and exotic ground-dwelling arthropods show contrasting responses to land-use change in an oceanic island, Terceira, Azores

François Rigal, Pedro Cardoso, Jorge Lobo, Kostas A Triantis, Robert J. Whittaker, Isabel R Amorim, Paulo Alexandre Vieira Borges

► To cite this version:

François Rigal, Pedro Cardoso, Jorge Lobo, Kostas A Triantis, Robert J. Whittaker, et al.. Functional traits of indigenous and exotic ground-dwelling arthropods show contrasting responses to land-use change in an oceanic island, Terceira, Azores. *Diversity and Distributions*, 2018, 24, pp.36-47. 10.1111/ddi.12655 . hal-01596143

HAL Id: hal-01596143

<https://hal.science/hal-01596143>

Submitted on 27 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Functional traits of indigenous and exotic ground-dwelling arthropods show**
2 **contrasting responses to land-use change in an oceanic island, Terceira, Azores**

3 François Rigal^{1,2*}, Pedro Cardoso^{1,3}, Jorge M. Lobo⁴, Kostas A. Triantis^{1,5}, Robert J.
4 Whittaker^{6,7}, Isabel R. Amorim¹ and Paulo A.V. Borges¹

5 ¹CE3c – Centre for Ecology, Evolution and Environmental Changes / Azorean
6 Biodiversity Group and Universidade dos Açores - Departamento de Ciências e
7 Engenharia do Ambiente, 9700-042 Angra do Heroísmo, Açores, Portugal

8 ²CNRS-Université de Pau et des Pays de l'Adour, Institut des Sciences Analytiques et
9 de Physico-Chimie pour l'Environnement et les Matériaux, MIRA, Environment and
10 Microbiology Team, UMR 5254, BP 1155, 64013 Pau Cedex, France

11 ³Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland.

12 ⁴Departament of Biogeography and Global Change, Museo Nacional de Ciencias
13 Naturales (CSIC), Madrid, Spain

14 ⁵Department of Ecology and Taxonomy, Faculty of Biology, National and
15 Kapodistrian University of Athens, Athens GR-15784, Greece

16 ⁶Conservation Biogeography and Macroecology Programme, School of Geography
17 and the Environment, University of Oxford, South Parks Road, Oxford, OX1 3QY,
18 United Kingdom

19 ⁷Center for Macroecology, Evolution and Climate, Department of Biology, University
20 of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen Ø, Denmark

21

22 *Corresponding author: François Rigal, email: francois.rigal@univ-pau.fr

23

24 **Keywords:** arthropods, Azores, community assembly, functional diversity, exotic
25 species, indigenous species

26 **Running head:** Functional traits of indigenous and exotic arthropods

27 **ABSTRACT**

28 **Aim** Land-use change typically goes hand-in-hand with the introduction of exotic
29 species, which mingle with indigenous species to form novel assemblages. Here we
30 compare the functional structure of indigenous and exotic elements of ground-
31 dwelling arthropod assemblages across four land-uses of varying management
32 intensity.

33 **Location** Terceira Island (Azores, North Atlantic).

34 **Methods** We used pitfall traps to sample arthropods in 36 sites across the four land-
35 uses and collated traits related to dispersal ability, body size and resource-use. For
36 both indigenous and exotic species, we examined the impact of land-uses on trait
37 diversity and tested for the existence of non-random assembly processes using null
38 models. We analysed differences in trait composition among land-uses for both
39 indigenous and exotic species with multivariate analyses. We used point-biserial
40 correlations to identify traits significantly correlated with specific land-uses for each
41 element.

42 **Results** We recorded 86 indigenous and 116 exotic arthropod species. Under high
43 intensity land-use, both indigenous and exotic elements showed significant trait
44 clustering. Trait composition strongly shifted across land-uses, with indigenous and
45 exotic species being functionally dissimilar in all land-uses. Large-bodied herbivores
46 dominated exotic species in low intensity land-uses, while small-bodied spiders
47 dominated exotics in high intensity land-uses. In contrast, with increasing land-use
48 intensity, indigenous species changed from functionally diverse to being dominated
49 by piercing and cutting herbivores.

50 **Main conclusions** Our study revealed two main findings: first, in high intensity land-
51 uses, trait clustering characterized both indigenous and exotic elements; second,

52 exotic species differed in their functional profile from indigenous species in all land-
53 use types. Overall, our results provide new insights into the functional role of exotic
54 species in a land-use context, suggesting that, in agricultural landscape, exotic species
55 may contribute positively to the maintenance of some ecosystem functions.
56

57 **INTRODUCTION**

58 Land-use change is inducing profound changes in biodiversity patterns throughout the
59 world (Mooney & Hobbs, 2000; Newbold et al., 2015). Different species may be
60 more or less sensitive to particular forms of disturbance, depending on their traits
61 (Flynn et al., 2009; Mayfield et al., 2010). Because species' traits influence
62 community assembly processes and ecosystem functioning, special attention has been
63 given to the consequences of the loss of indigenous species for functional diversity
64 (FD) (Diaz, Fargione, Chapin & Tilman, 2006; Flynn et al., 2009; Brown et al., 2013;
65 Forrest, Thorp, Kremen & Williams, 2015). Land-use changes frequently promote the
66 establishment of exotic species, often generating an increase in overall species
67 richness at particular scales of analysis (Sax, Gaines & Brown, 2002). However, the
68 role of exotic species in driving changes in FD in the context of land-use changes
69 remains understudied (Vilà et al., 2011). Particularly, it remains unclear how land-use
70 change shapes functional trait structure of exotic species and to what extent exotic
71 species could contribute to the provision and maintenance of ecosystem functions in
72 agricultural landscape (Rodgers & Parker, 2003; Didham, Tylianakis, Gemmill, Rand
73 & Ewers, 2007; Hobbs, Sayre & Gupta, 2008).

74 Many exotic species are known to be particularly successful in anthropogenic
75 habitats. Exotic species can possess functional traits that are not present in the
76 indigenous community, allowing them to occupy new ecological opportunities
77 provided by land-conversion and transformation (Mayfield & Levine, 2010). In
78 addition, some exotic species can possess similar functions to indigenous species and
79 may provide ecological “insurance” after the decline of less tolerant indigenous
80 species following land-use changes (e.g. Aslan, Zavaleta, Tershy, Croll & Robichaux,
81 2012). As land-use change intensifies, both indigenous and exotic species may,

82 however, be filtered out towards a restricted set of functional strategies due to
83 stressful environmental conditions (Tecco, Díaz, Cabido & Urcelay, 2010). This may
84 lead to functional homogenization whereby exotic and indigenous species fulfill
85 similar functional roles (Olden & Rooney, 2006; Drenovsky et al., 2012).

86 Arthropods are the most diverse group of organisms and form also a great part of the
87 exotic fauna worldwide (Kenis et al., 2009). Several studies have reported lower
88 indigenous arthropod species richness and abundance in agricultural land-uses than in
89 more natural habitats (Hendrickx et al., 2007, Attwood, Maron, House & Zammit,
90 2008), while exotic arthropods generally benefit from agricultural landscapes (Grez,
91 Rand, Zaviezo & Castillo-Serey, 2013). Although arthropods provide numerous
92 ecosystem services, including pollination, regulation of herbivorous pests and
93 decomposition (Losey & Vaughn, 2006; Isaacs, Tuell, Fiedler, Gardiner & Landis,
94 2009), there has been relatively little trait-based analysis of these functions compared
95 to plants or vertebrates (Kenis et al., 2009; Diekotter, Wamser, Wolters & Birkhofer,
96 2010). Moreover, we are unaware of any studies examining how traits of indigenous
97 and exotic arthropods change in communities sampled across contrasting land-use
98 types.

99 In this study, we evaluate and compare the functional trait structure of indigenous and
100 exotic ground-dwelling arthropods across four land-use types of varying management
101 intensity in the oceanic island of Terceira (Azores, Portugal). The four land-uses
102 range from pristine native forest patches subject to very low human management to
103 intensively exploited pastures subject to frequent cattle grazing and fertilisation
104 (Borges et al., 2006; Cardoso et al., 2009). Indigenous and exotic species have been
105 shown to co-occur at the local scale in all four land-uses, permitting comparisons of
106 trait assembly between indigenous and exotic elements of the same assemblages in

107 relation to these contrasting land-uses (Cardoso et al., 2009). Here, we focused on
108 body size, dispersal ability and resource-use traits, all of which are recognized as
109 important ecological traits, especially in a land-use context (Schweiger et al., 2005,
110 Simons, Weisser & Gossner, 2016).

111 Specifically, we first hypothesized that both indigenous and exotic species, analysed
112 separately, should show increased trait clustering in high intensity land-uses (H1).
113 The rationale is that only species with a specific set of traits should persist under
114 intensive and frequent environmental disturbances (i.e. cattle grazing and fertilization)
115 (Flynn et al., 2009). Our second hypothesis was that exotic species should show trait
116 dissimilarities when compared to co-occurring indigenous species under low and high
117 intensity land-uses (H2). The rationale is that, under low intensity land-uses (e.g.
118 native forest), exotic species exhibiting novel functional traits are favoured in
119 exploiting the vacant niches offered by the non-saturated nature of insular and/or
120 impoverished indigenous assemblages in Terceira (Whittaker et al., 2014). In contrast,
121 under high-intensity land-uses, exotic species have a competitive advantage for
122 resource exploitation and establishment space over the indigenous species (Grigulis,
123 Sheppard, Ash & Groves, 2001; Saar, de Bello, Pärtel & Helm, 2017).

124 This study expands on previous works on the impact of land-use change on arthropod
125 assemblages in the Azores (Borges, Ugland, Dinis & Gaspar, 2008; Cardoso et al.,
126 2009; Meijer, Whittaker & Borges, 2011; Florencio, Cardoso, Lobo, Azevedo &
127 Borges, 2013; Florencio, Lobo, Cardoso, Almeida-Neto & Borges, 2015; Florencio et
128 al. 2016; Rigal, Whittaker, Triantis & Borges, 2013), and by focusing on traits allows
129 for stronger inferences on the functional processes that might control the respective
130 role of indigenous and exotic species in community assembly in the Azores.

131

132 **METHODS**

133 **Study area**

134 The Azores (37° to 40° N; 25° to 31° W) is one of the world's most isolated
135 archipelagos. There are nine main islands aligned on a WNW–ESE axis in the
136 Atlantic Ocean. All islands are volcanic and of recent origin. At the time of human
137 colonization, around AD 1440, the Azorean archipelago was almost entirely forest-
138 covered. By c. AD 1850, human activities had restricted the native forest to areas
139 above 500 m a.s.l.. During the 20th century, the development of an economy based on
140 dairy has led to a further reduction of native forest area, with the clearing of large
141 fragments at mid- and high-elevations for pasture development. In addition, the
142 plantation of forests of exotic species was incentivized during the 1950s in areas
143 unsuitable for cattle grazing, thus promoting additional deforestation of natural forest
144 habitat. Thus, in < 600 years, 95% of the original native forest has been destroyed
145 (Gaspar, Borges & Gaston, 2008).

146 Our study was conducted on Terceira Island (402 km², 3.52 Ma), on which the native
147 forest is restricted to 23 km², about 6% of its original area (Triantis et al., 2010 and
148 references herein). The non-urban landscape of Terceira is dominated by four major
149 habitats: (i) native forests, (ii) exotic forests (*Cryptomeria japonica* and *Eucalyptus*
150 *globulus* monoculture plantations, monocultures and mixed forest of the invasive
151 *Pittosporum undulatum*); (iii) semi-natural pastures (mid- and high- elevation pastures
152 maintaining some indigenous plants) subject to short-term cattle grazing activity and
153 low inputs of fertilizers and (iv) intensively managed pastures subject to intensive
154 cattle grazing and high inputs of fertilizers (See Appendix S1 and Table S1.1 in
155 Supporting Information for detailed features of the land-uses). The four land-uses
156 were ranked in an increasing order, according to their intensity uses with (1) native

157 forest, (2) exotic forest, (3) semi-natural pasture and (4) intensively managed pasture.
158 Exotic forests were considered as the closest to the native forest since, contrary to
159 pastures, they maintain some of the architectural and microclimatic conditions of the
160 natural forests (Cardoso et al., 2009, Cardoso, Rigal, Fattorini, Terzopoulou &
161 Borges, 2013).

162

163 **Sampling methodology**

164 Nine sampling sites were established in each of the four land-use types (total = 36),
165 distributed across the island without regard to the character of the surrounding land-
166 use matrix. Sampling sites were on average 1.6 Km (range 0.137–3.7) apart from each
167 other and 0.73 Km apart (0.13–3.8) from the edge of the sampled land-use type (Table
168 S1.2 and Fig. S1.1). The ground-dwelling fauna was sampled at each site using 30
169 pitfall traps, spaced 5 m apart, along a 150 m transect. Traps were left in the field for
170 two weeks in the summer months of June, July or September of different years (native
171 forests between 1999 and 2003, and the other habitats in 2008 and 2009). A potential
172 effect of the sampling years on species compositions across sites were discarded by a
173 between-year analysis using additional data (Florencio et al., 2013). The majority of
174 captured arthropods (excluding Crustacea, Collembola, Diplura, Diptera and
175 Hymenoptera) were sorted to species level; the remaining specimens were sorted to
176 morphospecies. All species were classified as indigenous or exotic. Indigenous
177 species comprise Azorean endemics and other native non-endemics. Exotic species
178 are those considered to have colonized via human mediation, many of which having a
179 cosmopolitan distribution (Borges et al., 2010 and see Appendix S1 for further
180 details in species classification).

181 All analyses were conducted using the transect (i.e. set of 30 pitfall traps) as the unit
182 of analysis.

183

184 **Traits characterization**

185 For all arthropod species, we collated body size, dispersal abilities and a set of
186 functional traits related to resource-use (Table 1). Apart from body size, which was
187 measured on the individuals sampled in this study, traits were collected from an
188 extensive literature search, including manuscripts with the first descriptions of the
189 species, first species records for the Azores, brief notes, and ecological studies.
190 Information was also obtained from experts who have identified the specimens or
191 from experts of a given taxonomic group when information for a particular species
192 was not available. Most of the literature was retrieved from the taxonomic catalogue
193 of the entomological bibliography for the Azores (See Vieira & Borges, 1993 and
194 Borges & Vieira, 1994), with the addition of some recent documentation (Barnard,
195 2011). Functional information was assigned to each species according to the adult
196 characteristics, except for Lepidoptera, where traits were assigned by reference to the
197 larvae. For the unidentified morphospecies, we assigned functional traits of the
198 nearest taxonomic resolution (genus, family), except for body size. Species trait
199 assignments, together with the species mean abundances per land-use are given in
200 Appendix S2.

201

202 **Data analysis**

203 Our first set of analyses aimed to test whether indigenous and exotic elements of
204 ground-dwelling arthropod assemblages each show trait-clustering in high land-use
205 intensity (H1). Therefore, unless otherwise stated, all the following analyses were

206 performed independently for indigenous and exotic species. To provide context for
207 our trait-based analyses, we first quantify species diversity in each site using species
208 richness and the Simpson index. Due to differences in sampling completeness
209 between sites among the land-uses (See Appendix S3 and Fig. S3.1), we calculated
210 rarefied richness to the lowest sample size ($n = 77$ individuals) to perform direct
211 comparisons of richness between our four land-uses. Prior to our FD analysis, we
212 built a functional dissimilarity matrix between species using the extension of the
213 Gower's distance metric provided by Pavoine, Vallet, Dufour, Gachet & Daniel
214 (2009). To assess clustering of species in trait space per site, we computed the mean
215 pairwise functional distance (MFD), i.e. the averaged functional distance between all
216 pairs of species co-occurring in a site (Webb, Ackerly, McPeck & Donoghue, 2002).
217 To account for abundance distributions within sites and to lower the weight of rare
218 and possibly 'vagrant' species, the distance values were weighted according to the
219 log-transformed species abundance. By construction, MFD is hardly, if at all,
220 sensitive to species richness (Pavoine, Gasc, Bonsall & Mason, 2013). To test
221 whether rarefied species richness, the Simpson index and MPD differ between land-
222 uses, we applied generalized least square models (GLS; Pinheiro & Bates, 2000) to
223 account for unequal variances among land-uses. We also tested the ability of the GLS
224 models to account for potential spatial structure by estimating the Moran's I spatial
225 autocorrelation index on residuals using latitude and longitude of each sample site.
226 We quantified the effect of land-uses by using Magee's pseudo- R^2 (Magee, 1990).
227 When the overall GLS model was statistically significant, the Tukey's *post hoc* test
228 was used to identify statistically significant pairwise differences between land-uses.
229 To test if the MFD was larger (functionally overdispersed) or smaller (functionally
230 clustered) than expected by chance, we compared the observed MFD value in each

231 site to a random distribution under the null hypothesis of no association between land-
 232 uses and traits (Cornwell & Ackerly, 2009). We then generated 999 null assemblages
 233 using the Independent Swap algorithm (Gotelli & Graves, 1996), which reshuffles the
 234 species-by-sites matrices keeping species richness, species frequency and the total
 235 abundance of species in the matrix constant. To measure deviation from the null
 236 expectation for each site, we computed the standardized effect size (SES) using the
 237 formula $SES_{MFD} = (MFD_{obs} - \mu_{null})/\sigma_{null}$ where MFD_{obs} is the observed MFD value at a
 238 given site, μ_{null} , the mean MFD of the simulated communities, and σ_{null} the associated
 239 standard deviation. Positive SES_{MFD} indicate that species occurring together in a site
 240 are less functionally related than expected by chance (functionally overdispersed)
 241 while negative SES_{MFD} indicate that co-occurring species are more functionally
 242 related than expected by chance (functionally clustered). We tested whether SES_{MFD}
 243 values for a given land-use presented significant deviations from the null expectation
 244 (median=0) using a one-sample Wilcoxon test (Swenson & Enquist, 2009). We also
 245 computed GLS for SES_{MFD} in the same way as specified for diversity metrics.

246 Our second set of analyses aimed to test whether exotic and indigenous elements of
 247 single ground-dwelling arthropods assemblages differ in their trait composition
 248 among land-uses (H2). For each individual site, we began by calculating the
 249 community-weighted means (CWM) for body size and each trait attribute for both
 250 indigenous and exotic species separately. The CWMs express the mean attribute value
 251 between species occurring at a site, weighted by the relative abundance of each
 252 species (Violle et al., 2007) and was calculated as follows:

253
$$CWM_j = \sum_{i=1}^S p_i x_{ij}$$

254 where CWM_j is the community-weighted mean value of trait attribute j , p_i is the
255 relative abundance of species i ($i = 1, 2, \dots, S$) and x_{ij} is the value of trait attribute j
256 for species i . Prior to CWM calculation, multi-choice nominal traits and nominal traits
257 were dummy-transformed to as many binary variables as there were trait attributes
258 and were handled as quantitative variables. This generated new “CWM-by-sites”
259 matrices separately for indigenous and exotic species. Based on these matrices, we
260 calculated the Bray-Curtis distance between all pairs of sites and conducted a non-
261 parametric permutational multivariate analysis of variance (PerMANOVA; Anderson,
262 2001) to test for difference in trait attributes composition among land-uses.
263 PerMANOVA was performed using permutation tests with 9999 iterations. When the
264 overall PerMANOVA was significant, *post-hoc t*-tests were performed to identify
265 statistically significant pairwise differences between land-uses (Anderson, Gorley &
266 Clarke, 2008). To test for differences between indigenous and exotic species among
267 land-uses, we computed a PerMANOVA to specifically test the interaction between
268 species origin (indigenous vs. exotic) and land-uses using a Bray-Curtis distance
269 matrix computed by combining the indigenous and exotics “CWM-by-sites” matrices.
270 Site was set as a random effect. When the interaction term was significant, *post-hoc t*-
271 tests were performed to identify land-uses where trait compositions of indigenous and
272 exotic species were significantly different from each other. Results of the
273 PerMANOVAs were visualized using non-metric multidimensional scaling
274 ordinations (NMDS) based on Bray-Curtis distance matrices.
275 Finally, to help the interpretation of our functional trait-based patterns, we conducted
276 an indicator analysis (Dufrêne & Legendre, 1997) to identify the degree of
277 correlations between trait attributes and land-uses for both indigenous and exotic
278 species. We then used the point-biserial correlation coefficient (r_{pb} , De Caceres &

279 Legendre, 2009) to test the correlation between each CWM and the four land-use
280 separately using a binary coding for land-uses, with 1 and 0 indicating whether sites
281 belong or not to the land-use under study. We calculated 95% confidence intervals for
282 r_{pb} values by bootstrapping the observed data 9999 times and considered values not
283 overlapping zero as significant.

284 Statistical analyses were implemented within the R programming environment (R
285 Development Core Team 2014) using the packages *vegan* (Oksanen et al., 2013),
286 *picante* (Kembel et al., 2010), *nlme* (Pinheiro, Bates, DebRoy & Sarkar, 2012),
287 *indicspecies* (De Caceres & Legendre, 2009) and *BAT* (Cardoso, Rigal & Carvalho,
288 2015). PERMANOVAs were performed using the software PRIMER v.6 (Anderson
289 et al., 2008).

290

291 **RESULTS**

292 A total of 20,800 soil-arthropod specimens were collected and 202 (morpho) species
293 were identified, representing 20 orders, 76 families and 161 genera. Of the 202
294 species, 86 were considered to be indigenous and 116 to be exotics (Appendix S2).
295 About 13% of the taxa were identified to morphospecies rather than species.

296 No spatial autocorrelation was detected in the residuals of the GLS models (Table
297 S4.1 in Appendix S4). Rarefied indigenous species richness decreased significantly
298 from low to high intensity land-uses (Fig. 1A and Table S4.2), while rarefied exotic
299 species richness increased (Fig. 1B and Table S4.2). Similar trends were observed for
300 the Simpson's index (See Fig. S4.1A, B and Table S4.2). MFD of indigenous species
301 decreased significantly from low to high intensity land-uses (Fig. 1C and Table S4.2).
302 For exotic species, MFD of semi-natural pastures was significantly lower than for
303 other land-use types but no other significant differences occurred (Fig. 1D and Table
304 S4.2). Both indigenous and exotic elements were significantly functionally

305 overdispersed in native forest (median $SES_{MFD} > 0$; Wilcoxon test, $P=0.003$ and $P =$
306 0.007 respectively, Fig. 1E, F). Indigenous species were significantly functionally
307 clustered in intensively managed pastures (median $SES_{MFD} < 0$; Wilcoxon test,
308 $P=0.003$, Fig. 1E) but not in semi-natural pastures (Wilcoxon test, $P=0.07$, Fig. 1E)
309 while exotic species were significantly functionally clustered in both semi-natural
310 pastures and intensively managed pastures (median $SES_{MFD} < 0$ Wilcoxon test, $P =$
311 0.003 and $P = 0.003$ respectively, Fig. 1F). Neither indigenous nor exotic species
312 showed significant deviation from the null expectation in exotic forest samples
313 (Wilcoxon test, $P = 0.73$ and $P = 0.12$ respectively)

314 Trait composition significantly differed between land-uses for both indigenous and
315 exotic species (PERMANOVA: $R^2 = 0.44$, pseudo- $F_{3,32} = 8.518$, $P < 0.001$ and $R^2 =$
316 0.76 , pseudo- $F_{3,32} = 33.741$, $P < 0.001$ respectively). Moreover, *post-hoc t* tests
317 revealed that land-uses were significantly different from each other in their trait
318 composition for both indigenous and exotic species ($P < 0.05$, Table S4.3), except
319 between semi-natural and intensively managed pastures for indigenous species ($P =$
320 0.056 , Table S4.3 and see Fig. S4.2A and B for the NMDS plots for indigenous and
321 exotic species, respectively). Analyses combining both indigenous and exotic
322 “CWM-by-sites” matrices showed that trait composition differed between indigenous
323 and exotic species among land-use types (PERMANOVA, Interaction species origin x
324 land-uses: $R^2 = 0.324$, pseudo- $F_{3,64} = 28.317$, $P < 0.001$). Moreover, *post-hoc t* tests
325 revealed that indigenous and exotic species were significantly different from each
326 other in their trait composition within all land-uses ($P < 0.05$, Table S4.3). This pattern
327 was clearly evident in the NMDS plot (Fig. 2) showing a clear absence of overlap for
328 each land-use between indigenous and exotic species in the two-dimensional space.

329 The point-biserial correlations (r_{pb}) analysis between trait attributes (i.e. CWMs) and
330 land-uses indicated striking differences between indigenous and exotic species (Fig.
331 3). For instance, the trait attributes plants (Trait: Type of food), detritus (Type of
332 food), chewing and cutting (Mode of ingestion), high active search (Way of getting
333 food) and body size showed significantly negative correlations with native forest for
334 indigenous species while significantly positive correlations were found for exotic
335 species. In contrast, the attribute plants (Type of food) and piercing and sucking
336 (Mode of ingestion) showed significantly positive correlations with intensive
337 managed pastures for indigenous species and significantly negative correlations for
338 exotic species, while the attributes animals (Type of food) and intestinal digestion and
339 sucking (Mode of ingestion) showed the opposite pattern. For exotic species, body
340 size was also negatively correlated with intensively managed pastures. Only the
341 attribute high dispersal ability exhibited a consistent pattern for both indigenous and
342 exotic species, with significantly negative correlations in native and exotic forests and
343 significantly positive correlations in intensive managed pastures.

344

345 **DISCUSSION**

346 Our study revealed two main findings: 1) in high intensity land-use (semi-natural and
347 intensively-managed pastures), trait clustering characterized both indigenous and
348 exotic elements (supporting H1) and 2) indigenous and exotic species were filtered
349 out towards two distinct functional profiles across the four land-use types with both
350 species origin differing in their trait composition in all land-uses (supporting H2).

351

352 **Prevalence of trait clustering for both indigenous and exotic species in high**
353 **intensity land-uses**

354 Both indigenous and exotic elements were found to display significant trait-clustering
355 in high intensity land-uses, highlighting the fact that species were functionally more
356 similar than expected by chance in those land-uses. Overall, these findings are in
357 accordance with recent trait-based studies claiming that land-use intensification acts
358 as a filter, favouring a limited array of trait combinations (Flynn et al., 2009; Laliberté
359 et al., 2011). Specifically, for arthropods, our results confirm recent observations
360 made in a land-use change context for insect pollinators (Rader, Bartomeus,
361 Tylianakis & Laliberté, 2014), beetles (Birkhofer, Smith, Weisser, Wolters &
362 Gossner, 2015), bees (Forrest et al., 2015) and arthropods as a whole (Simons et al.,
363 2016).

364 For indigenous species, MFD values lower than expected by chance in intensively
365 managed land-uses (Fig. 1E) were coupled with a decline of species diversity,
366 indicating that functions were lost at a higher rate than would be predicted by chance
367 if species were lost randomly (Flynn et al., 2009). In contrast, exotic species richness
368 increased as land-use intensity increased, but this was accompanied by lower than
369 expected MFD values under high intensity land-uses (Fig. 1F). These results reveal a
370 degree of apparent functional redundancy within the exotic element of ground-
371 dwelling arthropod assemblages under high intensity land-uses, with more species
372 executing similar functions.

373 Interestingly, we found significant trait divergence in native forests for both
374 indigenous and exotic species (Fig. 1E-F). Although this pattern is usually interpreted
375 as the result of biotic interactions, typically competitive exclusion (Webb et al., 2002),
376 caution is needed with such an interpretation (de Bello, 2012). Since for the null
377 model analysis, we considered all the species found across the four land-uses as our
378 regional species pool, it is more likely that species distribution across these land-uses

379 may result predominantly from environmental constraints overpowering the
380 importance of local species interactions (Münkemüller et al., 2013). It is possible that
381 different outcomes would be produced if we down-scaled the species pool to a
382 particular land-use (Münkemüller et al., 2013). Moreover, capturing evidence of
383 competitive exclusion would require us to focus on a specific feeding guild (e.g.,
384 competition between predators), whereas our study considers all ecological guilds and
385 trophic levels of ground-dwelling arthropods. Therefore, the functional overdispersion
386 observed herein might simply reflect the large range of niches occupied by species in
387 native forest comparatively to the other land-uses.

388

389 **Land-use change filtered out indigenous and exotic species towards distinct**
390 **functional profiles**

391 We found that trait composition of both indigenous and exotic species differed
392 between land-uses, reflecting a clear shift in the functional space occupation from low
393 to high land-use intensity. Moreover, our study reveals that the direction of the effect
394 of land-use change on the trait composition greatly differed between indigenous and
395 exotic species.

396 Indigenous species shift from functionally diverse elements in low intensity land-use
397 towards elements dominated mainly by piercing and sucking herbivores in high
398 intensity pasture land-uses (Fig. 3). Examples of the species involved include the
399 coleopterans *Hirticollis quadriguttatus* (Anthicidae, native) and *Heteroderes azoricus*
400 (Elateridae, Azorean endemic) (Appendix S2). This finding is consistent with
401 previous studies on Azorean herbivores that have reported a high proportion of
402 generalists in free-living phytophagous insects (Ribeiro et al., 2005). Such patterns
403 have been observed for other oceanic island guilds of phytophagous insects (Olesen,

404 Eskildsen & Venkatasamy, 2002; Kaiser-Bunbury, Traveset & Hansen, 2010) and are
405 usually explained as reflecting the relaxation of interspecific competition as
406 consequence of the young age and the isolation of such islands (Olesen et al., 2002).
407 An alternative and non-exclusive explanation is that some herbivorous species can
408 increase in abundance with habitat loss due to a reduction in specialist predator
409 populations (a top-down effect) (Ryall & Fahrig, 2006). The directional shift towards
410 herbivore-dominated elements was accompanied by a correspondingly low relative
411 abundance of carnivorous species in intensively-managed pastures (Fig. 3). This is in
412 accordance with several previous studies that have reported that high trophic-level
413 arthropod taxa were relatively sensitive to environmental stress (e.g. Holt, 1996,
414 Pearce & Venier, 2006). Web-building spiders in particular, often being dependent on
415 substantial habitat structure and vegetation architecture to anchor their webs (e.g.
416 Samu, Sunderland & Szinetar, 1999), cannot find suitable sites when clear-cutting or
417 establishment of pastures destroy all trees. Also, some Chilopoda species are known
418 to display strong preferences for low temperatures and high humidity (conditions
419 encountered in native and exotic forest). Examples include species of *Lithobius*
420 (Lewis, 2006), one of which, *Lithobius pilicornis pilicornis* (Lithobiidae; native), was
421 found to be abundant in forest but almost absent in pastures (Appendix S2). From a
422 bottom-up perspective, predator populations could have also been indirectly and
423 negatively affected by the decline of their prey populations (Ryall & Fahrig, 2006).
424 We also found a significant correlation between high dispersal ability and high
425 intensity land-uses for indigenous species (Fig. 3). This suggests the hypothesis that
426 pastures act as sinks for forest-associated species, with only the good dispersers being
427 able to colonize them from native forest patches (Borges et al, 2008). Native forest is
428 particularly characterised by small-bodied indigenous species, reflecting the strong

429 dispersal filter imposed by the remote location of the archipelago. This is particular
430 apparent for indigenous spiders, which have mainly reached the islands by ballooning
431 (Carvalho & Cardoso, 2014)

432 In contrast to indigenous species, the trait composition of exotic elements shifted from
433 a preponderance of large-bodied, chewing and cutting herbivores and detritivores in
434 low-intensity land-uses, towards elements dominated mainly by smaller-bodied
435 species, active throughout both day and night, carnivorous, with extra-intestinal
436 digestion and highly dispersive (mostly spider species) in high land-use intensity (Fig.
437 3). Native forest provides abundant resources for herbivorous species, thereby
438 allowing the successful establishment of exotic herbivores. Moreover, since
439 indigenous Azorean communities are largely unsaturated (Borges & Brown, 2004;
440 Whittaker et al., 2014), competition for food is likely to be reduced. Interestingly, the
441 most successful exotic herbivores were two Julidae species, *Ommatoiulus moreletii*
442 and *Cylindroiulus propinquus*, which possess the largest body size of all arthropods
443 found in Terceira Island (Appendix S2). Considering that ecological niches for large
444 species have not yet been filled by indigenous species (above), large-sized exotic
445 species may have been favoured in establishing populations in native forests. These
446 two Julidae species are also occasionally detritivores (Bailey & Mendonça, 1990),
447 explaining the dominance of this trait within exotic elements in native forest.

448 The directional shift within exotic elements towards small-sized spiders in sites of
449 high intensity land-use was accompanied by a clear pattern of functional redundancy
450 (Fig. 1F). In both semi-natural and intensively managed pastures, small-sized
451 Linyphiidae spiders featured prominently, especially members of the genera *Erigone*
452 (*E. autumnalis* and *E. dentipalpis*) and *Oedothorax* (*O. fuscus*), both well adapted to
453 grazed pastureland (Wise, 1995 and Appendix S2). Linyphiidae in general, and those

454 genera in particular, are mostly *r*-selected species, which are tiny (≈ 2 mm), efficient
455 dispersers, often with multiple generations per year (Bell, Wheeler & Cullen, 2001).
456 These features may therefore confer strong advantages in the re-colonization of
457 pasture lands after intensive grazing and mowing (Tscharntke, Klein, Kruess, Steffan-
458 Dewenter & Thies, 2005). The dominance of exotic Linyphiids could also be due to
459 the wide availability of prey. However, Linyphiids mainly feed on Collembola,
460 Diptera and Hymenoptera (Agusti et al., 2003), three orders that were not considered
461 in our study.

462

463 **Potential mechanisms involved in functional structure of both indigenous and**
464 **exotic elements**

465 Our study clearly shows that indigenous and exotic species display opposing
466 functional profiles across land-use types. There could be multiple alternative
467 underlying mechanisms for this pattern. Prior to human colonization, the native forest
468 in the archipelago acted as a filter, selecting species that were able to establish, some
469 of which subsequently speciated in the archipelago. Hence the indigenous species
470 pool did not probably include species pre-adapted to pasturelands and in this regard,
471 generalist herbivores may be the exceptions. However, 600 years of human
472 occupation of the Azores have led to the emergence of a large exotic species pool
473 (60% of the current arthropod fauna, Borges et al., 2010), including many species able
474 to cope with both anthropogenic and native habitats (e.g. Cardoso et al., 2013;
475 Florencio et al., 2016). To some extent it is possible that exotic species simply filled
476 the ecological space not occupied by indigenous species. In low land-use intensity,
477 historical factors and/or the extinction of disturbance-sensitive species (i.e.
478 particularly those of comparatively large body size, Terzopoulou, Rigal, Whittaker,

479 Borges & Triantis, 2015) might have led to ecological niche space being vacated,
480 while it is evident that unoccupied spaces in high land-use intensity were made
481 available for exotics due to the inability of indigenous species to persist in them. It is
482 also likely that some indigenous species have been wiped out, especially from high-
483 intensity land-uses due to interactions (competition, predation) with exotic species
484 (Cardoso, Arnedo, Triantis & Borges, 2010). It is also entirely plausible that some
485 indigenous species may have been driven to extinction without ever having been
486 scientifically documented (e.g. Cardoso et al., 2010).

487

488 **CONCLUSION**

489 Although arthropods play important functional roles in both natural and
490 anthropogenic habitats (Losey & Vaughn, 2006; Isaacs et al., 2009), trait-based
491 studies on arthropod assemblages are scarce and, as far as we are aware, our study is
492 the first to formally assess the impact of land-use changes on the functional trait
493 structure of both indigenous and exotic elements of arthropod assemblages.
494 In Terceira Island, the establishment of novel habitats as a result of human activities
495 has led to the formation of novel biotic assemblages in which indigenous and exotic
496 species co-exist. Our results clearly showed that, within these novel assemblages,
497 exotic species differed in their functional profile from indigenous species, suggesting
498 that exotic species might play important roles in providing and maintaining some key
499 ecosystem functions, particularly in high-intensity land-uses. To what degree these
500 assemblages are self-sustaining and stable remains, however, unknown, emphasizing
501 the need for long-term studies to reliably assess the functional trajectory of those
502 assemblages. Future studies will also need to integrate data on other components of
503 the arthropod communities, such as plant-dweller and sedentary species in order to

504 better understand the functioning and the role of these newly built arthropod
505 assemblages in both low and high intensity land-uses.

506

507 **ACKNOWLEDGEMENTS**

508 We are very thankful to all of the taxonomists who assisted in the identification of the
509 morphotypes: Andrew Polaszek, António Bivar Sousa, Artur Serrano, Arturo Baz,
510 Fernando Ilharco, Henrik Enghoff, Jordi Ribes, José Quartau, Jörg Wunderlich, Mário
511 Boieiro, Ole Karsholt, Richard zur Strassen, Volker Assing, Volker Manhart and
512 Virgílio Vieira.

513 **REFERENCES**

- 514 Agusti, N., Shayler, S. P., Harwood, J. D., Vaughan, I. P., Sunderland, K. D. &
515 Symondson, W. O. C. (2003). Collembola as alternative prey sustaining spiders
516 in arable ecosystems: prey detection within predators using molecular markers.
517 *Molecular Ecology*, 12, 3467-3475.
- 518 Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of
519 variance. *Austral Ecology*, 26, 32–46.
- 520 Anderson, M., Gorley, R. N. & Clarke, R. K. (2008) *Permanova+ for Primer: Guide*
521 *to Software and Statistical Methods*. PRIMER-E, Plymouth, UK.
- 522 Aslan, C. E., Zavaleta, E. S., Tershy, B., Croll, D., & Robichaux, R. H. (2014).
523 Imperfect replacement of native species by non-native species as pollinators of
524 endemic Hawaiian plants. *Conservation Biology*, 28, 478-488.
- 525 Attwood, S. J., Maron, M., House, A. P. N., & Zammit, C. (2008). Do arthropod
526 assemblages display globally consistent responses to intensified agricultural
527 land use and management? *Global Ecology and Biogeography*, 17, 585-599.
- 528 Bailey, P. T., & Mendonça, T. R. D. (1990). The distribution of the millipede
529 *Ommatoiulus moreleti* (Diplopoda, Julida: Julidae) in relation to other
530 *Ommatoiulus* species on the south-western Iberian Peninsula. *Journal of*
531 *Zoology*, 221, 99-111.
- 532 Barnard, P. C. (2011). *The Royal Entomological Society book of British insects*. John
533 Wiley & Sons, Oxford.
- 534 Bell, J. R., Wheeler, C. P., & Cullen, W. R. (2001). The implications of grassland and
535 heathland management for the conservation of spider communities: a review.
536 *Journal of Zoology*, 255, 377-387.
- 537 Birkhofer, K., Smith, H. G., Weisser, W. W., Wolters, V., & Gossner, M. M. (2015).
538 Land-use effects on the functional distinctness of arthropod communities.
539 *Ecography*, 38, 889-900.
- 540 Borges, P.A.V. & Vieira, V. (1994). The entomological bibliography from the
541 Azores. II - the taxa. *Boletim do Museu Municipal do Funchal*, 46, 5-75.
- 542 Borges, P. A. V. & Brown, V. K. (2004). Arthropod community structure in pastures
543 of an island archipelago (Azores): looking for local–regional species richness
544 patterns at fine-scales. *Bulletin of Entomological Research*, 94, 111-121.

- 545 Borges, P. A. V., Lobo, J. M., Azevedo, E. B., Gaspar, C. S., Melo, C., & Nunes, L.
546 V. (2006). Invasibility and species richness of island endemic arthropods: a
547 general model of endemic vs. exotic species. *Journal of Biogeography*, 33, 169-
548 187.
- 549 Borges, P. A. V., Ugland, K. I., Dinis, F., & Gaspar, C. (2008). Insect and spider
550 rarity in an oceanic island (Terceira, Azores): true rare and pseudo-rare species.
551 *Insect Ecology and Conservation* (ed. by S. Fattorini), pp. 47-70. Research
552 Signpost, Kerala, India.
- 553 Borges, P. A. V., Vieira, V., Amorim, I. R., Bicudo, N., Fritzén, N., Gaspar, C., ...
554 Torrão, E. (2010). List of arthropods (Arthropoda). *A list of the terrestrial and*
555 *marine biota from the Azores* (ed. by P. A. V. Borges, A. Costa, R. Cunha, R.
556 Gabriel, V. A. Gonçalves, F. Martins, I. Melo, M. Parente, P. Raposeiro, P.
557 Rodrigues, R. S. Santos, L. Silva, P. Vieira and V. Vieira), pp. 179–246.
558 Príncipe, Cascais.
- 559 Brown, K. A., Johnson, S. E., Parks, K. E., Holmes, S. M., Ivoandry, T., Abram, N.,
560 ... Wright, P. C. (2013). Use of provisioning ecosystem services drives loss of
561 functional traits across land use intensification gradients in tropical forests in
562 Madagascar. *Biological Conservation*, 161, 118-127.
- 563 Cardoso, P., Aranda, S. C., Lobo, J. M., Dinis, F., Gaspar, C., & Borges, P. A. (2009).
564 A spatial scale assessment of habitat effects on arthropod communities of an
565 oceanic island. *Acta Oecologica*, 35, 590-597.
- 566 Cardoso, P., Arnedo, M. A., Triantis, K. A., & Borges, P. A. V. (2010). Drivers of
567 diversity in Macaronesian spiders and the role of species extinctions. *Journal of*
568 *Biogeography*, 37, 1034-1046.
- 569 Cardoso, P., Rigal, F., Fattorini, S., Terzopoulou, S., & Borges, P. A. V. (2013).
570 Integrating landscape disturbance and indicator species in conservation studies.
571 *PLoS ONE*, 8, e63294.
- 572 Cardoso, P., Rigal, F., & Carvalho, J. C. (2015). BAT–Biodiversity Assessment
573 Tools, an R package for the measurement and estimation of alpha and beta
574 taxon, phylogenetic and functional diversity. *Methods in Ecology and*
575 *Evolution*, 6, 232-236.
- 576 Carvalho, J. C., & Cardoso, P. (2014). Drivers of beta diversity in Macaronesian
577 spiders in relation to dispersal ability. *Journal of Biogeography*, 41, 1859-1870.

578 Cornwell, W. K., & Ackerly, D. D. (2009). Community assembly and shifts in plant
579 trait distributions across an environmental gradient in coastal California.
580 *Ecological Monographs*, 79, 109-126.

581 Cáceres, M. D., & Legendre, P. (2009). Associations between species and groups of
582 sites: indices and statistical inference. *Ecology*, 90, 3566-3574.

583 de Bello, F. (2012). The quest for trait convergence and divergence in community
584 assembly: are null-models the magic wand? *Global Ecology and Biogeography*,
585 21, 312-317.

586 Díaz, S., Fargione, J., Chapin III, F. S., & Tilman, D. (2006). Biodiversity loss
587 threatens human well-being. *PLoS Biology*, 4, 1300-1305.

588 Didham, R. K., Tylianakis, J. M., Gemmill, N. J., Rand, T. A., & Ewers, R. M.
589 (2007). Interactive effects of habitat modification and species invasion on native
590 species decline. *Trends in Ecology & Evolution*, 22, 489-496.

591 Diekötter, T., Wamser, S., Wolters, V., & Birkhofer, K. (2010). Landscape and
592 management effects on structure and function of soil arthropod communities in
593 winter wheat. *Agriculture, Ecosystems & Environment*, 137, 108-112.

594 Drenovsky, R. E., Grewell, B. J., D'Antonio, C. M., Funk, J. L., James, J., Molinari,
595 N., ... Richards, C. L. (2012). A functional trait perspective on plant invasion.
596 *Annals of Botany*, 110, 141-153.

597 Dufrière, M., & Legendre, P. (1997). Species assemblages and indicator species: the
598 need for a flexible asymmetrical approach. *Ecological Monographs*, 67, 345-
599 366.

600 Florencio, M., Cardoso, P., Lobo, J. M., Azevedo, E. B., & Borges, P. A. V. (2013).
601 Arthropod assemblage homogenization in oceanic islands: the role of
602 indigenous and exotic species under landscape disturbance. *Diversity and*
603 *Distributions*, 19, 1450-1460.

604 Florencio, M., Lobo, J. M., Cardoso, P., Almeida-Neto, M., & Borges, P. A. V.
605 (2015). The colonisation of exotic species does not have to trigger faunal
606 homogenisation: lessons from the assembly patterns of arthropods on oceanic
607 islands. *PLoS ONE*, 10, e0128276.

608 Florencio, M., Rigal, F., Borges, P. A. V., Cardoso, P., Santos, A. M. C. & Lobo, J.
609 M. (2016). The role of plant fidelity and land-use changes on island exotic and

610 indigenou canopy spiders at local and regional scales. *Biological Invasions*, 18,
611 2309-2324.

612 Flynn, D. F., Gogol-Prokurat, M., Nogeire, T., Molinari, N., Richers, B. T., Lin, B.
613 B., ... DeClerck, F. (2009). Loss of functional diversity under land use
614 intensification across multiple taxa. *Ecology Letters*, 12, 22-33.

615 Forrest, J. R., Thorp, R. W., Kremen, C., & Williams, N. M. (2015). Contrasting
616 patterns in species and functional-trait diversity of bees in an agricultural
617 landscape. *Journal of Applied Ecology*, 52, 706-715.

618 Gaspar, C., Borges, P. A. V., & Gaston, K. J. (2008). Diversity and distribution of
619 arthropods in native forests of the Azores archipelago. *Arquipélago - Life and*
620 *Marine Sciences*, 25, 1-30.

621 Gotelli, N. J., & Graves, G. R. (1996). *Null models in ecology*, Smithsonian
622 Institution Press, Washington, DC.

623 Grez, A. A., Rand, T. A., Zaviezo, T., & Castillo-Serey, F. (2013). Land use
624 intensification differentially benefits alien over native predators in agricultural
625 landscape mosaics. *Diversity and Distributions*, 19, 749-759.

626 Grigulis, K., Sheppard, A. W., Ash, J. E., & Groves, R. H. (2001). The comparative
627 demography of the pasture weed *Echium plantagineum* between its native and
628 invaded ranges. *Journal of Applied Ecology*, 38, 281-290.

629 Hendrickx, F., Maelfait, J. P., Van Wingerden, W., Schweiger, O., Speelmans, M.,
630 Aviron, S., ... Burel, F. (2007). How landscape structure, land-use intensity and
631 habitat diversity affect components of total arthropod diversity in agricultural
632 landscapes. *Journal of Applied Ecology*, 44, 340-351.

633 Hobbs, P. R., Sayre, K., & Gupta, R. (2008). The role of conservation agriculture in
634 sustainable agriculture. *Philosophical Transactions of the Royal Society B:*
635 *Biological Sciences*, 363, 543-555.

636 Holt, R. D. (1996). *Food webs in space: an island biogeographic perspective*, In Food
637 Webs (eds G. A. Polis and K. O. Winemiller), pp. 313-323. Chapman & Hall,
638 New York.

639 Isaacs, R., Tuell, J., Fiedler, A., Gardiner, M., & Landis, D. (2008). Maximizing
640 arthropod-mediated ecosystem services in agricultural landscapes: the role of
641 native plants. *Frontiers in Ecology and the Environment*, 7(4), 196-203.

642 Kaiser-Bunbury, C. N., Traveset, A., & Hansen, D. M. (2010). Conservation and
643 restoration of plant–animal mutualisms on oceanic islands. *Perspectives in*
644 *Plant Ecology, Evolution and Systematics*, 12, 131-143.

645 Kembel, S. W. (2009). Disentangling niche and neutral influences on community
646 assembly: assessing the performance of community phylogenetic structure tests.
647 *Ecology Letters*, 12, 949-960.

648 Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly,
649 D. D., ... Webb, C. O. (2010). Picante: R tools for integrating phylogenies and
650 ecology. *Bioinformatics*, 26, 1463-1464.

651 Kenis, M., Auger-Rozenberg, M. A., Roques, A., Timms, L., Pere, C., Cock, M., ...
652 Lopez-Vaamonde, C. (2009). Ecological effects of invasive alien insects.
653 *Biological Invasions*, 11, 21–45.

654 Laliberte, E., Wells, J. A., DeClerck, F., Metcalfe, D. J., Catterall, C. P., Queiroz, C.,
655 ... McNamara, S. (2010). Land-use intensification reduces functional
656 redundancy and response diversity in plant communities. *Ecology Letters*, 13,
657 76-86.

658 Lewis, J. G. E. (2006). *The biology of centipedes*. Cambridge University Press,
659 Cambridge.

660 Losey, J. E., & Vaughan, M. (2006). The economic value of ecological services
661 provided by insects. *Bioscience*, 56, 311-323.

662 Magee, L. (1990). R² measures based on Wald and likelihood ratio joint significance
663 tests. *The American Statistician*, 44, 250-253.

664 Mayfield, M. M., & Levine, J. M. (2010). Opposing effects of competitive exclusion
665 on the phylogenetic structure of communities. *Ecology Letters*, 13, 1085-1093.

666 Mayfield, M. M., Bonser, S. P., Morgan, J. W., Aubin, I., McNamara, S., & Vesk, P.
667 A. (2010). What does species richness tell us about functional trait diversity?
668 Predictions and evidence for responses of species and functional trait diversity
669 to land-use change. *Global Ecology and Biogeography*, 19, 423-431.

670 Meijer, S. S., Whittaker, R. J., & Borges, P. A. V. (2011). The effects of land-use
671 change on arthropod richness and abundance on Santa Maria Island (Azores):
672 unmanaged plantations favour endemic beetles. *Journal of Insect Conservation*,
673 15, 505-522.

674 Mooney, H. A., & Hobbs, R. J. (2000). *Invasive species in a changing world*. Island
675 Press, Washington, DC.

676 Münkemüller, T., Gallien, L., Lavergne, S., Renaud, J., Roquet, C., Abdulhak, S., ...
677 Svenning, J. C. (2014). Scale decisions can reverse conclusions on community
678 assembly processes. *Global Ecology and Biogeography*, 23, 620-632.

679 Newbold, T., Hudson, L. N., Hill, S. L., Contu, S., Lysenko, I., Senior, R. A., ... Day,
680 J. (2015). Global effects of land use on local terrestrial biodiversity. *Nature*,
681 520, 45-50.

682 Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B.,
683 ... Oksanen, M. J. (2013). *vegan: Community ecology package*. R package
684 version 2.3-5.

685 Olden, J. D., & Rooney, T. P. (2006). On defining and quantifying biotic
686 homogenization. *Global Ecology and Biogeography*, 15, 113-120.

687 Olesen, J. M., Eskildsen, L. I. & Venkatasamy, S. (2002). Invasion of pollination
688 networks on oceanic islands: importance of invader complexes and endemic
689 super generalists. *Diversity and Distributions*, 8, 181-192.

690 Pavoine, S., Vallet, J., Dufour, A. B., Gachet, S., & Daniel, H. (2009). On the
691 challenge of treating various types of variables: application for improving the
692 measurement of functional diversity. *Oikos*, 118, 391-402.

693 Pavoine, S., Gasc, A., Bonsall, M. B., & Mason, N. W. (2013). Correlations between
694 phylogenetic and functional diversity: mathematical artefacts or true ecological
695 and evolutionary processes? *Journal of Vegetation Science*, 24, 781-793.

696 Pearce, J. L., & Venier, L. A. (2006). The use of ground beetles (Coleoptera:
697 Carabidae) and spiders (Araneae) as bioindicators of sustainable forest
698 management: a review. *Ecological Indicators*, 6, 780-793.

699 Pinheiro, J. C. & Bates, D. M. (2000). *Mixed-effects Models in S and S-Plus*.
700 Springer, New York.

701 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & the R Development Core Team
702 (2012). *nlme: Linear and Nonlinear Mixed Effects Models*. R package version
703 3.1-113.

704 R Development Core Team (2014). *R: A language and environment for statistical*
705 *computing*. R Foundation for Statistical Computing, Vienna, Austria.

706 Rader, R., Bartomeus, I., Tylianakis, J. M., & Laliberté, E. (2014). The winners and
707 losers of land use intensification: pollinator community disassembly is non-

708 random and alters functional diversity. *Diversity and Distributions*, 20, 908-
709 917.

710 Ribeiro, S. P., Borges, P. A. V., Gaspar, C., Melo, C., Serrano, A. R., Amaral, J., ...
711 Quartau, J. A. (2005). Canopy insect herbivores in the Azorean Laurisilva
712 forests: key host plant species in a highly generalist insect community.
713 *Ecography*, 28, 315-330.

714 Rigal, F., Whittaker, R. J., Triantis, K. A., & Borges, P. A. V. (2013). Integration of
715 non-indigenous species within the interspecific abundance–occupancy
716 relationship. *Acta Oecologica*, 48, 69-75.

717 Rodgers, J. C., & Parker, K. C. (2003). Distribution of alien plant species in relation
718 to human disturbance on the Georgia Sea Islands. *Diversity and Distributions*,
719 9, 385-398.

720 Ryall, K. L., & Fahrig, L. (2006). Response of predators to loss and fragmentation of
721 prey habitat: a review of theory. *Ecology*, 87, 1086-1093.

722 Saar, L., de Bello, F., Pärtel, M., & Helm, A. (2017). Trait assembly in grasslands
723 depends on habitat history and spatial scale. *Oecologia*, 184, 1-12.

724 Samu, F., Sunderland, K. D., & Szinetar, C. (1999). Scale-dependent dispersal and
725 distribution patterns of spiders in agricultural systems: a review. *Journal of*
726 *Arachnology*, 27, 325-332.

727 Sax, D. F., Gaines, S. D., & Brown, J. H. (2002). Species invasions exceed
728 extinctions on islands worldwide: a comparative study of plants and birds. *The*
729 *American Naturalist*, 160, 766-783.

730 Scherber, C., Eisenhauer, N., Weisser, W. W., Schmid, B., Voigt, W., Fischer, M., ...
731 Tschardtke, T. (2010). Bottom-up effects of plant diversity on multitrophic
732 interactions in a biodiversity experiment. *Nature*, 468, 553–556.

733 Schweiger, O., Maelfait, J. P., Wingerden, W. V., Hendrickx, F., Billeter, R.,
734 Speelmans, M., ... Bukacek, R. (2005). Quantifying the impact of
735 environmental factors on arthropod communities in agricultural landscapes
736 across organizational levels and spatial scales. *Journal of Applied Ecology*, 42,
737 1129-1139.

738 Simons, N. K., Weisser, W. W., & Gossner, M. M. (2016). Multi-taxa approach
739 shows consistent shifts in arthropod functional traits along grassland land-use
740 intensity gradient. *Ecology*, 97, 754-764.

741 Swenson, N. G., & Enquist, B. J. (2009). Opposing assembly mechanisms in a
742 Neotropical dry forest: implications for phylogenetic and functional community
743 ecology. *Ecology*, 90, 2161-2170.

744 Tecco, P. A., Díaz, S., Cabido, M., & Urcelay, C. (2010). Functional traits of alien
745 plants across contrasting climatic and land-use regimes: do aliens join the locals
746 or try harder than them? *Journal of Ecology*, 98, 17-27.

747 Terzopoulou, S., Rigal, F., Whittaker, R. J., Borges, P. A. V., & Triantis, K. A.
748 (2015). Drivers of extinction: the case of Azorean beetles. *Biology Letters*, 11,
749 20150273.

750 Triantis, K. A., Borges, P. A. V., Ladle, R. J., Hortal, J., Cardoso, P., Gaspar, C., ...
751 Whittaker, R.J. (2010). Extinction debt on oceanic islands. *Ecography*, 33, 285-
752 294.

753 Tschardtke, T., Klein, A. M., Kruess, A., Steffan-Dewenter, I., & Thies, C. (2005).
754 Landscape perspectives on agricultural intensification and biodiversity–
755 ecosystem service management. *Ecology Letters*, 8, 857-874.

756 Vieira, V. & Borges, P. A. V. (1993). The entomological bibliography of the Azores.
757 I- Thematic: General (mainly biogeography), applied entomology, ecology and
758 biospeleology. *Boletim do Museu Municipal do Funchal*, 45, 5-28

759 Vilà, M., Espinar, J. L., Hejda, M., Hulme, P.E., Jarošík, V., Maron, J. L., ... Pyšek,
760 P. (2011). Ecological impacts of invasive alien plants: a meta-analysis of their
761 effects on species, communities and ecosystems. *Ecology Letters*, 14, 702-708.

762 Violle, C., Navas, M. L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier,
763 E. (2007). Let the concept of trait be functional! *Oikos*, 116, 882-892.

764 Webb, C. O., Ackerly, D. D., McPeck, M. A., & Donoghue, M. J. (2002).
765 Phylogenies and community ecology. *Annual Review of Ecology and*
766 *Systematics*, 33, 475-505.

767 Whittaker, R. J., Rigal, F., Borges, P. A. V., Cardoso, P., Terzopoulou, S., Casanoves,
768 F., ... Triantis, K. A. (2014). Functional biogeography of oceanic islands and
769 the scaling of functional diversity in the Azores. *Proceedings of the National*
770 *Academy of Sciences of the United States of America*, 111, 13709-13714.

771 Wise, D. H. (1995). *Spiders in ecological webs*. Cambridge University Press,
772 Melbourne.

773 **SUPPORTING INFORMATION**

774 Additional Supporting Information may be found online in the sup-orting
775 information tab for this article.

776

777 **FUNDING**

778 Financial support was provided by the project ‘Consequences of land use change on
779 Azorean fauna and flora - the 2010 Target’ (Ref: Direcção Regional da Ciência e
780 Tecnologia M.2.1.2/I/003/2008). FR was supported by FCT project PTDC/BIA-
781 BIC/119255/2010.

782

783 **AUTHOR CONTRIBUTIONS**

784 F.R., P.C. and P.A.V.B. conceived the ideas. P.A.V.B., P.C. and I.R.A. collected the
785 data; F.R. analysed the data; F.R. led the writing. P.C., P.A.V.B., I.R.A., J.M.L.,
786 K.A.T., and R.J.W. provided ideas, critique, revisions and guidance at all steps of the
787 writing process.

788

789 **BIOSKETCH**

790 François Rigal is an assistant professor at Université de Pau et des Pays de l’Adour.
791 His research interests include island biogeography, macroecology and functional
792 ecology.

793 All the authors are interested in understanding the processes that shape Azorean
794 arthropod biodiversity through a range of approaches, including molecular biology,
795 taxonomy, community ecology and biogeography.

796 **Table 1.** Selection, definition and ecological relevance of the functional traits used in the study.

Traits	Data type	Attributes (Abbreviations)	Definition	Ecological relevance
Body size	Continuous	Absolute body length in mm	Defined as mean body length measured from up to 10 individuals per species. Males and females were incorporated when clear distinction was available. Measures recorded using digital photography <i>via</i> a stereoscopic microscope	Body size is related to many life-history traits such as growth rate, fecundity/clutch size, foraging ability, dispersal and life span.
Dispersal abilities	Nominal	High dispersal ability (Hdisp ¹) vs low dispersal ability (Ldisp)	Based on the presence of active wings for coleopteran and Hemipteran, ballooning for spiders and based on descriptions of flying ability for endemics and general guides for the other species. Species subsequently classified as possessing either high or low dispersal ability.	Dispersal abilities condition potential colonisation/re-colonisation.
Type of food	Multi-choice nominal	Plants (FoodPl); Animals (FoodAni); Fungi (FoodFg); Detritus (FoodDet)	Refers to the main food consumed by species during their adult stages except for Lepidoptera, where traits were assigned by reference to the larvae.	Species can co-occur in the same site but differ in their feeding strategies and resource use. Feeding guilds can also react differently to land-use changes, such as herbivores being sensitive to change in plant diversity and biomass and predators such as spiders reacting to changes of habitat architecture (Pearce & Venier, 2006, Scherber et al., 2010)
Way of getting food	Nominal	High active search (GetHact ¹) vs low active search (GetLact)	Refers to the mobility of the species in getting their food. Species classified as having active search or low active search such as species with ambush tactics or using traps	Land-use changes can impact the architecture at micro-scale and reduce potential foraging sites selected by low active search species and affect their feeding opportunities.
Mode of ingestion	Nominal	Chewing and cutting (IngCC); Piercing and sucking (IngPS); extra intestinal digestion and sucking (IngEDS)	Defined as the way nutrients are ingested.	Mode of ingestion can be related to host specificity. For instance, land-use changes may alter the kind of plants available and consequently alter the availability of resources for chewing and sucking species.
Period of activity	Multi-choice nominal	Day (ActDay); Night (ActNig); Twilight (ActTwi)	Classified as species being active during the day, during the twilight or during the night or to a combination of those periods.	Species can co-occur in the same sites but have separated temporal niches. Circadian activities also play important roles in species interaction (e.g. prey-predator). Land-use changes may promote high proportion of open micro-habitats, which are less favourable for species with day activity.

797 ¹ Attributes used in the CWM analysis to avoid collinearity




Figure 1. Rarefied species richness (A, B), mean pairwise functional distance (MFD) (C, D) and standardized effect sizes of MPD (SES_{MFD}) (E, F) across the four land-use types for both indigenous (A, C, E) and exotic arthropod elements (B, D, F) in Terceira Island. The Magee's pseudo- R^2 and the P values from the generalized least squares (GLS) models are given on the top right of each panel. Land-use types accompanied by a different letter are significantly different from each other (*post hoc* tests; $P < 0.05$, see Table S4). For the SES_{MFD} values, asterisks indicate significant deviations from 0 according to the Wilcoxon test with $**P < 0.01$. For clarity, results of the *post-hoc* tests following the GLS are not given for SES_{MFD} (but see Table S4). Land-use types are ranked according to their intensity of use/transformation, represented by either indigenous (green) or exotic (grey) gradation, with NAT = Native forest, EXO = exotic forest, SEM = Semi-natural pastures and INT = intensive managed pastures. See Table S1.1 in Appendix S1 for detailed land-use features.




Figure 2. Two-dimensional ordination solution using non-metric multidimensional scaling (NMDS) based on the Bray-Curtis distance computed by combining the “community-weighted mean (CWM) x sites” matrices of indigenous and exotic arthropod elements across the land-use gradient in Terceira island. Dots indicate elements while lines delimit the smallest polygon that encloses all elements for a given land-use. The stress value of NMDS was 0.13. The black squares and their associated labels indicate the weighted mean scores of the attributes (i.e. CWMs) in the two-dimensional ordination configuration. Land-use types are ranked according to their intensity of transformation and use, represented by either green (i.e. indigenous) or grey (exotic) gradation with NAT = Native forest, EXO = exotic forest, SEM = Semi-natural pastures and INT = intensive managed pastures. See Table S1.1 in Appendix S1 for land-use features and Table 1 for the abbreviations of the trait attributes.




Figure 3. Results of the indicator analysis showing correlations between trait attributes (CWM) and land-uses for both indigenous and exotic arthropod species in Terceira Island. Correlations were computed using point-biserial correlation coefficients (r_{pb}) between CWM and each land-use separately using a binary coding for land-uses, with 1 and 0 indicating respectively whether sites belong or not to the land-use type indicated. For each r_{pb} value, the 95% confidence interval was calculated by bootstrapping the observed data 9999. Large dots indicate values not overlapping zero (i.e. significant r_{pb}) while small dots indicate values overlapping zero (i.e. non-significant r_{pb}). Land-use types are ranked according to their intensity of transformation, represented by either green (i.e. indigenous) or grey (exotic) gradation, with NAT = Native forest, EXO = exotic forest, SEM = Semi-natural pastures and INT = intensive managed pastures. See Table S1.1 in Appendix S1 for land-use features and Table 1 for the abbreviations of the trait attributes.

SUPPORTING INFORMATION

Rigal, F., Cardoso, P., Lobo, J.M., Triantis, K.A., Whittaker, R.J., Amorim, I.R. and Borges, P.A.V. Functional traits of indigenous and exotic ground-dwelling arthropods show contrasting responses to land-use change in an oceanic island, Terceira, Azores.

Appendix S1. Sampling details across transects, taxonomical identifications and features of the four land-use types.

Nine sites (our sampling units) were randomly selected in each of the land-uses ($n = 36$). Detailed features of each land-use are given in Table S1.1 and a land-use distribution map on the Terceira Island is provided in Figure S1 based on the Cartographic information from DROTRH (2008) and Gaspar (2007). Information on spatial distribution of the sampling sites is given in Table S1.2 and in Figure S1.1

The sites in native forests were sampled during summer between 1999 (one site) and 2003 (8 sites) (Borges et al., 2005). Data were collected in the other three habitats during the summer of 2008 (see Cardoso et al., 2009). In each site, a transect of 150 m was set up with 30 pitfall traps spaced 5 m apart. The pitfall traps were plastic cups with diameters of 42 mm and depths of 78 mm; they were buried in the ground so that the rims of the cups were level with the surface. For each transect, half of the traps were filled with approximately 60 ml of an antifreeze liquid (diluted ethylene glycol), and the other half were filled with the same volume of an attractive Turquin solution made of dark beer, and for each litre of beer, 10 g of chloral hydrate, 5 ml formalin and 5ml glacial acetic acid. The ethylene and Turquin traps were alternated in each transect. Traps were left in the field for 2 weeks. The arthropods collected in the traps were preserved at 70% ethanol for identification in the laboratory at the species level for the taxa Araneae, Opiliones, Pseudoscorpiones, Diplopoda, Chilopoda and Insects (excluding Collembola, Diplura, Diptera and Hymenoptera). Taxonomic identification was performed in two steps: i) trained parataxonomists sorted samples into morphospecies (or RTUs, i.e. recognizable taxonomic units, sensu Oliver & Beattie 1996) using a non-complete reference collection; ii) experienced taxonomists assisted in the identification of the morphospecies: Andrew Polaszek, António Bivar Sousa, Artur Serrano, Arturo Baz, Fernando Ilharco, Henrik Enghoff, Jordi Ribes, José Quartau, Jörg Wunderlich, Mário Boieiro, Ole Karsholt, Richard zur Strassen, Volker Assing, Volker Manhart and Virgílio Vieira.

All species were classified as indigenous or exotic. Indigenous species comprise Azorean endemics and other native non-endemics. Exotic species are those considered to have colonized via human mediation, many of which having a cosmopolitan distribution (Borges et al., 2010). As in some of our previous studies (e.g. Borges et al., 2005, Gaspar, Borges & Gaston, 2008), we dealt with unidentified morphospecies as follows. When other species in the same genus, subfamily or family were present in the archipelago and all belonged to the same colonization category (according to Borges et al.,

2010), the unknown morphospecies were classified similarly. If no information was available, we assumed the species to be native since exotics are usually widespread and easier to identify (Borges et al., 2010).

Table S1.1. Features of land-use types





Land-use (Rank)	Type of disturbance	Disturbance intensity/frequency	Types of vegetation	Complexity of habitats for epigeal arthropods
<p>Native forest (1)</p> 	<p>Subject to a very low degree of, or no, human management. In some sites, occurrence of invasive plants with modification of soil cover (decreasing the cover of mosses and ferns)</p>	<p>Generally low, but variable depending on the spread of invasive plant species</p>	<p><i>Juniperus brevifolia</i> <i>Laurus azorica</i>, <i>Ilex perado</i> subsp <i>azorica</i>, <i>Erica azorica</i></p>	<p>The evergreen native forest of the Azores is complex in architecture and offers a great variety of substrates for epigeal arthropods. Most remnant patches are characterized by a dense tree and shrub cover of small stature.</p>
<p>Exotic forest (2)</p> 	<p>Subject to some human management and sensitivity to wind-storms</p>	<p>Low to medium depending on management type</p>	<p><i>Cryptomeria japonica</i> and <i>Eucalyptus globulus</i></p>	<p>Medium to low if the understory is completely absent.</p>
<p>Semi-natural pastures (3)</p> 	<p>Cattle grazing activity concentrated mainly in spring and particularly in summer. Fertilizers are added in Spring by hand or in some specifically large sites using machinery.</p>	<p>Medium</p>	<p><i>Lotus uliginosus</i> + exotic and native grasses, rushes, sedges and ferns.</p>	<p>Medium, depending on soil characteristics. Some sites can be badly drained, with the formation of "hummocks" as an adaptation of plants to periodic flooding, topographic conditions, cattle trampling or a combination of some of the former together with a fortuitous result of the morphology of some of the dominant species.</p>
<p>Intensive-managed pastures (4)</p> 	<p>Cattle grazing and fertilization. In the summer some sites are used for frequent hay cut or corn production. Fertilizers are added using machinery</p>	<p>High</p>	<p>Poor vascular flora of dominant species, mainly composed of introduced grasses and leguminous forbs</p>	<p>Uniform monocultures with low complexity.</p>

Table S1.2. Information on the spatial distribution of the 36 sites sampled in this study. For each site, the distance to the closest site and the distance from the edge of the sampled land-use type as well as its geographical coordinates (Longitude and Latitude) and the land-use type. Land-use types are ranked according to their intensity uses with NAT = native forest, EXO = exotic forest, SEM = semi-natural pastures and INT = intensive managed pastures (See Table S1.1 in Appendix S1 for detailed land-use features).

Site Code	Distance to the closest site (m)	Distance from the edge of the sampled land-use type (m)	Longitude	Latitude	Land-use types
T-46	783	130	38.73339	-27.28363	NAT
T-57	1818	160	38.75712	-27.3086	NAT
T-7	1243	506	38.73908	-27.29111	NAT
TE-45	2613	677	38.73864	-27.29475	NAT
TE-48	872	168	38.75121	-27.33176	NAT
TE-49	1223	1266	38.74745	-27.31885	NAT
TE-50	3432	905	38.75499	-27.31732	NAT
TY-5	2232	250	38.72592	-27.32619	NAT
TY-6	2039	190	38.75057	-27.33285	NAT
T-25	406	150	38.72638	-27.22258	EXO
T-26	340	180	38.7252	-27.22912	EXO
T-36	733	157	38.73765	-27.13682	EXO
T-37	1818	170	38.74598	-27.1343	EXO
T-62	872	152	38.73981	-27.2657	EXO
T-63	783	150	38.73186	-27.28639	EXO
T-64	3727	140	38.68872	-27.25919	EXO
T-65	765	158	38.76637	-27.2494	EXO
T-66	990	164	38.75594	-27.18915	EXO
T-61	1833	200	38.75454	-27.20551	SEM
T-67	3390	157	38.73399	-27.22935	SEM
T-69	2049	180	38.70443	-27.20951	SEM
T-72	3261	900	38.71488	-27.32124	SEM
T-73	340	935	38.72635	-27.26478	SEM
T-76	2234	150	38.73714	-27.25905	SEM
T-77	137	718	38.70727	-27.23324	SEM
T-78	765	873	38.70962	-27.25162	SEM
T-86	872	329	38.72529	-27.17557	SEM
T-70	2464	1855	38.76603	-27.14744	INT

T-74	1223	2600	38.67475	-27.3074	INT
T-75	2191	1000	38.71507	-27.10464	INT
T-80	406	1487	38.78593	-27.29137	INT
T-81	733	870	38.67444	-27.14455	INT
T-82	3085	695	38.75131	-27.10527	INT
T-83	3594	3800	38.66696	-27.10976	INT
T-87	2191	2900	38.69309	-27.13394	INT
T-90	137	934	38.69633	-27.31322	INT




Figure S1.1. Land-use distribution map of Terceira Island with the selected sampling sites. Sampling sites are marked with black dots.

References

- Borges, P.A., Aguiar, C., Amaral, J., Amorim, I.R., André, G., Arraiol, A., ... Ilharco, F. (2005). Ranking protected areas in the Azores using standardised sampling of soil epigean arthropods. *Biodiversity & Conservation*, 14, 2029-2060.
- Borges, P. A. V., Vieira, V., Amorim, I. R., Bicudo, N., Fritzen, N., Gaspar, C., ... Torrão, E. (2010). List of arthropods (Arthropoda). A list of the terrestrial and marine biota from the Azores (ed. by P. A. V. Borges, A.

- Costa, R. Cunha, R. Gabriel, V. A. Gonçalves, F. Martins, I. Melo, M. Parente, P. Raposeiro, P. Rodrigues, R. S. Santos, L. Silva, P. Vieira and V. Vieira), pp. 179–246. Príncipe, Cascais.
- DROTRH. Direcção Regional do Ordenamento do Território e Recursos Hídricos (2008). Carta de ocupação do solo da Região Autónoma dos Açores. Secretaria Regional do Ambiente. Direcção Regional do Ordenamento do Território e dos Recursos Hídricos, Ponta Delgada, Portugal.
- Cardoso, P., Aranda, S. C., Lobo, J. M., Dinis, F., Gaspar, C., & Borges, P. A. (2009) A spatial scale assessment of habitat effects on arthropod communities of an oceanic island. *Acta Oecologica*, 35, 590-597.
- Gaspar, C.S. (2007). Arthropod diversity and conservation planning in native forests of the Azores Archipelago. Doctoral Dissertation, University of Sheffield, Sheffield, UK.
- Gaspar, C., Borges, P. A. V., & Gaston, K. J. (2008). Diversity and distribution of arthropods in native forests of the Azores archipelago. *Arquipélago - Life and Marine Sciences*, 25, 1-30.
- Oliver, I. & Beattie, A.J. (1996). Invertebrate morphospecies as surrogates for species: a case study. *Conservation Biology*, 10, 99–109.

Appendix S2. Indigenous and exotic species sampled across the four land-uses considered in the study. Species taxonomy, species origin, trait attributes and mean abundance per land-use are given. Traits and their associated attributes are: Body size in (mm); Dispersal abilities (Disp) with; Type of food with Plants (FoodPl); Animals (FoodAni); Fungi (FoodFg); Detritus (FoodDet); Way of getting food (Get) with High active search (GetHact) vs low active search (GetLact); Mode of ingestion (Ing) with Chewing and cutting (IngCC); Piercing and sucking (IngPS); External digestion and sucking (IngEDS) and period of activities with Day (ActDay); Night (ActNig); Twilight (ActTwi). See Table 1 in main text for ecological relevance of the traits. Land-use types are ranked according to their intensity uses with NAT = native forest, EXO = exotic forest, SEM = semi-natural pastures and INT = intensive managed pastures. See Table S1 for detailed land-use features.

Taxonomy and species origin					Functional traits and their associated attributes											Land-uses			
Class	Order	Family	Species	Origin	Body size	Disp	FoodPl	FoodAni	FoodFg	FoodDet	Get	Ing	ActDay	ActTwi	ActNig	NAT	EXO	SEM	INT
Arachnida	Araneae	Agelenidae	<i>Lycosoides coarctata</i>	Exo	7	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0	0	0.11
Arachnida	Araneae	Dysderidae	<i>Dysdera crocata</i>	Exo	13	Ldisp	0	1	0	0	GetHact	IngEDS	0	0	1	1.78	4.67	0	0.89
Arachnida	Araneae	Gnaphosidae	<i>Zelotes aeneus</i>	Exo	6	Hdisp	0	1	0	0	GetHact	IngEDS	0	0	1	0.00	0.00	0	3.00
Arachnida	Araneae	Linyphiidae	<i>Agyneta decora</i>	Exo	2.75	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0.56	1.78	1.22	0
Arachnida	Araneae	Linyphiidae	<i>Erigone atra</i>	Exo	2.4	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0.11	7.78	1.00
Arachnida	Araneae	Linyphiidae	<i>Erigone autumnalis</i>	Exo	2.2	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0.44	8.44	39.89
Arachnida	Araneae	Linyphiidae	<i>Erigone dentipalpis</i>	Exo	2.3	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0	19.00	56.78
Arachnida	Araneae	Linyphiidae	<i>Meioneta fuscipalpa</i>	Exo	1.9	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0	0.11	18.67
Arachnida	Araneae	Linyphiidae	<i>Mermessus bryantae</i>	Exo	2.2	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0.22	0	2.44	2.89
Arachnida	Araneae	Linyphiidae	<i>Mermessus fradeorum</i>	Exo	2.2	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0.33	3.78	0.44
Arachnida	Araneae	Linyphiidae	<i>Microlinyphia johnsoni</i>	Exo	4	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0.11	0	0
Arachnida	Araneae	Linyphiidae	<i>Oedothorax fuscus</i>	Exo	2.75	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0.67	543.67	489.11
Arachnida	Araneae	Linyphiidae	<i>Ostearius melanopygius</i>	Exo	2.3	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0	0.56	7.22
Arachnida	Araneae	Linyphiidae	<i>Pelecopsis parallela</i>	Exo	1.65	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0	0	3.11
Arachnida	Araneae	Linyphiidae	<i>Prinerigone vagans</i>	Exo	2.1	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0	0.89	6.44
Arachnida	Araneae	Linyphiidae	<i>Tenuiphantes tenuis</i>	Exo	2.65	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	2.78	8.00	4.78	5.56
Arachnida	Araneae	Mimetidae	<i>Ero furcata</i>	Exo	2.88	Ldisp	0	1	0	0	GetLact	IngEDS	0	0	1	0.11	0.67	0	0.11
Arachnida	Araneae	Oecobiidae	<i>Oecobius navus</i>	Exo	2	Ldisp	0	1	0	0	GetLact	IngEDS	0	0	0	0.11	0	0	0

Arachnida	Araneae	Salticidae	<i>Gen. sp.1</i>	Exo	2.95	Hdisp	0	1	0	0	GetHact	IngEDS	1	0	0	0	0	0	1.44
Arachnida	Araneae	Tetragnathidae	<i>Pachygnatha degeeri</i>	Exo	3.38	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0	0	1.33	9.56
Arachnida	Araneae	Theridiidae	<i>Cryptachaea blattea</i>	Exo	3	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	1	0	0.33	0.11	0
Arachnida	Araneae	Theridiidae	<i>Steatoda grossa</i>	Exo	8.25	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	1	0	0.11	0	0
Arachnida	Araneae	Thomisidae	<i>Xysticus nubilus</i>	Exo	7	Hdisp	0	1	0	0	GetLact	IngEDS	1	0	0	0	0	1.00	0.11
Arachnida	Pseudoscorpiones	Chthoniidae	<i>Chthonius ischnocheles</i>	Exo	2	Ldisp	0	1	0	0	GetHact	IngCC	0	0	0	5.11	2.67	0.22	0
Arachnida	Pseudoscorpiones	Chthoniidae	<i>Chthonius tetrachelatus</i>	Exo	1.61	Ldisp	0	1	0	0	GetHact	IngCC	0	0	0	0.44	0.33	0	0
Chilopoda	Scutigermorpha	Scutigeridae	<i>Scutigera coleoptrata</i>	Exo	12.64	Ldisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0	1.67
Diplopoda	Julida	Blaniulidae	<i>Blaniulus guttullatus</i>	Exo	6.83	Ldisp	1	0	0	1	GetHact	IngCC	0	0	1	0	48.56	0	0
Diplopoda	Julida	Blaniulidae	<i>Choneiulus palmatus</i>	Exo	10.43	Ldisp	1	0	0	1	GetHact	IngCC	0	0	1	0.11	0.00	0	0
Diplopoda	Julida	Blaniulidae	<i>Proteroiulus fuscus</i>	Exo	11.28	Ldisp	1	0	0	1	GetHact	IngCC	0	0	1	0	1.44	0	0
Diplopoda	Julida	Julidae	<i>Brachyiulus pusillus</i>	Exo	7.53	Ldisp	1	0	0	1	GetHact	IngCC	0	0	1	0	1.11	0	0
Diplopoda	Julida	Julidae	<i>Cylindroiulus propinquus</i>	Exo	37.44	Ldisp	1	0	0	1	GetHact	IngCC	0	0	1	5.44	0.67	0.11	0
Diplopoda	Julida	Julidae	<i>Ommatoiulus moreletii</i>	Exo	40.04	Ldisp	1	0	0	1	GetHact	IngCC	0	0	1	19.22	19.22	3.67	0.67
Diplopoda	Polydesmida	Paradoxosomatidae	<i>Oxidus gracilis</i>	Exo	16	Ldisp	0	0	0	1	GetHact	IngCC	0	0	1	0	0.11	0.11	0
Diplopoda	Polydesmida	Polydesmidae	<i>Brachydesmus superus</i>	Exo	6.62	Ldisp	0	0	0	1	GetHact	IngCC	0	0	1	0.11	0	0	0
Diplopoda	Polydesmida	Polydesmidae	<i>Polydesmus coriaceus</i>	Exo	25.38	Ldisp	0	0	0	1	GetHact	IngCC	0	0	1	0.78	3.00	0.22	0
Insecta	Coleoptera	Anthicidae	<i>Gen. sp.2</i>	Exo	2.63	Hdisp	1	1	1	0	GetHact	IngCC	0	0	0	0	0	0	0.44
Insecta	Coleoptera	Carabidae	<i>Amara aenea</i>	Exo	7	Hdisp	1	1	0	0	GetHact	IngCC	1	0	0	0	0	0.22	6.11
Insecta	Coleoptera	Carabidae	<i>Anisodactylus binotatus</i>	Exo	11	Hdisp	1	1	0	0	GetHact	IngCC	0	0	1	0	0.11	1.22	1.56
Insecta	Coleoptera	Carabidae	<i>Paranchus albipes</i>	Exo	10	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	4.44	2.00	0	0
Insecta	Coleoptera	Carabidae	<i>Pseudophonus rufipes</i>	Exo	13	Hdisp	1	1	0	0	GetHact	IngCC	0	0	1	0.00	0	0.33	3.33
Insecta	Coleoptera	Carabidae	<i>Pterostichus vernalis</i>	Exo	7	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0.33	7.67	1.56
Insecta	Coleoptera	Chrysomelidae	<i>Chaetocnema hortensis</i>	Exo	2.01	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0.11	8.89
Insecta	Coleoptera	Chrysomelidae	<i>Epitrix cucumeris</i>	Exo	1.85	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0	0.67
Insecta	Coleoptera	Corylophidae	<i>Sericoderus lateralis</i>	Exo	0.5	Hdisp	0	0	1	0	GetHact	IngCC	1	0	0	0	0	0.11	1.11

Insecta	Coleoptera	Cryptophagidae	<i>Cryptophagus sp.1</i>	Exo	1.44	Hdisp	0	0	1	0	GetHact	IngCC	0	0	0	0	0	0.44	2.78
Insecta	Coleoptera	Cryptophagidae	<i>Cryptophagus sp.8</i>	Exo	1.64	Hdisp	0	0	1	0	GetHact	IngCC	0	0	0	0	0	0.11	0.11
Insecta	Coleoptera	Cryptophagidae	<i>Cryptophagus sp.9</i>	Exo	2.2	Hdisp	0	0	1	0	GetHact	IngCC	0	0	0	0	0.11	0.11	0.44
Insecta	Coleoptera	Curculionidae	<i>Coccotrypes carpophagus</i>	Exo	2	Hdisp	1	0	0	0	GetHact	IngCC	0	0	0	0.11	1.22	0	0
Insecta	Coleoptera	Curculionidae	<i>Gymnetron pascuorum</i>	Exo	2.44	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0	0.11
Insecta	Coleoptera	Curculionidae	<i>Otiorhynchus rugosostriatus</i>	Exo	11.62	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0.33	0	0
Insecta	Coleoptera	Curculionidae	<i>Otiorhynchus sulcatus</i>	Exo	7	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0.11	0	0
Insecta	Coleoptera	Curculionidae	<i>Pantomorus cervinus</i>	Exo	9.75	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0
Insecta	Coleoptera	Curculionidae	<i>Sitona discoideus</i>	Exo	5.79	Hdisp	1	0	0	0	GetHact	IngCC	0	1	0	0	0	0.44	4.67
Insecta	Coleoptera	Curculionidae	<i>Sitona puberulus</i>	Exo	6	Hdisp	1	0	0	0	GetHact	IngCC	0	1	0	0	0	0.11	0
Insecta	Coleoptera	Curculionidae	<i>Sitona sp.1</i>	Exo	4.27	Hdisp	1	0	0	0	GetHact	IngCC	0	1	0	0	0	1.56	17.22
Insecta	Coleoptera	Curculionidae	<i>Sitona sp.3</i>	Exo	4.74	Hdisp	1	0	0	0	GetHact	IngCC	0	1	0	0	0	3.22	0.22
Insecta	Coleoptera	Curculionidae	<i>Tychius picirostris</i>	Exo	2	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0.00	0.56
Insecta	Coleoptera	Curculionidae	<i>Xyleborinus alni</i>	Exo	3.14	Hdisp	0	0	1	0	GetHact	IngCC	0	0	0	0	0	0.56	0.22
Insecta	Coleoptera	Dryophthoridae	<i>Sphenophorus abbreviatus</i>	Exo	10.45	Ldisp	1	0	0	0	GetHact	IngCC	0	0	0	0	0.11	0.56	1.11
Insecta	Coleoptera	Elateridae	<i>Aeolus melliculus</i>	Exo	8	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0	0.78
Insecta	Coleoptera	Elateridae	<i>Heteroderes vagus</i>	Exo	6.55	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0	8.22
Insecta	Coleoptera	Hydrophilidae	<i>Cercyon haemorrhoidalis</i>	Exo	2	Hdisp	0	0	0	1	GetHact	IngCC	0	0	0	0.22	0	0.11	0
Insecta	Coleoptera	Hydrophilidae	<i>Cercyon sp.1</i>	Exo	2.22	Hdisp	0	0	0	1	GetHact	IngCC	0	0	0	0	0	0.11	0
Insecta	Coleoptera	Hydrophilidae	<i>Cercyon sp.2</i>	Exo	2.43	Hdisp	0	0	0	1	GetHact	IngCC	0	0	0	0	0	0.11	0
Insecta	Coleoptera	Hydrophilidae	<i>Sphaeridium bipustulatum</i>	Exo	4	Hdisp	0	0	0	1	GetHact	IngCC	0	0	0	0	0	0.11	0.22
Insecta	Coleoptera	Mycetophagidae	<i>Typhaea stercorea</i>	Exo	2.27	Hdisp	0	0	1	0	GetHact	IngCC	1	0	0	0	0	0	22.44
Insecta	Coleoptera	Nitidulidae	<i>Carpophilus fumatus</i>	Exo	3.65	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0.11	0.56
Insecta	Coleoptera	Nitidulidae	<i>Carpophilus hemipterus</i>	Exo	3.22	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0	0.11
Insecta	Coleoptera	Nitidulidae	<i>Carpophilus sp.1</i>	Exo	2.42	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0.11	0.22
Insecta	Coleoptera	Nitidulidae	<i>Epuraea biguttata</i>	Exo	2.98	Hdisp	1	0	1	0	GetHact	IngCC	1	0	0	0	0.78	0.11	0.89

Insecta	Coleoptera	Nitidulidae	<i>Phenolia limbata</i>	Exo	5.95	Hdisp	1	0	1	0	GetHact	IngCC	1	0	0	0	12.89	0	0
Insecta	Coleoptera	Nitidulidae	<i>Stelidota geminata</i>	Exo	2.17	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0.11	12.78	0	0.56
Insecta	Coleoptera	Ptiliidae	<i>Ptenidium pusillum</i>	Exo	1	Hdisp	0	0	1	0	GetHact	IngCC	0	0	0	0	0	0.33	0.11
Insecta	Coleoptera	Scarabaeidae	<i>Gen. sp.1</i>	Exo	4.19	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0	0.11
Insecta	Coleoptera	Scarabaeidae	<i>Onthophagus taurus</i>	Exo	7.82	Hdisp	1	0	0	0	GetHact	IngCC	0	0	0	0	0	0.33	0
Insecta	Coleoptera	Scarabaeidae	<i>Popillia japonica</i>	Exo	11	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0.11	0.11
Insecta	Coleoptera	Silvanidae	<i>Cryptamorpha desjardinsii</i>	Exo	4	Hdisp	0	0	1	0	GetHact	IngCC	0	0	1	0	0	0	0.11
Insecta	Coleoptera	Staphylinidae	<i>Aleochara bipustulata</i>	Exo	3	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0.11	0	0.44	2.11
Insecta	Coleoptera	Staphylinidae	<i>Amischa analis</i>	Exo	3	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	2.56	3.44
Insecta	Coleoptera	Staphylinidae	<i>Anotylus nitidifrons</i>	Exo	2	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	22.78	0.33	0.33
Insecta	Coleoptera	Staphylinidae	<i>Carpelimus corticinus</i>	Exo	3.02	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0.56	0	0	0
Insecta	Coleoptera	Staphylinidae	<i>Atheta atramentaria</i>	Exo	3	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	3.89	0.33	0.00	0
Insecta	Coleoptera	Staphylinidae	<i>Atheta fungi</i>	Exo	3	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0.22	0.00	0.44
Insecta	Coleoptera	Staphylinidae	<i>Atheta atramentaria</i>	Exo	3.36	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.22	0
Insecta	Coleoptera	Staphylinidae	<i>Cordalia obscura</i>	Exo	2	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.89	3.44
Insecta	Coleoptera	Staphylinidae	<i>Euplectus infirmus</i>	Exo	1.19	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0.11	0	0
Insecta	Coleoptera	Staphylinidae	<i>Gabrius nigrifulus</i>	Exo	5.61	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0
Insecta	Coleoptera	Staphylinidae	<i>Gyrophypnus fracticornis</i>	Exo	8.34	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0	0.33
Insecta	Coleoptera	Staphylinidae	<i>Oligota parva</i>	Exo	1	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.22	0.67
Insecta	Coleoptera	Staphylinidae	<i>Oxytelus sculptus</i>	Exo	5	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0	0.11
Insecta	Coleoptera	Staphylinidae	<i>Philonthus sp.</i>	Exo	8.15	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0	0.11
Insecta	Coleoptera	Staphylinidae	<i>Tachyporus chrysomelinus</i>	Exo	3	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0	0.11
Insecta	Coleoptera	Staphylinidae	<i>Xantholinus longiventris</i>	Exo	7	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.78	1.56
Insecta	Coleoptera	Tenebrionidae	<i>Blaps lethifera</i>	Exo	19.17	Ldisp	1	0	0	1	GetHact	IngCC	0	0	1	0	0	0	0.11
Insecta	Dermaptera	Anisolabididae	<i>Euborellia annulipes</i>	Exo	14.64	Ldisp	1	1	0	1	GetHact	IngCC	0	0	1	0	0	1.11	1.00
Insecta	Dermaptera	Forficulidae	<i>Forficula auricularia</i>	Exo	16	Ldisp	1	1	0	0	GetHact	IngCC	0	0	1	0	0.22	0.56	13.67

Insecta	Hemiptera	Aphididae	<i>Dysaphis plantaginea</i>	Exo	2.5	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0.11	0.33
Insecta	Hemiptera	Aphididae	<i>Rhopalosiphonimus latsiphon</i>	Exo	1.18	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	1.44	0	0	0
Insecta	Hemiptera	Aphididae	<i>Rhopalosiphum oxyacanthae</i>	Exo	1.7	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0.11	0	0.44	0
Insecta	Hemiptera	Aphididae	<i>Rhopalosiphum rufiabdominalis</i>	Exo	0.88	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0.22	0	0	0
Insecta	Hemiptera	Aphididae	<i>Uroleucon erigeronense</i>	Exo	0.92	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0.11	0	0	0
Insecta	Lepidoptera	Choreutidae	<i>Gen. sp.26</i>	Exo	12.73	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0.11	0	0
Insecta	Lepidoptera	Choreutidae	<i>Tebenna micalis</i>	Exo	10	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0	0.11
Insecta	Lepidoptera	Noctuidae	<i>Gen. sp.12</i>	Exo	22	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0.33	0
Insecta	Lepidoptera	Noctuidae	<i>Gen. sp.13</i>	Exo	21.62	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0	0.44
Insecta	Lepidoptera	Tineidae	<i>Oinophila v-flava</i>	Exo	8.3	Hdisp	1	0	1	0	GetHact	IngCC	0	0	0	0	0	0.11	0
Insecta	Lepidoptera	Tortricidae	<i>Gen. sp.1</i>	Exo	16.41	Ldisp	1	0	0	0	GetLact	IngCC	1	0	0	12.33	0.89	1.11	0.11
Insecta	Lepidoptera	Tortricidae	<i>Gen. sp.3</i>	Exo	9.41	Ldisp	1	0	0	0	GetLact	IngCC	1	0	0	1.89	0	0	0
Insecta	Lepidoptera	Tortricidae	<i>Gen. sp.6</i>	Exo	11.61	Ldisp	1	0	0	0	GetLact	IngCC	1	0	0	3.00	0	0	0
Insecta	Lepidoptera	Tortricidae	<i>Rhopobota naevana</i>	Exo	5.57	Hdisp	1	0	0	0	GetLact	IngCC	1	0	0	0.78	0	0	0
Insecta	Orthoptera	Gryllidae	<i>Eumodicogryllus bordigalensis</i>	Exo	10.21	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0.22	0.11	0.22
Insecta	Orthoptera	Gryllidae	<i>Gryllus bimaculatus</i>	Exo	32.69	Hdisp	1	1	0	1	GetHact	IngCC	1	0	1	0	0	0.11	17.89
Insecta	Psocoptera	Ectopsocidae	<i>Ectopsocus briggsi</i>	Exo	2	Hdisp	1	0	1	0	GetHact	IngCC	0	0	0	0	0.11	0	0
Insecta	Psocoptera	Lachesillidae	<i>Lachesilla greeni</i>	Exo	0.93	Hdisp	1	0	1	0	GetHact	IngCC	0	0	0	0	0.11	0	0
Insecta	Thysanoptera	Phlaeothripidae	<i>Nesothrips propinquus</i>	Exo	0.76	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0	0.11
Insecta	Thysanoptera	Thripidae	<i>Aptinothrips rufus</i>	Exo	1.04	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0.11	0
Arachnida	Araneae	Clubionidae	<i>Clubiona decora</i>	Ind	6	Ldisp	0	1	0	0	GetHact	IngEDS	0	0	1	0	0.22	0	0
Arachnida	Araneae	Linyphiidae	<i>Acorigone acoreensis</i>	Ind	1.4	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0.44	0	0.11	0
Arachnida	Araneae	Linyphiidae	<i>Lepthyphantes acoreensis</i>	Ind	3.25	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	1.44	0.33	0	0
Arachnida	Araneae	Linyphiidae	<i>Palliduphantes schmitzi</i>	Ind	1.95	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	1.78	0.89	1.11	0

Arachnida	Araneae	Linyphiidae	<i>Porrhomma borgesii</i>	Ind	2	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0.67	0.22	0	0
Arachnida	Araneae	Linyphiidae	<i>Tenuiphantes miguelensis</i>	Ind	2.65	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	35.22	13.89	1.00	0
Arachnida	Araneae	Linyphiidae	<i>Walckenaeria grandis</i>	Ind	2.3	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0.67	0	0	0
Arachnida	Araneae	Lycosidae	<i>Pardosa acorensis</i>	Ind	6	Hdisp	0	1	0	0	GetHact	IngEDS	1	0	0	7.67	7.89	51.44	12.11
Arachnida	Araneae	Pisauridae	<i>Pisaura acorensis</i>	Ind	10.5	Ldisp	0	1	0	0	GetHact	IngEDS	0	0	0	0.67	0	0	0
Arachnida	Araneae	Salticidae	<i>Neon acorensis</i>	Ind	2	Hdisp	0	1	0	0	GetHact	IngEDS	1	0	0	0.11	0	0.11	0
Arachnida	Araneae	Tetragnathidae	<i>Sancus acorensis</i>	Ind	4.05	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	0	0.33	0	0.11	0
Arachnida	Araneae	Theridiidae	<i>Lasaeola oceanica</i>	Ind	2	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	1	0.11	0	0	0
Arachnida	Araneae	Theridiidae	<i>Rugathodes acorensis</i>	Ind	1.8	Hdisp	0	1	0	0	GetLact	IngEDS	0	0	1	4.56	0.44	0	0
Arachnida	Araneae	Thomisidae	<i>Xysticus cor</i>	Ind	7	Hdisp	0	1	0	0	GetLact	IngEDS	1	0	0	0.89	0	0	0
Arachnida	Opiliones	Phalangiidae	<i>Homalenotus coriaceus</i>	Ind	3.82	Ldisp	1	1	1	1	GetHact	IngCC	0	0	1	0	17.89	7.22	0.22
Arachnida	Opiliones	Phalangiidae	<i>Leiobunum blackwalli</i>	Ind	6	Hdisp	1	1	1	1	GetHact	IngCC	0	0	1	4.67	32.22	14.78	3.44
Chilopoda	Geophilomorpha	Geophilidae	<i>Geophilus truncorum</i>	Ind	13.78	Ldisp	0	1	0	0	GetHact	IngCC	0	0	1	0.67	0.11	0	0
Chilopoda	Lithobiomorpha	Lithobiidae	<i>Lithobius pilicornis</i>	Ind	21.83	Ldisp	0	1	0	0	GetHact	IngCC	0	0	1	98.00	34.78	3.44	0.67
Chilopoda	Lithobiomorpha	Lithobiidae	<i>Lithobius sp.2</i>	Ind	8.97	Ldisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	16.11	13.11
Chilopoda	Scolopendromorpha	Cryptopidae	<i>Cryptops hortensis</i>	Ind	15.39	Ldisp	0	1	0	0	GetHact	IngCC	0	0	1	0.22	0.11	0.11	0.00
Insecta	Blattaria	Polyphagidae	<i>Zetha vestita</i>	Ind	5	Hdisp	0	0	0	1	GetHact	IngCC	1	0	0	0.67	0.56	0	0
Insecta	Coleoptera	Anthicidae	<i>Hirticollis quadriguttatus</i>	Ind	2.61	Hdisp	1	1	1	0	GetHact	IngCC	0	0	0	0	0	0	95.22
Insecta	Coleoptera	Carabidae	<i>Calosoma olivieri</i>	Ind	23	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.67	0.78
Insecta	Coleoptera	Carabidae	<i>Cedrorum azoricus</i>	Ind	10.71	Ldisp	0	1	0	0	GetHact	IngCC	0	0	1	11.56	0	0	0
Insecta	Coleoptera	Carabidae	<i>Ocys harpaloides</i>	Ind	6	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0.00	0.89	0	0
Insecta	Coleoptera	Carabidae	<i>Pterostichus aterrimus</i>	Ind	9	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0.11	0	0	0.11
Insecta	Coleoptera	Carabidae	<i>Stenolophus teutonius</i>	Ind	7	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0	0.11
Insecta	Coleoptera	Carabidae	<i>Trechus terrabravensis</i>	Ind	1.12	Ldisp	0	1	0	0	GetHact	IngCC	0	0	1	0.33	0	0	0
Insecta	Coleoptera	Chrysomelidae	<i>Chrysolina bankii</i>	Ind	11.78	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0.11	0
Insecta	Coleoptera	Chrysomelidae	<i>Psylliodes marcidus</i>	Ind	2.56	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0	0.11	1.33


Insecta	Coleoptera	Coccinellidae	<i>Scymnus interruptus</i>	Ind	2	Hdisp	0	1	0	0	GetHact	IngCC	1	0	0	0	0	0.11	
Insecta	Coleoptera	Curculionidae	<i>Drouetius borgesii</i>	Ind	8.12	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	6.33	0.11	0	0
Insecta	Coleoptera	Elateridae	<i>Alestrus dolosus</i>	Ind	5.01	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0.22	0	0	0.00
Insecta	Coleoptera	Elateridae	<i>Heteroderes azoricus</i>	Ind	7.65	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0	58.56
Insecta	Coleoptera	Laemophloeidae	<i>Placonotus sp.1</i>	Ind	2.37	Hdisp	1	0	1	0	GetHact	IngCC	0	0	0	0	0	0.11	0.11
Insecta	Coleoptera	Phalacridae	<i>Gen. sp.2</i>	Ind	1.34	Hdisp	1	0	1	0	GetHact	IngCC	1	0	0	0	0	0.22	0.33
Insecta	Coleoptera	Scraptiidae	<i>Anaspis proteus</i>	Ind	2	Hdisp	1	0	0	0	GetHact	IngCC	1	0	0	0	0.11	0	0
Insecta	Coleoptera	Staphylinidae	<i>Aloconota sulcifrons</i>	Ind	2.74	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	1.67	0	0.00	0.00
Insecta	Coleoptera	Staphylinidae	<i>Astenus lyonessius</i>	Ind	4	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0.33
Insecta	Coleoptera	Staphylinidae	<i>Carpelimus corticinus</i>	Ind	3	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0.11	0	0.11	0
Insecta	Coleoptera	Staphylinidae	<i>Habrocerus capillaricornis</i>	Ind	3.55	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0.11	0	0	0
Insecta	Coleoptera	Staphylinidae	<i>Ocyopus aethiops</i>	Ind	9	Hdisp	0	1	0	1	GetHact	IngCC	0	0	1	11.78	3.00	5.44	0
Insecta	Coleoptera	Staphylinidae	<i>Ocyopus olens</i>	Ind	17	Hdisp	0	1	0	1	GetHact	IngCC	0	0	1	0	0.67	3.44	3.33
Insecta	Coleoptera	Staphylinidae	<i>Phloeonomus punctipennis</i>	Ind	2.3	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0.11	0	0
Insecta	Coleoptera	Staphylinidae	<i>Proteinus atomarius</i>	Ind	1.52	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0.22	0.11	0	0
Insecta	Coleoptera	Staphylinidae	<i>Pseudoplectus perplexus</i>	Ind	0.92	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0
Insecta	Coleoptera	Staphylinidae	<i>Quedius curtipennis</i>	Ind	11	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.33	0
Insecta	Coleoptera	Staphylinidae	<i>Quedius simplicifrons</i>	Ind	11	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.44	0.33
Insecta	Coleoptera	Staphylinidae	<i>Rugilus orbiculatus</i>	Ind	4	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	3.22	3.67
Insecta	Coleoptera	Staphylinidae	<i>Scopaeus portai</i>	Ind	1.72	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0
Insecta	Coleoptera	Staphylinidae	<i>Coproporus pulchellus</i>	Ind	3.36	Hdisp	0	1	0	0	GetHact	IngCC	0	0	1	0	0.11	0	0
Insecta	Coleoptera	Zopheridae	<i>Tarphius azoricus</i>	Ind	2.84	Ldisp	0	0	1	0	GetHact	IngCC	0	0	0	0	0.11	0	0
Insecta	Hemiptera	Aphididae	<i>Acyrtosiphon pisum</i>	Ind	0.63	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0.22	0.11
Insecta	Hemiptera	Cicadellidae	<i>Anoscopus albifrons</i>	Ind	4	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0.22	1.22	6.00	2.44
Insecta	Hemiptera	Cicadellidae	<i>Aphrodes hamiltoni</i>	Ind	2.31	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	8.00	0.33	0.11	0
Insecta	Hemiptera	Cicadellidae	<i>Euscelidius variegatus</i>	Ind	3.56	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0.44	10.11

Insecta	Hemiptera	Cixiidae	<i>Cixius azoterceirae</i>	Ind	5	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	1.67	0	0	0
Insecta	Hemiptera	Cydnidae	<i>Geotomus punctulatus</i>	Ind	4.61	Hdisp	1	0	0	0	GetHact	IngPS	0	0	0	0	0	0	0.11
Insecta	Hemiptera	Delphacidae	<i>Megamelodes quadrimaculatus</i>	Ind	3	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	3.44	0.11	0	0
Insecta	Hemiptera	Delphacidae	<i>Muellerianella sp.1</i>	Ind	3.02	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0	0.22
Insecta	Hemiptera	Flatidae	<i>Cyphopterus adscendens</i>	Ind	5	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0.22	0	0.11	0
Insecta	Hemiptera	Lachnidae	<i>Cinara juniperi</i>	Ind	1.34	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0.44	0	0	0
Insecta	Hemiptera	Lygaeidae	<i>Nysius atlantidum</i>	Ind	3.34	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0	0.78
Insecta	Hemiptera	Lygaeidae	<i>Scolopostethus decoratus</i>	Ind	0.48	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0.11	0.11
Insecta	Hemiptera	Nabidae	<i>Nabis pseudoferus</i>	Ind	9	Hdisp	0	1	0	0	GetHact	IngPS	1	0	0	0	0	2.00	3.22
Insecta	Hemiptera	Saldidae	<i>Saldula palustris</i>	Ind	4.5	Hdisp	0	1	0	0	GetHact	IngPS	0	0	1	0	0	0.11	0
Insecta	Hemiptera	Tingidae	<i>Acalypta parvula</i>	Ind	1.86	Hdisp	1	0	0	0	GetHact	IngPS	1	0	0	0	0	0.44	0
Insecta	Lepidoptera	Crambidae	<i>Gen. sp.21</i>	Ind	8.57	Ldisp	1	0	0	0	GetHact	IngCC	1	0	0	0.11	0	0	0
Insecta	Lepidoptera	Crambidae	<i>Gen. sp.1</i>	Ind	1.54	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	2.11	0	0	0
Insecta	Lepidoptera	Crambidae	<i>Scoparia coecimaculalis</i>	Ind	25.4	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0.44	0.11	0.11	0
Insecta	Lepidoptera	Crambidae	<i>Scoparia semiampalis</i>	Ind	6	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0.11	0	0	0
Insecta	Lepidoptera	Geometridae	<i>Ascotis fortunata</i>	Ind	14.12	Ldisp	1	0	0	0	GetHact	IngCC	1	0	0	0.11	0	0	0
Insecta	Lepidoptera	Geometridae	<i>Cyclophora azorensis</i>	Ind	13	Ldisp	1	0	0	0	GetHact	IngCC	1	0	0	0.33	0	0	0
Insecta	Lepidoptera	Noctuidae	<i>Agrotis sp.1</i>	Ind	34	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0
Insecta	Lepidoptera	Noctuidae	<i>Gen. sp.1</i>	Ind	28	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0
Insecta	Lepidoptera	Noctuidae	<i>Gen. sp.14</i>	Ind	18.77	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0
Insecta	Lepidoptera	Noctuidae	<i>Gen. sp.5</i>	Ind	15.96	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0	0.11	0
Insecta	Lepidoptera	Noctuidae	<i>Mythimna unipuncta</i>	Ind	21.01	Hdisp	1	0	0	0	GetHact	IngCC	0	0	1	0.11	0.44	11.89	5.33
Insecta	Lepidoptera	Noctuidae	<i>Noctua pronuba</i>	Ind	24.96	Hdisp	1	0	0	0	GetLact	IngCC	0	0	0	0	0	0.11	0
Insecta	Lepidoptera	Noctuidae	<i>Phlogophora interrupta</i>	Ind	40	Ldisp	1	0	0	0	GetHact	IngCC	0	0	1	0	0.11	0.56	0.67
Insecta	Lepidoptera	Yponomeutidae	<i>Argyresthia atlanticella</i>	Ind	3.44	Hdisp	1	0	0	0	GetLact	IngCC	1	0	0	4.56	0.78	0.33	0

Insecta	Microcoryphia	Machilidae	<i>Trigoniophthalmus borgesii</i>	Ind	14.11	Ldisp	0	0	1	0	GetHact	IngCC	0	1	0	0.56	0	0	0
Insecta	Psocoptera	Ectopsocidae	<i>Ectopsocus strauschi</i>	Ind	1.44	Hdisp	1	0	1	0	GetHact	IngCC	0	0	0	0	0.22	0	0
Insecta	Thysanoptera	Phlaeothripidae	<i>Hoplothrips corticis</i>	Ind	2.76	Hdisp	0	0	1	0	GetHact	IngPS	1	0	0	0.11	0.22	0.22	0
Insecta	Trichoptera	Limnephilidae	<i>Limnephilus atlanticus</i>	Ind	9.14	Hdisp	0	1	0	1	GetHact	IngCC	0	0	0	1.67	0	0	0
Symphyla	Symphyla	Scutigereidae	<i>Scutigereella immaculata</i>	Ind	3.89	Ldisp	1	0	0	0	GetHact	IngCC	0	0	0	0	0.11	0	0

1 **APPENDIX S3. Sampling completeness.**

2 We calculated species accumulation curves for each sampling unit (transect; N = 36), using
3 the inventories obtained in their corresponding 30 pitfall traps. We use the number of
4 individuals caught as a measure of survey effort. Species accumulation curves were
5 calculated using 1000 randomizations. The degree of completeness was estimated by
6 calculating the slope at the end of the obtained accumulative curve (i.e. the rate of species
7 addition per individual or the tangent line using the last two points in the species accumulation
8 curves). The slopes so calculated were submitted to a non-parametric Kruskal-Wallis one-way
9 analysis of variance (KW) in order to examine if there were differences in the completeness of
10 the inventories across the four land-uses. KW was followed by Nemenyi–Damico–Wolfe–
11 Dunn *post-hoc* tests (Nemenyi, 1963) to identify statistically significant pairwise differences
12 between land-uses. Analyses were performed within the R programming environment (R
13 Development Core Team 2014) using the package *BAT* (Cardoso, Rigal & Carvalho. 2015).
14 The results obtained suggest that the completeness values vary according to the land-uses
15 (KW: $H=15.569$, $P= 0.001$, Figure S2). Multiple comparisons showed that the statistically
16 significant differences occur between the completeness value of native forests and intensive
17 pastures (*Post-hoc* test: $P= 0.003$) and between exotic forest and intensive pastures (*Post-*
18 *hoc* test: $P=0.017$); all other multiple comparisons being non-significant.
19




21 **Figure S3.1.** Box plots displaying the differences between the slope values reflecting
22 completeness in the sampling units (sites) between the four land-uses. Land-use types
23 accompanied by a different letter are significantly different from each other (*Post hoc* tests; P
24 < 0.05). Land-use types are ranked according to their intensity uses with NAT = native forest,
25 EXO = exotic forest, SEM = semi-natural pastures and INT = intensive managed pastures
26 (See Table S1 for detailed land-use features).

27

28 **References**

- 29 Cardoso, P., Rigal, F., & Carvalho, J. C. (2015). BAT–Biodiversity Assessment Tools, an R
30 package for the measurement and estimation of alpha and beta taxon, phylogenetic
31 and functional diversity. *Methods in Ecology and Evolution*, **6**, 232-236.
- 32 Nemenyi, P.B. (1963). Distribution-free multiple comparisons. PhD thesis, Princeton
33 University.
- 34 R Development Core Team (2014). *R: A language and environment for statistical computing*.
35 R Foundation for Statistical Computing, Vienna, Austria.


36



38

39 **Figure S4.1.** Box plots displaying difference in Simpson index values between the four land-
 40 use types for both indigenous (A) and exotic elements (B). The Magee 's pseudo-R² and the
 41 P values from the generalized least squares (GLS) models are given on the top right of each
 42 panel. Land-use types accompanied by a different letter are significantly different (*post hoc*
 43 tests; P < 0.05, see Table S4). Land-use types are ranked according to their intensity uses
 44 represented by either indigenous (green) or exotic (grey) gradation, with NAT = Native forest,
 45 EXO = exotic forest, SEM = Semi-natural pastures and INT = intensive managed pastures
 46 (See Table S1.1 in Appendix S1 for detailed land-use features).

47



49

50 **Figure S4.2.** A two-dimensional non-metric multidimensional scaling (NMDS) ordination
 51 based on the Bray-Curtis distance measure, applied to the “CWMs x sites” matrix for
 52 indigenous (A) and exotic (B) elements. Dots indicate site types (i.e. where assemblages
 53 were sampled) while lines delimit the smallest polygon that encloses all elements for a given
 54 land-use. The stress value of the NMDS was 0.1 and 0.07 for indigenous and exotics,
 55 respectively. The black squares and their associated labels indicate the weighted averages
 56 scores of the attributes (i.e. CWMs) in the two-dimensional ordination configuration. Land-use
 57 types are ranked according to their intensity uses represented by either green (indigenous) or
 58 grey (exotic) gradation with NAT = native forest, EXO = exotic forest, SEM = semi-natural
 59 pastures and INT = intensive managed pastures (See Table S1.1 in Appendix S1 for detailed
 60 land-use features).

61

62 **Table S4.1.** Moran's I values and their associated P values for both the raw diversity metrics
 63 and the corresponding residuals from the generalized linear square models (GLS) with land-
 64 use type as a factor for both indigenous and exotic arthropod species. Significant results are
 65 marked in bold.

	Raw metrics		Residuals	
	I	P	I	P
Indigenous species				
Rarefied species richness	0.192	<0.001	-0.064	0.472
Simpson index	0.153	<0.001	0.063	0.054
MFD	0.214	<0.001	0.013	0.334
SES _{MFD}	0.133	0.001	-0.046	0.723
Exotic species				
Rarefied species richness	0.028	0.239	-0.054	0.593
Simpson index	0.116	0.003	-0.051	0.642
MFD	-0.028	0.988	-0.059	0.519
SES _{MFD}	0.072	0.043	-0.033	0.923

66

67 **Table S4.2.** Results of the Tukey *post hoc* tests performed when the generalized least square
68 (GLS) models were significant. Results are given for each diversity metric and for both
69 indigenous and exotic arthropod species. The statistic *z* and its associated *P* values are
70 given. Land-use types are ranked according to their intensity uses with NAT = native forest,
71 EXO = exotic forest, SEM = semi-natural pastures and INT = intensive managed pastures.
72 Significant results are marked in bold (See Table S1.1 in Appendix S1 for detailed land-use
73 features).

74

	Indigenous species		Exotic species	
Rarefied species richness	<i>z</i>	<i>P</i>	<i>z</i>	<i>P</i>
NAT - EXO	5.809	<0.001	-0.668	0.909
NAT - SEM	-7.386	<0.001	-0.100	1.000
NAT - INT	7.663	<0.001	-4.230	<0.001
EXO - SEM	-1.577	0.392	-0.768	0.869
EXO - INT	-1.707	0.248	3.562	0.002
SEM - INT	0.277	0.993	-4.330	<0.001
Simpson index				
NAT - EXO	5.397	<0.001	-0.438	0.972
NAT - SEM	-2.403	0.077	2.815	0.025
NAT - INT	2.598	0.046	-6.403	<0.001
EXO - SEM	2.994	0.014	2.377	0.082
EXO - INT	2.799	0.027	5.965	<0.001
SEM - INT	0.195	0.997	-3.588	0.002
MFD				
NAT - EXO	4.697	<0.001	0.082	1.000
NAT - SEM	-3.501	0.003	-5.233	<0.001
NAT - INT	5.091	<0.001	2.036	0.175
EXO - SEM	1.196	0.629	-5.151	<0.001
EXO - INT	-0.394	0.979	-1.954	0.206
SEM - INT	1.590	0.385	-3.197	0.008
SES_{MFD}				
NAT - EXO	2.435	0.071	0.170	0.998
NAT - SEM	-3.334	0.005	-6.712	<0.001
NAT - INT	5.765	<0.001	6.262	<0.001

EXO - SEM	-0.899	0.805	-6.542	<0.001
EXO - INT	-3.330	0.005	-6.092	<0.001
SEM - INT	2.431	0.071	-0.450	0.970

75

76 **Table S4.3.** Results of the post hoc *t* tests performed for significant PERMANOVAs. The post
77 hoc *t* tests were performed to highlight which land-uses differ between each other in their
78 trait's attributes compositions. Analysis were performed for both indigenous and exotic
79 arthropod elements separately. The post hoc *t* tests were also performed within each land-
80 use type to highlight differences in trait's attributes compositions between indigenous and
81 exotic arthropod elements. The statistic *t* and its associated *P* values are given. Land-use
82 types are ranked according to their intensity uses with NAT = native forest, EXO = exotic
83 forest, SEM = semi-natural pastures and INT = intensive managed pastures. Significant
84 results are marked in bold (See Table S1.1 in Appendix S1 for detailed land-use features).

85

Pairwise comparisons between land-uses for indigenous		
	<i>t</i>	<i>P</i>
NAT - EXO	3.382	<0.001
NAT - SEM	2.697	<0.001
NAT - INT	4.044	<0.001
EXO - SEM	2.210	0.012
EXO - INT	3.457	<0.001
SEM - INT	1.567	0.056
Pairwise comparisons between land-uses for exotics		
	<i>t</i>	<i>P</i>
NAT - EXO	2.423	0.004
NAT - SEM	9.595	<0.001
NAT - INT	9.480	<0.001
EXO - SEM	5.247	<0.001
EXO - INT	4.477	<0.001
SEM - INT	2.702	0.004
Comparisons between indigenous and exotics within land-uses		
	<i>t</i>	<i>P</i>
NAT	15.396	<0.001
EXO	5.4323	<0.001
SEM	16.301	<0.001
INT	22.494	<0.001

86

87

88