



HAL
open science

Families of rational solutions of order 5 to the KPI equation depending on 8 parameters

Pierre Gaillard

► **To cite this version:**

Pierre Gaillard. Families of rational solutions of order 5 to the KPI equation depending on 8 parameters. *New Horizons in Mathematical Physics*, 2017, 1 (1), pp.26-32. 10.22606/nhmp.2017.11004 . hal-01598396

HAL Id: hal-01598396

<https://hal.science/hal-01598396>

Submitted on 29 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rational solutions of order 5 to the KPI equation depending on 8 parameters.

P. Gaillard,

Université de Bourgogne,
Institut de mathématiques de Bourgogne,
9 avenue Alain Savary BP 47870
21078 Dijon Cedex, France :
E-mail : Pierre.Gaillard@u-bourgogne.fr

Abstract

In this paper, we go on with the study of rational solutions to the Kadomtsev-Petviashvili equation (KPI). We construct here rational solutions of order 5 as a quotient of 2 polynomials of degree 60 in x , y and t depending on 8 parameters. The maximum modulus of these solutions at order 5 is checked as equal to $2(2N + 1)^2 = 242$. We study their modulus patterns in the plane (x, y) and their evolution according to time and parameters $a_1, a_2, a_3, a_4, b_1, b_2, b_3, b_4$. We get triangle and ring structures as obtained in the case of the NLS equation.

PACS numbers :
33Q55, 37K10, 47.10A-, 47.35.Fg, 47.54.Bd

1 Introduction

We consider the Kadomtsev-Petviashvili equation (KPI) in the following form

$$(4u_t - 6uu_x + u_{xxx})_x - 3u_{yy} = 0, \quad (1)$$

where subscripts x , y and t denote partial derivatives.

Kadomtsev and Petviashvili [1] first proposed this equation in 1970. This equation is a model for example, for surface and internal water waves [2], and in nonlinear optics [3]. Zakharov extended the inverse scattering transform (IST) to this KPI equation, and obtained several exact solutions.

The first rational solutions were found in 1977 by Manakov, Zakharov, Bordag and Matveev [4]. Other researches were led and more general rational solutions to the KPI equation were obtained. We can mention the following works by Krichever in 1978 [5, 6], Satsuma and Ablowitz in 1979 [7], Matveev in 1979 [8], Freeman and Nimmo in 1983 [9, 10], Pelinovsky and Stepanyants in 1993 [11],

Pelinovsky in 1994 [12], Ablowitz and Villarroel [13, 14] in 1997-1999, Biondini and Kodama [15, 16, 17] in 2003-2007.

We recall the author's results about the representations of the solutions to the KPI equation, first in terms of Fredholm determinants of order $2N$ depending on $2N - 1$ parameters, then in terms of wronskians of order $2N$ with $2N - 1$ parameters. These representations allow to obtain an infinite hierarchy of solutions to the KPI equation, depending on $2N - 1$ real parameters .

Then we construct the rational solutions of order N depending on $2N - 2$ parameters without presence of a limit which can be written as a ratio of two polynomials of x , y and t of degree $2N(N + 1)$.

The maximum modulus of these solutions at order N is equal to $2(2N + 1)^2$. This method gives an infinite hierarchy of rational solutions of order N depending on $2N - 2$ real parameters. We construct here the explicit rational solutions of order 5, depending on 8 real parameters, and the representations of their modulus in the plane of the coordinates (x, y) according to the real parameters $a_1, b_1, a_2, b_2, a_3, b_3, a_4, b_4$ and time t .

2 Rational solutions to the KPI equation of order N depending on $2N - 2$ parameters

2.1 Fredholm representation

One defines real numbers λ_j such that $-1 < \lambda_\nu < 1$, $\nu = 1, \dots, 2N$ depending on a parameter ϵ that will be intended to tend towards 0; they can be written as

$$\lambda_j = 1 - 2\epsilon^2 j^2, \quad \lambda_{N+j} = -\lambda_j, \quad 1 \leq j \leq N, \quad (2)$$

The terms $\kappa_\nu, \delta_\nu, \gamma_\nu, \tau_\nu$ and $x_{r,\nu}$ are functions of λ_ν , $1 \leq \nu \leq 2N$; they are defined by the formulas :

$$\begin{aligned} \kappa_j &= 2\sqrt{1 - \lambda_j^2}, & \delta_j &= \kappa_j \lambda_j, & \gamma_j &= \sqrt{\frac{1 - \lambda_j}{1 + \lambda_j}}; \\ x_{r,j} &= (r - 1) \ln \frac{\gamma_j - i}{\gamma_j + i}, & r &= 1, 3, & \tau_j &= -12i\lambda_j^2 \sqrt{1 - \lambda_j^2} - 4i(1 - \lambda_j^2) \sqrt{1 - \lambda_j^2}, \\ \kappa_{N+j} &= \kappa_j, & \delta_{N+j} &= -\delta_j, & \gamma_{N+j} &= \gamma_j^{-1}, \\ x_{r,N+j} &= -x_{r,j}, & \tau_{N+j} &= \tau_j & j &= 1, \dots, N. \end{aligned} \quad (3)$$

e_ν $1 \leq \nu \leq 2N$ are defined in the following way :

$$\begin{aligned} e_j &= 2i \left(\sum_{k=1}^{1/2 M-1} a_k (je)^{2k-1} - i \sum_{k=1}^{1/2 M-1} b_k (je)^{2k-1} \right), \\ e_{N+j} &= 2i \left(\sum_{k=1}^{1/2 M-1} a_k (je)^{2k-1} + i \sum_{k=1}^{1/2 M-1} b_k (je)^{2k-1} \right), \quad 1 \leq j \leq N, \\ a_k, b_k &\in \mathbf{R}, \quad 1 \leq k \leq N - 1. \end{aligned} \quad (4)$$

ϵ_ν , $1 \leq \nu \leq 2N$ are real numbers defined by :

$$e_j = 1, \quad e_{N+j} = 0 \quad 1 \leq j \leq N. \quad (5)$$

Let I be the unit matrix and $D_r = (d_{jk})_{1 \leq j, k \leq 2N}$ the matrix defined by :

$$d_{\nu\mu} = (-1)^{\epsilon_\nu} \prod_{\eta \neq \mu} \left(\frac{\gamma_\eta + \gamma_\nu}{\gamma_\eta - \gamma_\mu} \right) \exp(i\kappa_\nu x - 2\delta_\nu y + \tau_\nu t + x_{r,\nu} + e_\nu). \quad (6)$$

Then we recall the following result :

Theorem 2.1 *The function v defined by*

$$v(x, y, t) = -2 \frac{|n(x, y, t)|^2}{d(x, y, t)^2} \quad (7)$$

where

$$n(x, y, t) = \det(I + D_3(x, y, t)), \quad (8)$$

$$d(x, y, t) = \det(I + D_1(x, y, t)), \quad (9)$$

and $D_r = (d_{jk})_{1 \leq j, k \leq 2N}$ the matrix

$$d_{\nu\mu} = (-1)^{\epsilon_\nu} \prod_{\eta \neq \mu} \left(\frac{\gamma_\eta + \gamma_\nu}{\gamma_\eta - \gamma_\mu} \right) \exp(i\kappa_\nu x - 2\delta_\nu y + \tau_\nu t + x_{r,\nu} + e_\nu). \quad (10)$$

is a solution to the KPI equation (1), depending on $2N - 1$ parameters a_k, b_k , $1 \leq k \leq N - 1$ and ϵ .

2.2 Wronskian representation

We use the following notations :

$$\phi_{r,\nu} = \sin \Theta_{r,\nu}, \quad 1 \leq \nu \leq N, \quad \phi_{r,\nu} = \cos \Theta_{r,\nu}, \quad N + 1 \leq \nu \leq 2N, \quad r = 1, 3, \quad (11)$$

with the arguments

$$\Theta_{r,\nu} = \frac{\kappa_\nu x}{2} + i\delta_\nu y - i\frac{x_{r,\nu}}{2} - i\frac{\tau_\nu}{2} t + \gamma_\nu w - i\frac{e_\nu}{2}, \quad 1 \leq \nu \leq 2N. \quad (12)$$

$W_r(w)$ denotes the wronskian of the functions $\phi_{r,1}, \dots, \phi_{r,2N}$ defined by

$$W_r(w) = \det[(\partial_w^{\mu-1} \phi_{r,\nu})_{\nu, \mu \in [1, \dots, 2N]}]. \quad (13)$$

We consider the matrix $D_r = (d_{\nu\mu})_{\nu, \mu \in [1, \dots, 2N]}$ defined in (10). Then we have the following statement :

Theorem 2.2 *The function v defined by :*

$$v(x, y, t) = -2 \frac{|W_3(\phi_{3,1}, \dots, \phi_{3,2N})(0)|^2}{(W_1(\phi_{1,1}, \dots, \phi_{1,2N})(0))^2}$$

is a solution to the KPI equation depending on $2N - 1$ real parameters a_k, b_k $1 \leq k \leq N - 1$ and ϵ , with ϕ_ν^r defined in (11)

$$\begin{aligned} \phi_{r,\nu}(w) &= \sin\left(\frac{\kappa_\nu x}{2} + i\delta_\nu y - i\frac{x_{r,\nu}}{2} - i\frac{\tau_\nu}{2}t + \gamma_\nu w - i\frac{e_\nu}{2}\right), & 1 \leq \nu \leq N, \\ \phi_{r,\nu}(w) &= \cos\left(\frac{\kappa_\nu x}{2} + i\delta_\nu y - i\frac{x_{r,\nu}}{2} - i\frac{\tau_\nu}{2}t + \gamma_\nu w - i\frac{e_\nu}{2}\right), & N + 1 \leq \nu \leq 2N, \quad r = 1, 3, \end{aligned}$$

$\kappa_\nu, \delta_\nu, x_{r,\nu}, \gamma_\nu, e_\nu$ being defined in(3), (2) and (4).

2.3 Rational solutions

We recall the last result concerning the rational solutions to the KPI equation as a quotient of two determinants.

We use the following notations :

$$\begin{aligned} X_\nu &= \frac{\kappa_\nu x}{2} + i\delta_\nu y - i\frac{x_{3,\nu}}{2} - i\frac{\tau_\nu}{2}t - i\frac{e_\nu}{2}, \\ Y_\nu &= \frac{\kappa_\nu x}{2} + i\delta_\nu y - i\frac{x_{1,\nu}}{2} - i\frac{\tau_\nu}{2}t - i\frac{e_\nu}{2}, \end{aligned}$$

for $1 \leq \nu \leq 2N$, with $\kappa_\nu, \delta_\nu, x_{r,\nu}$ defined in (3) and parameters e_ν defined by (4).

We define the following functions :

$$\begin{aligned} \varphi_{4j+1,k} &= \gamma_k^{4j-1} \sin X_k, & \varphi_{4j+2,k} &= \gamma_k^{4j} \cos X_k, \\ \varphi_{4j+3,k} &= -\gamma_k^{4j+1} \sin X_k, & \varphi_{4j+4,k} &= -\gamma_k^{4j+2} \cos X_k, \end{aligned} \quad (14)$$

for $1 \leq k \leq N$, and

$$\begin{aligned} \varphi_{4j+1,N+k} &= \gamma_k^{2N-4j-2} \cos X_{N+k}, & \varphi_{4j+2,N+k} &= -\gamma_k^{2N-4j-3} \sin X_{N+k}, \\ \varphi_{4j+3,N+k} &= -\gamma_k^{2N-4j-4} \cos X_{N+k}, & \varphi_{4j+4,N+k} &= \gamma_k^{2N-4j-5} \sin X_{N+k}, \end{aligned} \quad (15)$$

for $1 \leq k \leq N$.

We define the functions $\psi_{j,k}$ for $1 \leq j \leq 2N, 1 \leq k \leq 2N$ in the same way, the term X_k is only replaced by Y_k .

$$\begin{aligned} \psi_{4j+1,k} &= \gamma_k^{4j-1} \sin Y_k, & \psi_{4j+2,k} &= \gamma_k^{4j} \cos Y_k, \\ \psi_{4j+3,k} &= -\gamma_k^{4j+1} \sin Y_k, & \psi_{4j+4,k} &= -\gamma_k^{4j+2} \cos Y_k, \end{aligned} \quad (16)$$

for $1 \leq k \leq N$, and

$$\begin{aligned} \psi_{4j+1,N+k} &= \gamma_k^{2N-4j-2} \cos Y_{N+k}, & \psi_{4j+2,N+k} &= -\gamma_k^{2N-4j-3} \sin Y_{N+k}, \\ \psi_{4j+3,N+k} &= -\gamma_k^{2N-4j-4} \cos Y_{N+k}, & \psi_{4j+4,N+k} &= \gamma_k^{2N-4j-5} \sin Y_{N+k}, \end{aligned} \quad (17)$$

for $1 \leq k \leq N$.

Then we get the following result :

Theorem 2.3 *The function v defined by :*

$$v(x, y, t) = -2 \frac{|\det((n_{jk})_{j,k \in [1, 2N]})|^2}{\det((d_{jk})_{j,k \in [1, 2N]})^2} \quad (18)$$

is a rational solution to the KPI equation (1).

$$(4u_t - 6uu_x + u_{xxx})_x - 3u_{yy} = 0,$$

where

$$\begin{aligned} n_{j1} &= \varphi_{j,1}(x, y, t, 0), \quad 1 \leq j \leq 2N & n_{jk} &= \frac{\partial^{2k-2} \varphi_{j,1}}{\partial \epsilon^{2k-2}}(x, y, t, 0), \\ n_{jN+1} &= \varphi_{j,N+1}(x, y, t, 0), \quad 1 \leq j \leq 2N & n_{jN+k} &= \frac{\partial^{2k-2} \varphi_{j,N+1}}{\partial \epsilon^{2k-2}}(x, y, t, 0), \\ d_{j1} &= \psi_{j,1}(x, y, t, 0), \quad 1 \leq j \leq 2N & d_{jk} &= \frac{\partial^{2k-2} \psi_{j,1}}{\partial \epsilon^{2k-2}}(x, y, t, 0), \\ d_{jN+1} &= \psi_{j,N+1}(x, y, t, 0), \quad 1 \leq j \leq 2N & d_{jN+k} &= \frac{\partial^{2k-2} \psi_{j,N+1}}{\partial \epsilon^{2k-2}}(x, y, t, 0), \\ & & & 2 \leq k \leq N, \quad 1 \leq j \leq 2N \end{aligned} \quad (19)$$

The functions φ and ψ are defined in (14), (15), (16), (17).

3 Explicit expression of rational solutions of order 5 depending on 8 parameters

In the the following, we explicitly construct rational solutions to the KPI equation of order 5 depending on 8 parameters.

Because of the length of the expression, we cannot give it in this text. We only give the expression without parameters in the appendix.

We give patterns of the modulus of the solutions in the plane (x, y) of coordinates in function of parameters $a_1, a_2, a_3, a_4, b_1, b_2, b_3, b_4$ and time t .

When at least one parameter is not equal to 0, we observe the presence of 15 peaks. The maximum modulus of these solutions is checked in this case $N = 5$, equal to $2(2N + 1)^2 = 2 \times 11^2 = 242$.




Figure 1. Solution of order 5 to KPI, on the left for $t = 0$; in the center for $t = 0,01$; on the right for $t = 0,1$; all parameters equal to 0.




Figure 2. Solution of order 5 to KPI, on the left for $t = 0,2$; in the center for $t = 20$; on the right for $t = 50$; all parameters equal to 0.




Figure 3. Solution of order 5 to KPI for $t = 0$, on the left for $a_1 = 10^4$; in the center for $b_1 = 10^4$; on the right for $a_2 = 10^6$; all other parameters equal to 0.




Figure 4. Solution of order 5 to KPI for $t = 0$, on the left for $b_2 = 10^6$; in the center for $a_3 = 10^8$; on the right for $b_3 = 10^8$; all other parameters equal to 0.




Figure 5. Solution of order 5 to KPI for $t = 0$, on the left for $a_4 = 10^8$; in the center for $b_4 = 10^8$; on the right for $b_4 = 10^8$, sight on top; all other parameters equal to 0.

4 Conclusion

We obtain N -th order rational solutions to the KPI equation depending on $2N - 2$ real parameters. These solutions can be expressed in terms of a ratio of two polynomials of degree $2N(N + 1)$ in x, y and t . The maximum modulus of these solutions is equal to $2(2N + 1)^2$. This gives a new approach to find explicit solutions for higher orders and try to describe the structure of these rational solutions. Here, we have given a complete description of rational solutions of order 5 with 8 parameters by giving explicit expressions of polynomials of those solutions.

We construct the modulus of solutions in the (x, y) plane of coordinates; different structures appear. For a given t , when one parameter grows and the other ones are equal to 0 we obtain triangular or rings or concentric rings. There are four types of patterns. For $a_1 \neq 0$ or $b_1 \neq 0$, and other parameters equal to zero, we obtain a triangle with 15 peaks. For $a_2 \neq 0$ or $b_2 \neq 0$, and other parameters equal to zero, we obtain three concentric rings of 5 peaks on each of them. For $a_3 \neq 0$ or $b_3 \neq 0$, and other parameters equal to zero, we obtain two concentric rings of 7 peaks on each of them with a central peak; in the last case, when $a_4 \neq 0$ or $b_4 \neq 0$, and other parameters equal to zero, we obtain one ring with 9 peaks with the lump L_3 in the center.

References

- [1] B.B. Kadomtsev, V.I. Petviashvili, On the stability of solitary waves in weakly dispersing media, *Sov. Phys. Dokl.*, V. 15, N. 6, 539-541, 1970
- [2] M.J. Ablowitz, H. Segur On the evolution of packets of water waves, *J. Fluid Mech.*, V. 92, 691-715, 1979
- [3] D.E. Pelinovsky, Y.A. Stepanyants, Y.A. Kivshar, Self-focusing of plane dark solitons in nonlinear defocusing media, *Phys. Rev. E*, V. 51, 5016-5026, 1995
- [4] S.V. Manakov, V.E. Zakharov, L.A. Bordag, V.B. Matveev, Two-dimensional solitons of the Kadomtsev-Petviashvili equation and their interaction, *Phys. Letters*, V. 63A, N. 3, 205-206, 1977
- [5] I. Krichever, Rational solutions of the Kadomtsev-Petviashvili equation and integrable systems of n particles on a line, *Funct. Anal. and Appl.*, V. 12, N. 1 : 59-61, 1978
- [6] I. Krichever, S.P. Novikov Holomorphic bundles over Riemann surfaces and the KPI equation, *Funkt. Ana. E Pril.*, V. 12, 41-52, 1979
- [7] J. Satsuma, M.J. Ablowitz, Two-dimensional lumps in nonlinear dispersive systems, *J. Math. Phys.*, V. 20, 1496-1503, 1979
- [8] V.B. Matveev, Darboux transformation and explicit solutions of the Kadomtsev-Petviashvili equation depending on functional parameters, *Letters in Mathematical Physics*, V. 3, 213-216, 1979
- [9] N. C Freeman, J.J.C. Nimmo Rational solutions of the KdV equation in wronskian form, *Phys. Letters*, V. 96 A, N. 9, 443-446, 1983
- [10] N. C Freeman, J.J.C. Nimmo The use of Bäcklund transformations in obtaining N -soliton solutions in wronskian form, *J. Phys. A : Math. Gen.*, V. 17 A, 1415-1424, 1984
- [11] D.E. Pelinovsky, Y.A. Stepanyants, New multisolitons of the Kadomtsev-Petviashvili equation, *Phys. JETP Lett.*, V. 57, 24-28, 1993
- [12] D.E. Pelinovsky, Rational solutions of the Kadomtsev-Petviashvili hierarchy and the dynamics of their poles. I. New form of a general rational solution, *J.Math.Phys.*, V. 35, 5820-5830, 1994
- [13] M.J Ablowitz, J. Villarroel, Solutions to the time dependent Schrödinger and the Kadomtsev-Petviashvili equations, *Phys. Rev. Lett.*, V. 78, 570, 1997
- [14] J. Villarroel, M.J Ablowitz, On the discrete spectrum of the nonstationary Schrödinger equation and multipole lumps of the Kadomtsev-Petviashvili I equation, *Commun. Math. Phys.*, V. 207, 1-42, 1999

- [15] G. Biondini, Y. Kodama, On a family of solutions of the Kadomtsev-Petviashvili equation which also satisfy the Toda lattice hierarchy, *J. Phys. A: Math. Gen.*, V. 36, 1051910536, 2003
- [16] Y. Kodama, Young diagrams and N solitons solutions to the KP equation, *J. Phys. A: Math. Gen.*, V. 37, 1116911190, 2004
- [17] G. Biondini, Line Soliton Interactions of the Kadomtsev-Petviashvili Equation, *PRL*, V. 99, 064103-1-4, 2007
- [18] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, *Max-Planck-Institut für Mathematik, MPI 02-31*, V. **161**, 2002
- [19] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, *Lett. Math. Phys.*, V. **68**, 77-90, 2004
- [20] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. **89**, 1-12, 2009
- [21] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *RIMS Kyoto*, N. 1653, 1-19, 2009
- [22] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. **42**, 1-16, 2009
- [23] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Spe. Top.*, V. **185**, 247-258, 2010
- [24] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A : Meth. Theor.*, V. **44**, 1-15, 2011
- [25] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *J. Math. Sciences : Adv. Appl.*, V. **13**, N. 2, 71-153, 2012
- [26] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, V. **54**, 013504-1-32, 2013
- [27] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. **4**, N. 4, 246-266, 2013
- [28] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. **2013**, ID 645752, 1-10, 2013
- [29] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. **2013**, 1-111, 2013

- [30] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. **7**, N. **45**, 1-6, 2013
- [31] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. **54**, 073519-1-22, 2013
- [32] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. **88**, 042903-1-9, 2013
- [33] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. **89**, 015004-1-7, 2014
- [34] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. **61**, 365-369, 2014
- [35] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. **482**, 012016-1-7, 2014
- [36] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. **26**, N. 2, 1550016-1-14, 2014
- [37] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. **5**, 093506-1-12, 2014
- [38] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. **574**, 012031-1-5, 2015
- [39] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. **355**, 293-298, 2015
- [40] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation *Phys. Lett. A*, V. **379**, 1309-1313, 2015
- [41] P. Gaillard, Other $2N-2$ parameters solutions to the NLS equation and $2N+1$ highest amplitude of the modulus of the N -th order AP breather, *J. Phys. A: Math. Theor.*, V. **48**, 145203-1-23, 2015
- [42] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. **4**, 346-364, 2015
- [43] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, *Jour. Phys. : Conf. Ser.*, V. **633**, 012106-1-6, 2016

- [44] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, *Journal of Theoretical and Applied Physics*, V. **10**,1-7, 2016
- [45] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, *Commun. Theor. Phys*, V. **65**, 136-144, 2016
- [46] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, *Annals Of Physics*, V. **367**, 1-5, 2016
- [47] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, *Adv. Res.*, V. **10**, 83-89, 2016
- [48] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, *Theor. And Math. Phys.*, V. **189**, 1440-1449, 2016
- [49] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, *Jour. of Math. Phys.*, V. **57**, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [50] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. 3, 130-139, 2017
- [51] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on $2N^2$ parameters, *Int. Jour. of Appl. Sci. And Math.*, V. **4**, N. 3, 60-70, 2017

Appendix : Because of the length of the complete expression, we only give in this appendix the explicit expression of the rational solution of order 5 to KPI equation without parameters. They can be written as

$$v(x, y, t) = -2 \frac{|n(x, y, t)|^2}{(d(x, y, t))^2}$$

with

$$\frac{n(x, y, t)}{d(x, y, t)} = \frac{F(2x, 4y, 4t) - iH(2x, 4y, 4t)}{Q(2x, 4y, 4t)}$$

with

$$\begin{aligned} F(X, Y, T) &= \sum_{k=0}^{30} f_k(Y, T) X^k, \\ H(X, Y, T) &= \sum_{k=0}^{30} h_k(Y, T) X^k, \\ Q(X, Y, T) &= \sum_{k=0}^{30} q_k(Y, T) X^k. \end{aligned}$$

$$\begin{aligned} f_{30} &= 1, \quad f_{29} = -90T, \quad f_{28} = 3915T^2 + 15Y^2 - 45, \quad f_{27} = -109620T^3 + (-1260Y^2 + 5460)T, \quad f_{26} = \\ &2219805T^4 + 105Y^4 + (51030Y^2 - 289170)T^2 - 3150Y^2 - 1575, \quad f_{25} = -34628958T^5 + (-1326780Y^2 + 9287460)T^3 + \\ &(-8190Y^4 + 260820Y^2 + 77490)T, \quad f_{24} = 432861975T^6 + 455Y^6 + (24877125Y^2 - 207309375)T^4 - 35175Y^4 + \\ &(307125Y^4 - 10347750Y^2 - 297675)T^2 + 67725Y^2 - 80325, \quad f_{23} = -4452294600T^7 + (-358230600Y^2 + 3462895800)T^5 + \\ &(-7371000Y^4 + 261954000Y^2 - 77225400)T^3 + (-32760Y^6 + 2583000Y^4 - 6388200Y^2 + 5027400)T, \quad f_{22} = \\ &38401040925T^8 + 1365Y^8 + (4119651900Y^2 - 45316170900)T^6 - 192780Y^6 + (127149750Y^4 - 4753444500Y^2 + \\ &3163094500)T^4 + 1943550Y^4 + (1130220Y^6 - 90852300Y^4 + 279814500Y^2 - 169136100)T^2 + 1379700Y^2 - 3472875, \quad f_{21} = \\ &-281607633450T^9 + (-38842432200Y^2 + 479056663800)T^7 + (-1678376700Y^4 + 65844009000Y^2 - 70879139100)T^5 + \\ &(-24864840Y^6 + 2037004200Y^4 - 7622861400Y^2 + 4308406200)T^3 + (-90090Y^8 + 12760440Y^6 - 133364700Y^4 - \\ &72160200Y^2 + 211519350)T, \quad f_{20} = 1774128090735T^{10} + 3003Y^{10} + (305884153575Y^2 - 418041676525)T^8 - \\ &654885Y^8 + (17622955350Y^4 - 723896781300Y^2 + 1102112515350)T^6 + 14442750Y^6 + (391621230Y^6 - 32685310350Y^4 + \\ &145679168250Y^2 - 91054615050)T^4 - 6284250Y^4 + (2837835Y^8 - 403118100Y^6 + 4390970850Y^4 + 788640300Y^2 - \\ &6550139925)T^2 + 55466775Y^2 - 59634225, \quad f_{19} = -9677062313100T^{11} + (-2039227690500Y^2 + 30588415357500)T^9 + \\ &(-151053903000Y^4 + 6483698298000Y^2 - 12960424877400)T^7 + (-4699454760Y^6 + 399453654600Y^4 - 2085756334200Y^2 + \\ &1576774949400)T^5 + (-56756700Y^8 + 8085646800Y^6 - 92211777000Y^4 + 31696434000Y^2 + 137638966500)T^3 + \\ &(-180180Y^{10} + 39015900Y^8 - 847513800Y^6 + 752787000Y^4 - 3105742500Y^2 + 2173027500)T, \quad f_{18} = 45966045987225T^{12} + \\ &5005Y^{12} + (11623597835850Y^2 - 189852097985550)T^{10} - 1522290Y^{10} + (1076259058875Y^4 - 48183290174250Y^2 + \\ &120557572425675)T^8 + 53961075Y^8 + (44644820220Y^6 - 3863494057500Y^4 + 23308627626900Y^2 - 21898019625300)T^6 - \\ &248327100Y^6 + (808782975Y^8 - 115552275300Y^6 + 1385054984250Y^4 - 1381491814500Y^2 - 21521292262625)T^4 + \\ &279979875Y^4 + (5135130Y^{10} - 1104052950Y^8 + 23701500900Y^6 - 36565735500Y^4 + 80147009250Y^2 - 24936234750)T^2 + \\ &3162300750Y^2 + 3524968125, \quad f_{17} = -190935883331550T^{13} + (-57061298466900Y^2 + 1008082939581900)T^{11} + \\ &(-6457554353250Y^4 + 301021379851500Y^2 - 911707327688850)T^9 + (-344402898840Y^6 + 30333947628600Y^4 - \\ &208974233853000Y^2 + 243008235124200)T^7 + (-8734856130Y^8 + 1251548103960Y^6 - 15817012325100Y^4 + 27698656368600Y^2 + \\ &25683997326750)T^5 + (-92432340Y^{10} + 19730749500Y^8 - 420105155400Y^6 + 1008423675000Y^4 - 1224986206500Y^2 - \\ &146950240500)T^3 + (-270270Y^{12} + 81039420Y^{10} - 2740396050Y^8 + 13812913800Y^6 - 18309989250Y^4 - 192268282500Y^2 - \\ &310728048750)T, \quad f_{16} = 695552146422075T^{14} + 6435Y^{14} + (242510518484325Y^2 - 4607699851202175)T^{12} - \\ &2546775Y^{12} + (32933527201575Y^4 - 1596009395153250Y^2 + 5707633598857575)T^{10} + 120860775Y^{10} + (2195568480105Y^6 - \\ &196756713794025Y^4 + 1532235106692675Y^2 - 2171399870551275)T^8 - 1303717275Y^8 + (74246277105Y^8 - 10668618894780Y^6 + \\ &142516507710150Y^4 - 367316581823100Y^2 - 233903236078575)T^6 + 2575186425Y^6 + (1178512335Y^{10} - 249753960225Y^8 + \\ &5293914711750Y^6 - 18307533431250Y^4 + 11947241194875Y^2 + 7339410799875)T^4 + 23261812875Y^4 + (6891885Y^{12} - \\ &2036817090Y^{10} + 65683505475Y^8 - 379658267100Y^6 + 296991835875Y^4 + 53066024130750Y^2 + 11990299390125)T^2 + \\ &134851240125Y^2 + 46414974375, \quad f_{15} = -2225766868550640T^{15} + (-895423452865200Y^2 + 18206943541592400)T^{13} + \\ &(-143709936879600Y^4 + 7229715286096800Y^2 - 29946939923602800)T^{11} + (-11709698560560Y^6 + 1067384061097200Y^4 - \\ &9313909900218000Y^2 + 15794795481771600)T^9 + (-509117328720Y^8 + 73365112497600Y^6 - 1037922900991200Y^4 + \\ &3582548274110400Y^2 + 1598676194631600)T^7 + (-11313718416Y^{10} + 2380232297520Y^8 - 50421277418400Y^6 + \\ &237401051580000Y^4 - 73786397806800Y^2 - 84105766054800)T^5 + (-110270160Y^{12} + 32114063520Y^{10} - 987433902000Y^8 + \\ &6788816798400Y^6 + 215286498000Y^4 - 78733894428000Y^2 - 281497554450000)T^3 + (-308880Y^{14} + 119805840Y^{12} - \\ &5246957520Y^{10} + 58975938000Y^8 - 11206868400Y^6 - 1359946098000Y^4 - 9136788822000Y^2 - 1877852970000)T, \quad f_{14} = \\ &6259969317798675T^{16} + 6435Y^{16} + (2878146812781000Y^2 - 62359847610255000)T^{14} - 3148200Y^{14} + (538912263298500Y^4 - \\ &28106347270491000Y^2 + 132796272634647300)T^{12} + 172386900Y^{12} + (52693643522520Y^6 - 4884295418818200Y^4 + \\ &47389913750579400Y^2 - 94499857473625800)T^{10} - 3227477400Y^{10} + (2863784974050Y^8 - 413853643942200Y^6 + \\ &6209473531495500Y^4 - 27032928566931000Y^2 - 7792736255214750)T^8 + 514127250Y^8 + (84852888120Y^{10} - \\ &17721199326600Y^8 + 376649696833200Y^6 - 2311896626298000Y^4 + 251203467375000Y^2 + 281809220175000)T^6 + \\ &126059409000Y^6 + (1240539300Y^{12} - 359939353000Y^{10} + 10435219105500Y^8 - 87520569822000Y^6 - 67443242422500Y^4 + \\ &672149733255000Y^2 + 4605837252502500)T^4 + 994204732500Y^4 + (6949800Y^{14} - 2640745800Y^{12} + 106033422600Y^{10} - \\ &1306648665000Y^8 - 1706880357000Y^6 + 26975963385000Y^4 + 291783952215000Y^2 + 16749041085000)T^2 - 1834094745000Y^4 + \\ &637151011875, \quad f_{13} = -1546580654985550T^{17} + (-8058811075786800Y^2 + 185352654743096400)T^{15} + (-1741101158349000Y^4 + \\ &94019462550846000Y^2 - 500392472909502600)T^{13} + (-201193911631440Y^6 + 18958657057578000Y^4 - 203148655154146800Y^2 + \end{aligned}$$

$$\begin{aligned}
& (5670 Y^{26} + 3052350 Y^{24} + 718823700 Y^{22} + 79112092500 Y^{20} + 4620841769250 Y^{18} + 221043771632250 Y^{16} + \\
& 7141636482435000 Y^{14} + 103508757238635000 Y^{12} + 1015327719169706250 Y^{10} - 412507894719768750 Y^8 + 768374499128512500 Y^6 + \\
& 17751118754923312500 Y^4 + 7990846504579968750 Y^2 - 346311617783906250)T^2 - 5861578332093750 Y^2 + 299060118984375, \quad \mathbf{q}_1 = \\
& -2058911320946490 T^{29} + (-3202750943694540 Y^2 + 112096283029308900)T^{27} + (-2313097903779390 Y^4 + 120636952212494340 Y^2 - \\
& 1963820120308702110)T^{25} + (-1028043512790840 Y^6 + 54169985097055800 Y^4 - 1899112689205540200 Y^2 + 13391057556802837800)T^{23} + \\
& (-314124406686090 Y^8 + 11888400622273560 Y^6 - 923152321047757500 Y^4 + 3708205670816703000 Y^2 - 77034760175239533450)T^{21} + \\
& (-69805423708020 Y^{10} + 617509517417100 Y^8 - 318229746639827400 Y^6 - 2614896527607945000 Y^4 - 93336429316395226500 Y^2 - \\
& 347767826255658238500)T^{19} + (-11634237284670 Y^{12} - 384824771723700 Y^{10} - 87149793466987650 Y^8 - 2085383065602979800 Y^6 - \\
& 30755778146008193250 Y^4 + 865772990916574135500 Y^2 + 4306816518950882081250)T^{17} + (-1477363464720 Y^{14} - \\
& 128076048056880 Y^{12} - 1881011900419280 Y^{10} - 723245780863494000 Y^8 - 5720253447152286000 Y^6 + 514177793785476198000 Y^4 + \\
& 564801286096701738000 Y^2 - 16761364588803306402000)T^{15} + (-143632559070 Y^{16} - 21655370444400 Y^{14} - 3069297212272200 Y^{12} - \\
& 157738187047971600 Y^{10} - 1345528904618776500 Y^8 + 97961457598829622000 Y^6 - 6962735025427630005000 Y^4 - \\
& 9074968983265939110000 Y^2 - 17589445369515278358750)T^{13} + (-10639448820 Y^{18} - 2384054954820 Y^{16} - 366632430267600 Y^{14} - \\
& 23432517013726800 Y^{12} - 3592387997640091800 Y^{10} + 4718379529937997000 Y^8 - 289697132657308698000 Y^6 - 3612815584880104890000 Y^4 - \\
& 22341223720366514932500 Y^2 - 18223480285848493582500)T^{11} + (-591080490 Y^{20} - 180506888100 Y^{18} - 31284257630850 Y^{16} - \\
& 2402965397070000 Y^{14} - 63344801316352500 Y^{12} - 1105794713723175000 Y^{10} - 55865245593916552500 Y^8 - 643450736493112590000 Y^6 - \\
& 4748165160242207681250 Y^4 + 15394968789630573937500 Y^2 + 8254010444065136493750)T^9 + (-23882040 Y^{22} - \\
& 9396664200 Y^{20} - 1849930374600 Y^{18} - 167095588678200 Y^{16} - 6623492829534000 Y^{14} - 220140941074866000 Y^{12} - \\
& 69755094704730000 Y^{10} - 81901126660142790000 Y^8 - 6950892686313625000 Y^6 + 272300604511889675000 Y^4 + \\
& 2664473085690924375000 Y^2 - 5448473494388544375000)T^7 + (-663390 Y^{24} - 323121960 Y^{22} - 71929948860 Y^{20} - \\
& 7505303589000 Y^{18} - 401975673293250 Y^{16} - 17425508594130000 Y^{14} - 526836051310653000 Y^{12} - 6559820577961938000 Y^{10} - \\
& 627996662283831250 Y^8 + 87598972192506075000 Y^6 - 296602185070281037500 Y^4 - 1474549884565353225000 Y^2 - \\
& 376367433547989318750)T^5 + (-11340 Y^{26} - 6633900 Y^{24} - 1653258600 Y^{22} - 195980488200 Y^{20} - 13106441722500 Y^{18} - \\
& 662673648514500 Y^{16} - 20887179980310000 Y^{14} - 282058529536350000 Y^{12} - 3018223686271972500 Y^{10} - 4384821363343312500 Y^8 - \\
& 21525129857522025000 Y^6 - 103471877025208125000 Y^4 - 11989971456564937500 Y^2 + 5774359109837062500)T^3 + \\
& (-90 Y^{28} - 61740 Y^{26} - 17013150 Y^{24} - 2253598200 Y^{22} - 176788190250 Y^{20} - 9754930804500 Y^{18} - 318801378750750 Y^{16} - \\
& 4447241995770000 Y^{14} - 52651311898578750 Y^{12} - 180623197653412500 Y^{10} - 526027698056306250 Y^8 - 2572888193311275000 Y^6 - \\
& 772890971503218750 Y^4 - 247781277249187500 Y^2 - 12959271822656250)T, \quad \mathbf{q}_0 = 90 T X^{30} + 205891132094649 T^{30} + \\
& X^{30} + Y^{30} + (343151886824415 Y^2 - 12467851887953745)T^{28} + 855 Y^{28} + (266895911974545 Y^4 - 14412379246625430 Y^2 + \\
& 245811134928555945)T^{26} + 275625 Y^{26} + (128505439098855 Y^6 - 6968948812668675 Y^4 + 257554555055435925 Y^2 - \\
& 1922391943749986625)T^{24} + 44441775 Y^{24} + (42835146366285 Y^8 - 1638718932781980 Y^6 + 136644116908449150 Y^4 - \\
& 560086013795471100 Y^2 + 12730753775566554525)T^{22} + 4060783125 Y^{22} + (10470813556203 Y^{10} - 76517483679945 Y^8 + \\
& 51930597815513550 Y^6 + 446668065138963150 Y^4 + 17134239772133241975 Y^2 + 68083614964471020075)T^{20} + 207533751075 Y^{20} + \\
& (1939039547445 Y^{12} + 72490247696790 Y^{10} + 15872801459062275 Y^8 + 387335760139055700 Y^6 + 4966260292078792875 Y^4 - \\
& 219391313625118514250 Y^2 - 987121566914058583875)T^{18} + 5923312282125 Y^{18} + (277005649635 Y^{14} + 26201893371885 Y^{12} + \\
& 3876302771947935 Y^{10} + 150552748144346625 Y^8 + 611528589858893625 Y^6 - 132315060718915853625 Y^4 + 11896115762319067125 Y^2 + \\
& 5091352640433006100875)T^{16} + 77461769896875 Y^{16} + (30778405515 Y^{16} + 5019247668600 Y^{14} + 727212650994900 Y^{12} + \\
& 37793261862001800 Y^{10} + 214511736599417250 Y^8 - 25126138303645491000 Y^6 + 274776697273223092500 Y^4 + 2735159588700616035000 Y^2 + \\
& 9163014664668879616875)T^{14} + 1691986493491875 Y^{14} + (2659862205 Y^{18} + 642663629685 Y^{16} + 101952332313300 Y^{14} + \\
& 6644086914269700 Y^{12} + 101346793849844550 Y^{10} - 298739643719564250 Y^8 + 126859497950110306500 Y^6 + 1474770758254926772500 Y^4 + \\
& 12353324750841035233125 Y^2 + 4701616168042373698125)T^{12} + 21127132873153125 Y^{12} + (177324147 Y^{20} + 58335097590 Y^{18} + \\
& 1050264260575 Y^{16} + 8324833300912200 Y^{14} + 24428227005948150 Y^{12} + 670167595543868100 Y^{10} + 27909592318881519750 Y^8 + \\
& 284868817499929005000 Y^6 + 2474446523044839564375 Y^4 - 9797539843874926946250 Y^2 + 711575050142526766875)T^{10} + \\
& 60580010182426875 Y^{10} + (8955765 Y^{22} + 3796555455 Y^{20} + 781214825475 Y^{18} + 73798662229425 Y^{16} + 3320791358519250 Y^{14} + \\
& 131842615029441750 Y^{12} + 4001251919029611750 Y^{10} + 38622239875976321250 Y^8 + 354175782842632235625 Y^6 - \\
& 191841030332196253125 Y^4 - 1751845493630807465625 Y^2 + 9443971835072962228125)T^8 + 225021251512378125 Y^8 + \\
& (331695 Y^{24} + 174216420 Y^{22} + 40774449870 Y^{20} + 4511008624500 Y^{18} + 271665478688625 Y^{16} + 12732511726557000 Y^{14} + \\
& 375130032565018500 Y^{12} + 4045763811379941000 Y^{10} + 46685866420475690625 Y^8 + 17034748127560012500 Y^6 + \\
& 644222760196600068750 Y^4 + 3366662881162066762500 Y^2 + 898711974965844309375)T^6 + 50098108080234375 Y^6 + \\
& (8505 Y^{26} + 5372325 Y^{24} + 1416167550 Y^{22} + 180484519950 Y^{20} + 13304000185875 Y^{18} + 682841792091375 Y^{16} + \\
& 20712824439772500 Y^{14} + 255790716490732500 Y^{12} + 3468407818261719375 Y^{10} + 16693527887869021875 Y^8 + 48029919060345468750 Y^6 + \\
& 240167299660771218750 Y^4 - 115468740156070546875 Y^2 + 17221204161424265625)T^4 + 67806897644390625 Y^4 + \\
& (135 Y^{28} + 100170 Y^{26} + 29387925 Y^{24} + 4246092900 Y^{22} + 358633184175 Y^{20} + 19028361381750 Y^{18} + 581147239609125 Y^{16} + \\
& 7773885322875000 Y^{14} + 126384813847648125 Y^{12} + 1109621551203033750 Y^{10} + 3306804953883834375 Y^8 + 12758673814194262500 Y^6 + \\
& 2447767199204578125 Y^4 + 2944864928980406250 Y^2 + 295212212120109375)T^2 + 5881515673359375 Y^2 + 19937341265625.
\end{aligned}$$