



HAL
open science

Chronic ingestion of nitrate and nitrite increases in vitro cholinergic contractility and decreases neutrophil contents of the jejunum in rats

Rafael Garcia Villar, Pauline Anton, Stéphane Marty, Vassilia Theodorou, Jean Fioramonti, Lionel Bueno

► **To cite this version:**

Rafael Garcia Villar, Pauline Anton, Stéphane Marty, Vassilia Theodorou, Jean Fioramonti, et al.. Chronic ingestion of nitrate and nitrite increases in vitro cholinergic contractility and decreases neutrophil contents of the jejunum in rats. Digestive Disease Week / 100th Annual AGA meeting, May 1999, Orlando, United States. 1999, Gastroenterology 116 (4 Part 2) A996. <hal-01600591>

HAL Id: hal-01600591

<https://hal.science/hal-01600591v1>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-SA 4.0 - Attribution - ShareAlike - International License

Dr R. Garcia-Villan

PROGRAM OF THE ANNUAL MEETING OF

THE AMERICAN

GASTROENTEROLOGICAL ASSOCIATION

AND

DIGESTIVE DISEASE WEEK

May 16-19, 1999
Orlando, Florida



This supplement contains the abstracts and program of the American Gastroenterological Association, American Association for the Study of Liver Disease and the Society for Surgery of the Alimentary Tract. Consult the DDW program book for information on the American Society for Gastrointestinal Endoscopy program sessions and abstracts.

A bullet before the title of an abstract (see pp. A1-A1373) signifies that the abstract is intended, at press time, for oral or poster presentation during Digestive Disease Week.

SEE INSTRUCTIONS BEFORE COMPLETING. BE SURE TO COMPLETE BOTH SIDES OF THIS FORM.
TYPE ABSTRACT IN THE SPACE BELOW. USE TYPE NO SMALLER THAN 10-POINT FONT SIZE.

F. ABSTRACT REVIEW DESCRIPTOR

A coded list of abstract descriptors appears on page 4 of this booklet. Enter the code of the descriptor that best fits the subject of your abstract in this box. Select only one descriptor.

72

G. KEY WORDS

Enter the codes for up to four(4) key words in this box. (Listing on pages 5-10)

1. 462 2. 450
3. 590 4. 605
5. 185

Add a keyword you would like added to future lists on line #5.

H. CLINICAL RESEARCH

Indicate if this research is clinical.

Clinical Basic

I. POSTER ONLY

Check below if you want your abstract presented only as a poster

J. CERTIFICATION BY PRINCIPAL AUTHOR

I certify that this work and any substantial part herein has not been submitted concurrently to other DDW societies, will not be published as an article before May 16, 1999, and has not been presented at other U.S. national meetings. I further certify that the studies described in the abstract that involved human subjects conformed to the principles of the Declaration of Helsinki of the World Medical Association (Clinical Research 1966; 14:03) and met all of the requirements governing informed consent of the country in which they were performed.

CHRONIC INGESTION OF NITRATE AND NITRITE INCREASES IN VITRO CHOLINERGIC CONTRACTILITY AND DECREASES NEUTROPHIL CONTENT OF THE JEJUNUM IN RATS. R. Garcia-Villar, P. Anton, S. Marty, V. Theodorou*, J. Fioramonti, L. Buéno. INRA, Digestive Pharmacology Unit; *ESAP, Toulouse, France.

Nitrate and nitrite present in foods and drinking water subserve a controversial human health issue. Thus, consumers blame these compounds and their byproducts (nitrosamines, nitric oxide) for insidious functional and/or structural alterations of the digestive tract that may lead to more severe pathologies.

Specific aims: The present study was therefore undertaken to investigate the influence on jejunal contractility, mast cells, and neutrophils, of a two-week oral administration of nitrate or nitrite (sodium and potassium salts) to rats at doses possibly found in food.

Methods: Five groups of 8 male Wistar rats (250-300 g) received respectively once daily by intragastric gavage (mg/kg/day; treated groups): NaNO₃ (500), KNO₃ (250), NaNO₂ (8), KNO₂ (5) and distilled water (1ml/day; controls). After 14 days rats were sacrificed; segments of jejunum (20 cm aboral to the LT) were collected for: (i) myeloperoxidase (MPO) spectrophotometric assay for neutrophil infiltration; (ii) mast cell numbers determination (alcian blue/safranin staining); (iii) in vitro contractility studies (construction of concentration-contractile responses curves to carbachol (CCH; a muscarinic cholinergic agonist), serotonin (5-HT), and neurokinin A (NKA); 0.1 nM-1 mM).

Results: MPO activity in jejunal tissue was reduced to less than 15% of controls (p<0.05) in rats treated with nitrate but not in those treated with nitrite. No differences in jejunal mast cell numbers were found between the five groups. Maximal contraction induced by CCH was significantly increased reaching 153-159% of controls (p<0.05), while no differences were measured in the responses to 5-HT and NKA, in the four treated groups compared with controls.

Conclusion: Ingestion of nitrate or nitrite for two weeks enhanced jejunal smooth muscle responsiveness to cholinergic, but not to serotonergic or tachykinergic stimulation in rats. It can be suggested that alteration of parasympathetic control of GI tract motility and neutropenia resulting from chronic ingestion of nitrate may pave the way for intestinal functional disorders.

This research was made possible by a grant of Ministère de l'Environnement (France) and the institutional support of INRA.

I affirm that this abstract meets all of the requirements stated in this booklet, that the authors named are familiar with the presented data, and that Box B has been signed. On behalf of all of the authors, I hereby transfer copyright in the above abstract to American Gastroenterological Association.

Signature Rafael Garcia-Villar Print Name Rafael GARCIA-VILLAR

Phone +33 561285151 Fax +33 561285310

M. MAILING

Submit original abstract to:
AGA
c/o Medical Support Systems
1000 Massachusetts Ave., 3rd flr
Cambridge, MA 02138
Phone: 617/868-7767

G4325

QUANTIFICATION OF CAECOILEAL REFLUX BY GAMMA-SCINTIGRAPHY IN CONSCIOUS PIGS.

Cuche gaelle, Malbert Charles-Henri, Station de Ress Porcines, Saint-Gilles France

We evidenced recently frequent caecoileal reflux events in pigs (Cuche and Malbert, 1998). However, the method used, i. e. measurement of intraluminal ileal pH, was not quantitatively informative. Thus, this study aimed to quantify the amount of caecal fluids invading terminal ileum using gamma scintigraphy imaging in conscious pigs. Under anaesthesia and aseptic conditions, an infusion silicone tube (ID 5 mm) was inserted in the apex of the caecum and a Y-shaped silicone cannula was located 20 cm proximal to the caecoileal sphincter in 10 pigs (30 ± 2 kg). The animals were allowed to recover for a week after surgery. While the pig was standing in a sling, a guide (ID 10 mm) filled with 40 MBq ⁶⁷Ga was introduced into the terminal ileum pointing towards the caecum. This guide will serve to outline the ileal lumen on gamma scans. 200 ml saline labelled with 40 MBq ^{99m}Tc-DTPA were infused in the caecum at 40 ml.min⁻¹. Repetitive one minute scans of the ileocecolonic area were acquired in multi-isotope mode for 30 minutes with high resolution collimator (Apex 500G). Since porcine ileum follows the spiral shape of the colon, the only practicable incidence allowing simultaneous imaging of ileum and caecum was from the rear of the animal. For each frame, the ileum was outlined on the Ga scan and the number of counts (Ic) within a copy of this ROI on the Tc scan was evaluated. Cecoileal reflux was expressed as Ic divided by the total number of counts on the Tc scan. A significant amount of Tc radioactivity was found in the ileal ROI during 4 ± 3 min on each pig. It is uncommon to observe more than one episode during the recording session. None of these refluxes invade the ileum 15 cm proximal to the caecoileal junction and 50 % of them were limited to the first 5 cm of the ileum. For each event, reflux amounted to 13 ± 2 %, a value remarkably constant along frames once the reflux event was detected. In conclusion, this work confirms the existence of cecoileal reflux in pigs. They amount about 13 % of caecal contents. Cuche G., Malbert, C. H. 1998. Relationships between caecoileal reflux and ileal motor patterns in conscious pigs. American Journal of Physiology, 274, G35-G40, 1998.

● G4326

CHRONIC INGESTION OF NITRATE AND NITRATE INCREASES IN VITRO CHOLINERGIC CONTRACTILITY AND DECREASES NEUTROPHIL CONTENT OF THE JEJUNDUM IN RATS

R Garcia-Villal, INRA, Digest Pharmacology Unit, Toulouse Cedex 9 France; P Anton, INRA, Toulouse France; S Marty, INRA, Digest Pharmacology Unit, Toulouse France; V Theodorou, ESAP, Toulouse France; J Fioramonti, L Bueno, INRA, Digest Pharmacology Unit, Toulouse France

Nitrate and nitrite present in foods and drinking water subserve a controversial human health issue. Thus, consumers blame these compounds and their byproducts (nitrosamines, nitric oxide) for insidious functional and/or structural alterations of the digestive tract that may lead to more severe pathologies. **Specific aims:** The present study was therefore undertaken to investigate the influence on jejunal contractility, mast cells, and neutrophils, of a two-week oral administration of nitrate or nitrite (sodium and potassium salts) to rats at doses possibly found in food. **Methods:** Five groups of 8 male Wistar rats (250-300 g) received respectively once daily by intragastric gavage (mg/kg/day; treated groups); NaNO₃ (500), KNO₃ (250), NaNO₂ (8), KNO₂ (5) and distilled water (1 ml/day; controls). After 14 days rats were sacrificed; segments of jejunum (20 cm aboral to the LT) were collected for: (i) myeloperoxidase (MPO) spectrophotometric assay for neutrophil infiltration; (ii) mast cell numbers determination (alcian blue/safranin staining); (iii) in vitro contractility studies (construction of concentration-contractile responses curves to carbachol (CCH; a muscarinic cholinergic agonist), serotonin (5-HT), and neurokinin A (NKA); 0.1 nM-1 mM). **Results:** MPO activity in jejunal tissue was reduced to less than 15% of controls (p<0.05) in rats treated with nitrate but not in those treated with nitrite. No differences in jejunal mast cell numbers were found between the five groups. Maximal contraction induced by CCH was significantly increased reaching 153-159% of controls (p<0.05), while no differences were measured in the responses to 5-HT and NKA, in the four treated groups compared with controls. **Conclusion:** Ingestion of nitrate or nitrite for two weeks enhanced jejunal smooth muscle responsiveness to cholinergic, but not to serotonergic or tachykinergic stimulation in rats. It can be suggested that alteration of parasympathetic control of GI tract motility and neutropenia resulting from chronic ingestion of nitrate may pave the way for intestinal functional disorders.

G4327

ROLE OF TACHYKININ NK₁, NK₂ AND NK₃ RECEPTORS IN MODULATION OF VISCERAL HYPERSENSITIVITY IN THE RAT.

Genevieve-Anne Gaudreau, Janie McDuff, Victor Plourde, CHUM, Saint-Luc - Univsite de Montreal, Montreal, PQ Canada

Tachykinins are involved in visceral pain but the relative contribution of tachykinin receptors is unclear. In animal models of visceral sensitization, repetitive colorectal distensions (CDs) result in visceral afferent nerve sensitization. **AIMS:** To investigate the effects of NK₁, NK₂ and NK₃ receptor blockade in visceral hypersensitivity induced by CDs. **METHODS:** Visceral pain thresholds (VPT) were determined in male Sprague Dawley rats before or following a sensitization paradigm consisting of 10 repetitive CDs (duration 1 min followed by 30 sec rest) at 30 or 50mmHg. VPT were evaluated by measuring the pressure threshold (increments of 5mmHg, duration 3 min, 1 min rest) required to evoke the first abdominal muscle contraction (mean of 3 measurements). NK receptors antagonists, RP67580 (6.5 nmol), SR48968 (6.5 nmol) and R820 (6.5 nmol), acting at NK₁, NK₂ and NK₃ receptors respectively, were administered intrathecally 15 min prior to the assessment of VPT. **RESULTS:** VPT (31.0±11.0mmHg) were significantly lowered following either 30 or 50mmHg CDs. VPT were significantly increased by RP67580 in rats sensitized with 30 (12.9±2.5 to 27.7±3.9 mmHg; p<0.05) or 50mmHg (12.61.7± to 32.1±8.8 mmHg; p<0.05). VPT were also significantly increased by R820 in rats sensitized with 30 (14.0±2.9 to 28.7±8.2mmHg; p<0.05) or 50mmHg (16.3±2.3 to 33.2±10.6 mmHg; p<0.05). SR48968 (≤ 65 nmol) did not affect VPT. **SUMMARY:** Repetitive CDs at 30 or 50mmHg significantly lowered VPT suggesting visceral sensitization and this effect was reversed by NK₁ and NK₃ receptor antagonists but unaffected by NK₂ blockade. **CONCLUSIONS:** Our results suggest that tachykinins with actions at spinal NK₁ and NK₃ receptors are involved in visceral sensitization induced by repetitive CDs.

G4328

INVOLVEMENT OF NMDA RECEPTORS IN RAT VISCERAL HYPERALGESIA.

Genevieve-Anne Gaudreau, Victor Plourde, CHUM, Saint-Luc - Univsite de Montreal, Montreal, PQ Canada

N-methyl-D-aspartate (NMDA) receptors are involved in spinal transmission of nociceptive input. In the rat, repetitive colorectal distensions (CDs) result in visceral afferent nerve sensitization. **AIMS:** To investigate a possible role of NMDA receptors in the development of visceral hypersensitivity. **METHODS:** Visceral pain thresholds were determined in male Sprague Dawley rats before or following a sensitization paradigm consisting of 10 repetitive CDs (duration 1 min followed by 30 sec rest) at 30 or 50mmHg. Visceral pain thresholds were evaluated by measuring the pressure threshold (increments of 5mmHg, duration 3 min, 1 min rest) required to evoke the first abdominal muscle contraction (mean of 3 measurements). To assess the role of NMDA receptors, the NMDA receptor antagonist, MK-801, was administered intraperitoneally (i.p.; 30nmol/kg) or intrathecally (i.t.; 10nmol) under light halothane anesthesia, 15 min before the sensory thresholds were determined in conscious unanesthetized animals. **RESULTS:** Normal visceral pain thresholds (31.0±11.0mmHg) were significantly lowered following either 30 or 50mmHg CDs. MK-801 administered i.p. to rats sensitized by 30mmHg increased the pain threshold from 17.0±3.3 to 30.8±7.5mmHg (p<0.05) and in rats sensitized by 50mmHg from 13.3±2.5 to 22.3±6.9mmHg (p<0.05). Following i.t. MK-801 administration, pain thresholds in rats sensitized with 30mmHg were significantly increased from (13.9±2.0)mmHg to (27.0±9.0mmHg) (p<0.05) and with 50mmHg from (15.0±3.9)mmHg to (25.9±6.9)mmHg; p<0.05). **SUMMARY:** Both 30 and 50mmHg-induced hypersensitivity was abolished by i.p. or i.t. MK-801. **CONCLUSIONS:** Those results support a role for NMDA receptors in visceral hypersensitivity induced by repetitive CDs.

● G4329

NEMATODE-INDUCED HYPERMASTOCYTOSIS MODIFIES THE JEJUNAL MOTOR RESPONSE TO CCK-8 IN RATS: ROLE OF MAST CELLS AND CCK RECEPTORS

J Gay, J Fioramonti, R Garcia-Villar, L Bueno, INRA, Digest Pharmacology Unit, Toulouse France

Intestinal hypermastocytosis is a common feature of various digestive disturbances such as inflammatory bowel diseases. In rats, jejunal hypermastocytosis accompanied by enteric nerves remodeling can be observed during several weeks after infection with the nematode *Nippostrongylus brasiliensis* (Nb). The aim of this study was, first to evaluate whether jejunal motor responses to CCK-8 were modified by post-infection hypermastocytosis, and second to determine the CCK receptors involved in the motor response. **METHODS:** 8 groups of 8 male Wistar rats were used: 4 control groups were matched with 4 groups infected at day 0 with 3500 third-stage larvae of Nb (in 0.5 ml sterile water; sc). At day 30, 3 groups of NiCr electrodes were implanted 10 cm apart in mid-jejunum. Myoelectrical recordings started 7 days after surgery. The duration of disruption of the migrating myoelectrical complex (MMC) pattern was evaluated in both control and post-infected groups after CCK-8 (50 µg/kg, ip), under baseline