



The plant cuticle as an interface between leaves and air-borne pollutants

M. Riederer, Pierre-Marie Badot, Jean-Pierre Garrec, Béatrice Richard, L. Schreiber, P. Sümmchen, M. Uhlig, O. Wienhaus

► To cite this version:

M. Riederer, Pierre-Marie Badot, Jean-Pierre Garrec, Béatrice Richard, L. Schreiber, et al.. The plant cuticle as an interface between leaves and air-borne pollutants. EUROSILVA Conference - Contribution to Forest Tree Physiol, Nov 1994, Dourdan, France. pp.101-118. hal-01601031

HAL Id: hal-01601031

<https://hal.science/hal-01601031>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The plant cuticle as an interface between leaves and air-borne pollutants

M. RIEDERER¹, P. M. BADOT², J. P. GARREC³, B. RICHARD³, L. SCHREIBER¹,
P. SUMMCHEH⁴, M. UHLIG⁴, O. WIENHAUS⁴

¹ Julius-von-Sachs-Institut für Biowissenschaften mit Botanischem Garten, Lehrstuhl für Botanik II, Universität Würzburg, Mittlerer Dallenbergrweg 64, D-97082 WÜRZBURG, Germany

² Laboratoire de Sciences Végétales, Institut des Sciences et Techniques de l'Environnement, Place Leclerc, F-25030 BESANÇON CEDEX, France

³ Laboratoire Pollution Atmosphérique, INRA, Centre de Recherches Forestières de Nancy, F-54280 CHAMPENOUX, France

⁴ Institut für Pflanzenchemie und Holzchemie, Technische Universität Dresden, Pierner Str. 21, D-01737 THARANDT, Germany

Abstract

Material deposited on leaf surfaces may be taken up into the interior either by the stomatal or the cuticular pathway. The uptake of non-volatile substances is restricted to the penetration of the cuticle. The objective of the collaborative work reported here was to characterise important aspects of the interactions between air-borne pollutants and the cuticle of trees. The amounts and composition of the cuticular waxes of the needles of *Picea abies* and the leaves of *Fagus sylvatica* were investigated with special attention to the effects of age and environmental factors. The transport properties of plant cuticles were studied using a variety of techniques covering uptake into intact needles, permeability of isolated cuticular membranes and the mobility of permeants in reconstituted cuticular waxes. Effects of polluted sea-spray on the needle surfaces of conifers growing close to the Mediterranean Sea were also investigated.

Keywords: air pollution, *Citrus aurantium*, cuticle, cuticular waxes, *Fagus sylvatica*, *Hordeum vulgare*, permeability, *Picea abies*, *Pinus halepensis*, sea spray, surfactants