



Is there a synergy between hedges and intercrops for pest biocontrol ?

Audrey Alignier, Emilie Andrieu, Laurent Bedoussac, Etienne-Pascal Journet, Annie Ouin, Jean-Pierre Sarthou, Aude Vialatte, Antoine Brin

► To cite this version:

Audrey Alignier, Emilie Andrieu, Laurent Bedoussac, Etienne-Pascal Journet, Annie Ouin, et al.. Is there a synergy between hedges and intercrops for pest biocontrol ?. 11. International Association for Ecology congress, Aug 2013, Londres, United Kingdom. 1 p., 2013. hal-01601774

HAL Id: hal-01601774

<https://hal.science/hal-01601774>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Is there a synergy between distance from the hedge and intercrop for pest biocontrol?

A. Alignier^{1,2,3}, E.Andrieu², L.Bedoussac^{3,4}, E.-P. Journet⁴, A.Ouin^{1,2}, J.-P. Sarthou^{1,3}, A.Vialatte^{1,2} & A.Brin^{2,5}

Contact: audrey.alignier@toulouse.inra.fr

Identifying alternative cropping systems: a major challenge in agroecology

To achieve sustainability, we need to identify alternative cropping systems that can both ensure sufficient levels of food production and reduction in pesticide inputs while maintaining ecosystem integrity and simultaneously delivering multiple ecosystem services.

What is intercropping ?



Intercropping creates structures of mixed plant populations with "barrier" or "dilution of resources" effects which can reduce the pressure of pathogens, weeds and pests compared with that exerted on sole crops.



How to promote biological control by natural enemies in crop fields ?

Promoting relevant management practices of semi-natural elements, i.e. hedgerows

Increase crop diversity both in time and space (rotation and intercropping respectively)


Combination = synergy ?

Absence of a global synergy between distance from the hedgerow and intercrop in supporting biocontrol of pests and weeds

	Aphid predation	Slug predation	Weed seed predation	Sitona eggs predation	Bruchids predation	Nb of pest enemies
Distance	+	-	-	ns	-	+
Crop [pea]	ns	-	+	-	-	+
Crop [intercrop]	ns	-	-	na	-	+
Interaction	-	ns	+/-	+	+	-

Results of GLMMs including Distance, Type of Crop (sole crops vs intercrop) and their interaction as fixed effect and the position of the survey as random effect (when available). na: not available; ns: not significant; +/- significant but depending on the weed species considered.

Field surveys of pest abundance and effectiveness of biological control





One field separated in 3 blocks:
 -Pea (sole crop)
 -Wheat (sole crop)
 - Pea and Wheat (intercrop)

Random sampling design according to the distance from the hedgerow.

Prey (aphids, slugs, eggs of *Sitona*) and weed seeds were stuck on sandpaper cards.

Pea bruchids were sampled on 10 cloves infields, at each distance. The number of bruchids and parasitoids were counted.

flora surveys and cards position



A great variability in pest biocontrol was observed. This study calls for a deepened **multi-species approach** in order to find **trade-offs** between agricultural practices and field edge management to **optimize regulating ecosystem services** such as pest biocontrol.

This work was conducted under the BQR project HaiesCAFARR.