

Impact of exercise intensity on 24h energy balance in obese children: an exploration in calorimetric chambers

David Thivel, Christophe Montaurier, Sylvie Rousset, Béatrice Morio, P.

Duché

▶ To cite this version:

David Thivel, Christophe Montaurier, Sylvie Rousset, Béatrice Morio, P. Duché. Impact of exercise intensity on 24h energy balance in obese children: an exploration in calorimetric chambers. 8. Journées Francophones de Nutrition (JFN), Dec 2010, Lille, France. Elsevier Mason SAS, Nutrition Clinique et Métabolisme, 2010, Nutrition Clinique et Métabolisme. hal-01602033

HAL Id: hal-01602033 https://hal.science/hal-01602033v1

Submitted on 3 Jun2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Impact of exercise intensity on 24h energy balance in obese children : an exploration in calorimetric chambers

D. Thivel^{1,2}, C. Montaurier¹, S. Rousset¹, B. Morio¹ and P. Duché²

¹Lipidic and Energetic Metabolism Team, UMR 1019 Human Nutrition, INRA; Auvergne University, France. ²Laboratory of Exercise Physiology, EA3533, Clermont Universities, Blaise Pascal, France.

Introduction

Physical exercise can dually impact the energy balance by increasing energy expenditure and indirectly modify energy intake. The actual available data mostly concern healthy adults or athletes but very few consider obese people, particularly youths.

Objective

To examine whether intensive vs. moderate or sedentary exercise can decrease spontaneous energy intake and thus induce negative 24h energy balance in obese adolescents.


Methods

15 obese adolescents (12-15 years old)

Body composition (DXA) and maximal oxygen consumption (VO₂max) were assessed. The participants entered a metabolic chamber at 3 different occasions randomly assigned (Fig1):

- SED: Sedentary day

- LIE: Low Intensity Exercise day (40%VO₂max)
- HIE: High Intensity Exercise day (75%VO2max)



The cycling exercises during LIE and HIE were individually isoenergetics (330±20 Kcal). 24h energy intake (*ad libitum*), expenditure and appetite sensations (using Visual Analogue Scales) were assessed.

Results




Figure 2. LIE, HIE and SED 24h Energy Expenditure, Intake and Balance results (*p<0.05; **p<0.01)

Total energy intake at lunch and dinner was significantly after intensive exercise in comparison to moderate exercise and sedentary sessions (p<0.01). Energy intake at dinner was the most affected after intensive exercise. Energy intake the next morning at breakfast was similar between all conditions.

Energy balance was significantly reduced after intensive exercise in comparison with moderate exercise and sedentary sessions (p<0.05).

Subjective appetite rates were similar between sessions.

Conclusion

By dually affecting energy expenditure and energy intake, intensive exercise favors a decreased energy balance in comparison with moderate exercise and sedentary conditions. This is associated with no change in appetite sensations, suggesting that adolescents are not at risk of food frustration.

References

Martins C, Morgan L, Truby H. A review of the effects of exercise on appetite regulation: an obesity perspective. Int J Obes (Lond) 2008; 32: 1337-1347.

Thivel D., et al. Intensive exercise: a remedy for childhood obesity. Physiology and Behavior 2010. In press.

Blundell JE, King NA. Physical activity and regulation of food intake: current evidence. Med Sci Sports Exerc 1999; 31: S573-583