



HAL
open science

Cereulide production by *Bacillus weihenstephanensis* strains during growth at different pH values and temperatures

Alizée Guérin, Helene Thorsen Rønning, Claire Dargaignaratz, Thierry Clavel, Veronique Broussolle, Jacques Mahillon, Per Einar Granum, Christophe Nguyen-The

► To cite this version:

Alizée Guérin, Helene Thorsen Rønning, Claire Dargaignaratz, Thierry Clavel, Veronique Broussolle, et al.. Cereulide production by *Bacillus weihenstephanensis* strains during growth at different pH values and temperatures. *Food Microbiology*, 2017, 65, pp.130-135. 10.1016/j.fm.2017.02.006 . hal-01602612

HAL Id: hal-01602612

<https://hal.science/hal-01602612v1>

Submitted on 12 Oct 2021


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Cereulide production by *Bacillus weihenstephanensis* strains during growth at different pH values and temperatures

Alizée Guérin, Helene Thorsen Rønning, Claire Dargaignaratz, Thierry Clavel, Véronique Broussolle, Jacques Mahillon, Per Einar Granum, Christophe Nguyen-The



PII: S0740-0020(16)30884-X

DOI: [10.1016/j.fm.2017.02.006](https://doi.org/10.1016/j.fm.2017.02.006)

Reference: YFMIC 2739

To appear in: *Food Microbiology*

Received Date: 19 October 2016

Revised Date: 13 February 2017

Accepted Date: 15 February 2017

Please cite this article as: Guérin, A., Rønning, H.T., Dargaignaratz, C., Clavel, T., Broussolle, V., Mahillon, J., Granum, P.E., Nguyen-The, C., Cereulide production by *Bacillus weihenstephanensis* strains during growth at different pH values and temperatures, *Food Microbiology* (2017), doi: 10.1016/j.fm.2017.02.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 Cereulide production by *Bacillus weihenstephanensis* strains during growth
2 at different pH values and temperatures
3
4

5 Alizée Guérin¹, Helene Thorsen Rønning³, Claire Dargaignaratz¹, Thierry Clavel¹,

6 Véronique Broussolle¹, Jacques Mahillon², Per Einar Granum³ and Christophe Nguyen-the^{1*}
7

8 ¹UMR408 SQPOV « Sécurité et Qualité des Produits d'Origine Végétale », INRA,
9 University of Avignon, 84000 Avignon, France

10 ²Laboratory of Food and Environmental Microbiology, Université catholique de Louvain,
11 Louvain-la-Neuve, Belgium

12 ³Norwegian University of Life Sciences, NMBU School of Veterinary Science, Oslo,
13 Norway
14

15 *Corresponding author. Mailing address: INRA, UMR408, site Agroparc, 84914 Avignon
16 cedex 9, France. Phone: +33 (0)4 32 72 25 21, Fax: +33 (0)4 32 72 24 92, Email:

17 christophe.nguyen-the@inra.fr
18

19 **Abstract**

20

21 Besides *Bacillus cereus*, some strains of the psychrotolerant, potentially foodborne pathogen
22 *Bacillus weihenstephanensis* can produce the emetic toxin (cereulide). This toxin is a heat-
23 and acid-stable cyclic dodecadepsipeptide that causes food intoxication with vomiting.
24 However, some severe clinical cases with lethal outcomes have been described. If cereulide
25 can be produced during refrigerated storage, it will not be inactivated by reheating food,
26 representing an important risk of food intoxication for consumers. In this paper, we
27 determined the capacity of the *B. weihenstephanensis* strains BtB2-4 and MC67 to grow and
28 produce cereulide on agar media at temperatures from 8 °C to 25 °C and at a pH from 5.4 to
29 7.0. At 8 °C, strain BtB2-4 produced quantifiable amounts of cereulide, whereas the limit of
30 detection was reached for strain MC67. For BtB2-4, cereulide production increased 5-fold
31 between 8 °C and 10-15 °C and by more than 100-fold between 15 °C and 25 °C. At
32 temperatures of 10 °C and higher, cereulide concentrations were within the range of those
33 reported by previous works in foods implicated in emetic poisoning. At 25 °C, decreasing the
34 pH to 5.4 reduced cereulide production by strain BtB2-4 by at least 20-fold.

35

36

37 Key words: *Bacillus cereus*; psychrotolerant; cold; acid; emetic toxin

38

39 1 Introduction

40 Cereulide is the emetic toxin produced by some strains of the spore-forming pathogen
41 *Bacillus cereus* (Ehling-Schulz et al., 2004; Stenfors Arnesen et al., 2008). Cereulide is a
42 heat- and acid-stable cyclic dodecadepsipeptide, [D-O-Leu-D-Ala-O-Val-L-Val]₃, with a
43 molecular mass of 1,153 Da (Agata et al., 1995; Teplova et al., 2006). Cereulide is
44 synthesized by a non-ribosomal peptide synthase encoded by the cereulide synthase gene
45 (*ces*) cluster (Ehling-Schulz et al., 2005). The cluster is located on the megaplasmid pCER270
46 which is related to the toxin plasmid pXO1 of *Bacillus anthracis* (Ehling-Schulz et al., 2015;
47 Ehling-Schulz et al., 2006a; Hoton et al., 2005). The effects of cereulide on the human body
48 have not yet been completely elucidated. In the duodenum, this toxin acts on the 5-HT₃
49 receptor and causes vomiting by stimulation of the afferent vagus nerve 0.5-6 h after the
50 ingestion of contaminated food (Agata et al., 1995; Ehling-Schulz et al., 2004). It acts as a K⁺
51 ionophore with a mechanism of action similar to that of valinomycin. It interferes with
52 oxidative phosphorylation on mitochondrial membranes (Hoton et al., 2005; Mikkola et al.,
53 1999); (Paananen et al., 2002; Teplova et al., 2006). Cereulide has been involved in severe
54 clinical cases, leading in some cases to death of the patient (Dierick et al., 2005; Mahler et al.,
55 1997; Naranjo et al., 2011).

56 The majority of the *B. cereus* strains producing cereulide are mesophilic (Carlin et al., 2006)
57 and belong to phylogenetic group III, growing between 15 °C and 45 °C (Guinebretière et al.,
58 2008). These strains are not able to grow and produce cereulide during cold storage of food.
59 However, some psychrotolerant *Bacillus weihenstephanensis* strains, belonging to the *B.*
60 *cereus* group and classified into phylogenetic group VI (Guinebretière et al., 2008;
61 Guinebretiere et al., 2010), can produce cereulide (Hoton et al., 2009; Thorsen et al., 2006).
62 These psychrotolerant cereulide-producing strains represent a potentially serious food safety
63 risk for consumers of cooked-refrigerated foods. Indeed, if these strains grow and produce

64 cereulide during food storage at low temperatures, the emetic toxin will not be inactivated by
65 food reheating before consumption. However, the production of cereulide at low temperatures
66 by these psychrotolerant strains is still debated. For instance, a recent study showed that
67 *B. weihenstephanensis* MC67 grew at 8 °C and produced approximately 230 ppb of cereulide
68 per bacterial wet biomass after 9 days on PCA (Rossvoll et al., 2014). In contrast, a previous
69 study showed that this strain produced cereulide at a level close to the limit of detection at
70 8 °C after 1, 2 or 3 weeks on BHI-agar (Thorsen et al., 2009a). Furthermore, another
71 cereulide-producing strain, *B. weihenstephanensis* M118, did not produce cereulide in cooked
72 meat sausages stored at 8 °C (Thorsen et al., 2009b). At higher temperatures, the quantity of
73 cereulide produced varied according to the food products and seemed to depend on the food
74 pH. For example, a mesophilic emetic *B. cereus* strain produced 2.4 µg/g of cereulide in liver
75 sausages at pH 6.2 and 1.0 µg/g in a quark dessert at pH 5.1 (Dommel et al., 2010).
76 All studies to date regarding cereulide production by *B. weihenstephanensis* have been
77 performed using the psychrotolerant MC 67 and MC118 strains, which are phylogenetically
78 very similar, both harbouring the *ces* genes on a comparable plasmid (Mei *et al.* 2014).
79 The goal of the present study was to determine the impact of growth temperature and pH on
80 cereulide production by phylogenetically distant *B. weihenstephanensis* strains.(Hoton et al.,
81 2009; Rossvoll et al., 2014; Thorsen et al., 2006)(Hoton et al., 2009; Mei et al., 2014)

82

83 2 Materials and methods

84 2.1 Strains

85 The cereulide-producing strains used in this study were *B. weihenstephanensis* MC67,
86 isolated from soil in a sandy loam on the island of Møn, Denmark, and *B. weihenstephanensis*
87 BtB2-4, isolated from a forest soil (Hoton et al., 2009; Thorsen et al., 2006). Among the
88 cereulide producing *B. weihenstephanensis*, these two strains are phylogenetically distant

89 (Hoton et al., 2009; Mei et al., 2014). *B. weihenstephanensis* KBAB4, which was isolated
90 from forest soil and belongs to phylogenetic group VI, was used as a non cereulide-producing
91 strain (Sorokin et al., 2006; Vilas-Boas et al., 2002).

92 2.2 Growth conditions

93 Growth conditions were adapted from (Rossvoll et al., 2014): briefly, the stock cultures
94 consisted of exponential phase cells (OD₆₀₀ of 0.5) in 30 % glycerol, stored at -80 °C. Frozen
95 suspensions were streaked onto plate count agar (PCA; Oxoid) and incubated at 30 °C
96 overnight. Three colonies were transferred to 10 ml Tryptic Soy Broth (TSB; Oxoid) in
97 KIMAX tubes and cultured for 18 to 20 h at 20 °C while shaking at 200 rpm. The cultures
98 were then incubated at 4 °C for 18-21 h for cold adaptation prior to growth (Rossvoll et al.,
99 2014).

100 Bacterial growth occurred on PCA plates and on tryptic soy agar plates at 10 °C (TSA;
101 Sigma-aldrich). The pH of the PCA plates was 7.0 ± 0.2 and was adjusted to 6.0, 5.6 or 5.4
102 with 1 N or 6 N HCl when necessary, 5.4 being the lowest pH permitting growth of *B.*
103 *weihenstephanensis* at 8-10°C (Guérin et al., 2016).

104 Each PCA plate was inoculated with 100 µl of a 10³ CFU/ml dilution of the inoculum stored
105 at 4 °C, which resulted in approximately 10² cells per plate. The agar plates were incubated at
106 temperatures varying between 8 °C and 25 °C. For each condition, the test was performed
107 three times with an independently prepared inoculum (biological triplicates). Plates were
108 removed daily for the determination of viable cells. As 95% of cereulide is associated with the
109 bacterial biomass (Thorsen et al., 2009a), the cells were collected from the agar surface using
110 2 ml of sterile water and a sterile Drigalski spatula. An aliquot of 100 µl of each suspension
111 was used and plated in serial dilutions on Luria Bertani plates (LB; Biokar), and then the cells
112 were incubated at 30 °C. The rest of the suspension was stored at -20 °C for cereulide
113 extraction.

114 2.3 Extraction of cereulide

115 For cereulide extraction (Ronning et al., 2015), 1.4 ml of all samples were transferred into 2-
116 ml non-skirted microtubes with O-ring seal caps (Dutscher) with 0.3 g of 0.1 mm silica
117 spheres (MP Biomedical), and the cells were disrupted for 45 s at 1500 beats per min (bpm)
118 using a Mini-GTM (SPEX[®] SAMPLEPREP). After centrifugation (3 min at 19,000 × g), 1 ml
119 of each sample supernatant was recovered in 10 ml glass tubes. Ten nanograms of an internal
120 standard, ¹³C₆-cereulide (> 95% purity, 20 µg/ml in acetonitrile; Chiralix), and 1 ml of
121 methanol LC-MS Grade (Fischer chemical) were added to each glass tubes. All samples were
122 vigorously shaken for 30 s and evaporated to dryness by heating at 80 °C under a steady flow
123 of nitrogen. Then, 1 ml of methanol was added, and all samples were vigorously shaken for
124 30 s and centrifuged 5 min at 5,000 × g. The supernatant was recovered in a new glass tube
125 before a second evaporation step at 80 °C under a nitrogen flow. When the samples were
126 completely dry, 100 µl of methanol/sterile water (3/1 vol/vol) was added to the glass tubes,
127 which were immediately vortexed for 30 s. The 100-µl samples were spun down and
128 transferred through a 0.2-µm nylon Micro centrifuge filter (Sigma-Aldrich) and centrifuged 1
129 min at 10, 000 × g. Filtrates were then transferred into HPLC vials (screw vials 8-mm amber
130 glass 12*32 mm with label; Interchim) containing a 0.1-ml micro insert (Sigma-Aldrich) and
131 stored at -20 °C before analysis by LC-MS/MS.

132 2.4 Calibration curve

133 The calibration curve, for quantification of cereulide in the samples, was performed by adding
134 cereulide (cereulide of > 95% purity, 50 µg/ml in acetonitrile; Chiralix) (Biesta-Peters et al.,
135 2010) to the fresh cell biomass of the non-cereulide producer *B. weihenstephanensis* KBAB4
136 at final concentrations of 0.1, 1.0, 5.0, 10.0, 50.0 and 100.0 ng/ml (Ronning et al., 2015).
137 KBAB4 cells were grown on PCA plates at 30 °C overnight as described above. The cells
138 were collected from the plates, disrupted and centrifuged as described above for the emetic

139 strains. One millilitre of this supernatant was supplemented with cereulide and underwent the
140 subsequent steps of the cereulide extraction protocol described above, including the addition
141 of $^{13}\text{C}_6$ -cereulide as an internal standard. The 18 samples of KBAB4 biomass were divided
142 into the six cereulide concentration levels. Several controls were used for analysis, including
143 extracts of KBAB4 cells alone (negative control) and extracts of KBAB4 cells with either
144 cereulide at 10 ng/ml or $^{13}\text{C}_6$ -cereulide.

145 2.5 Quantification of cereulide by LC-MSMS

146 The LC-MS/MS analysis was performed according to Rønning et al (2015) using an Agilent
147 1200 SL LC-system (Agilent Technologies, Waldbronn, Germany) coupled with an Agilent
148 G6460 MS/MS (Agilent Technologies, Santa Clara, CA, USA) (Ronning et al., 2015). The
149 limit of detection (LOD) and the limit of quantification (LOQ) were calculated as 3 times and
150 10 times the standard deviation of the signal obtained with the negative control, respectively
151 (Ronning et al., 2015). They were determined as 0.1 ng/ml (LOD) and 0.33 ng/ml (LOQ).

152 2.6 Expression of results

153 The amount of cereulide was expressed as ng per ml of the bacterial suspension washed off
154 the agar plate, with the whole biomass from one agar plate (19.6 cm^2) corresponding to 2 ml.

155 2.7 Statistical analysis

156 The results were expressed as the means of three independent biological replicates. A
157 Student's T-test was used to compare mean values. The null hypothesis was rejected for
158 $p < 0.05$

159

160 3 Results

161 *B. weihenstephanensis* BtB2-4 and MC67 strains were able to grow under all tested
162 conditions (Fig. 1 and 2), except at 8 °C with a pH 5.6 (data not shown). In artificially
163 inoculated food samples or media, cereulide production usually started at the end of the

164 exponential phase (Dommel et al., 2011; Thorsen et al., 2009a). For this reason, extraction of
165 cereulide was not done during the earlier phase of cell growth, as in (Rossvoll et al., 2014).

166 3.1 Influence of growth temperature on cereulide production

167 *B. weihenstephanensis* BtB2-4 and MC67 strains grew between 8 °C and 25 °C on PCA
168 plates at pH 7.0. At 8°C the stationary growth phase was reached after 7 days and 9-10 days
169 for strain BtB2-4 and strain MC67, respectively, with a maximal population (N_{max}) of
170 $9.42 \pm 0.12 \log_{10}$ CFU/ml for strain BtB2-4 and $8.50 \pm 0.02 \log_{10}$ CFU/ml for strain MC67
171 (Fig 1). The time to reach the stationary phase decreased with the increase of incubation
172 temperature (Fig 1).

173 The MC67 strain produced cereulide at 8 °C just above the detection limit (0.1 ng/ml) at a
174 level not quantifiable. It clearly produced emetic toxin between 10° and 25 °C. The BtB2-4
175 strain produced cereulide beginning at 8 °C and at all tested temperatures. For both strains,
176 the production of cereulide increased with increasing temperature (up to 25°C). Furthermore,
177 BtB2-4 cereulide production was always higher than that of MC67 (Fig 1). At 8 °C, the
178 maximal cereulide production (2.17 ± 0.69 ng/ml) by BtB2-4 was reached at the beginning of
179 stationary growth phase (7 days), while cereulide production was near the detection limit
180 (0.16 ± 0.03 ng/ml) for MC67 over the 11 days of the experiment (Fig. 1). At 10°C growth and
181 cereulide production of strains BtB2-4 and MC67 were also measured on TSA, in addition to
182 PCA. TSA differs from PCA by higher peptone content, presence of NaCl and absence of
183 glucose, compounds known to influence cereulide production (Dommel et al., 2010; Dommel
184 et al., 2011). Growth at 10 °C and pH 7.0 for these two *B. weihenstephanensis* strains on PCA
185 and TSA was similar, with an entry into stationary phase after 7 days (Fig. 1). The maximal
186 population of BtB2-4 cells on PCA plates and TSA plates were not significantly different
187 ($p > 0.05$) at 9.32 ± 0.02 and $9.95 \pm 0.14 \log_{10}$ CFU/ml, respectively. However, the maximal
188 population for strain MC67 was $9.23 \pm 0.09 \log_{10}$ CFU/ml on PCA plates and

189 9.81±0.01 log₁₀ CFU/ml on TSA plates, which was statistically significant ($p < 0.05$) (Fig. 1
190 and 2). The maximal cereulide production of the two strains on TSA plates at 10 °C, 6.83±1.3
191 ng/ml for BtB2-4 and 4.30±4.18 ng/ml for MC67, was not significantly different ($p > 0.05$)
192 (Fig 1). While the cereulide production of strain BtB2-4 (9.01±2.91 ng/ml) was similar on
193 TSA and on PCA plates ($p > 0.05$), the production of strain MC67 was increased on TSA
194 plates (4.30±4.18 ng/ml) compared with PCA plates at 7 days (0.85±0.40 ng/ml). However,
195 this difference was not significant ($p > 0.05$) and the cereulide production of that strain was
196 similarly low on both TSA and PCA plates for other incubation times at 10 °C (< 0.6 ng/ml)
197 (Fig 1). Therefore, the two strains produced the same amount of cereulide at 10°C on both
198 TSA and PCA.

199 At 15 °C, the maximal cereulide concentration produced by strain BtB2-4 was not higher than
200 that at 10 °C on PCA plates (Fig 1). The MC67 strain produced between 0.74±0.23 ng/ml and
201 2.45±1.46 ng/ml of cereulide, with a maximal concentration obtained after 7 days (Fig 1). At
202 25 °C, the production of cereulide on PCA plates at pH 7.0 was high for the BtB2-4 strain,
203 with the maximal cereulide concentration ($> 1,000$ ng/ml) reached at the end of the stationary
204 growth phase (Fig.1). The same kinetic of cereulide production was observed for the MC67
205 strain, with however a lower maximal cereulide concentration, not exceeding
206 13.20±4.67 ng/ml after 4 days (Fig 1). Cereulide production increased during the stationary
207 phase at 25 °C, whereas it was maximal at the onset of the stationary phase for the lower
208 temperatures tested.

209 3.2 Influence of pH on cereulide production

210 The growth of strains MC67 and BtB2-4 on PCA plates at pH 6.0 at 25 °C (Fig 2) was similar
211 to that observed on PCA plates at pH 7.0 at 25 °C (Fig. 1). The growth of both strains on PCA
212 plates at pH 5.4 was slower than that observed at pH 6.0 or 7.0 (Fig 1 and 2). Cereulide
213 production increased during the stationary growth phase for both strains at pH 6.0 (Fig 2), as

214 was observed for pH 7.0 (Fig 1). At these pH levels, BtB2-4 cereulide production was greater
215 than MC67 cereulide production, as observed at pH 7.0 ($p < 0.05$). The maximal cereulide
216 production of strain BtB2-4 at pH 6.0 and 25 °C was approximately 400 ng/ml, lower than the
217 production at pH 7.0 (Fig 1 and 2), while the maximal cereulide production of the MC67
218 strain at pH 6.0 (11 ± 0.50 ng/ml) was not significantly different ($p > 0.05$) from that observed
219 at pH 7.0 (13.20 ± 4.67 ng/ml) (Fig 1 and 2). When the pH of PCA plates was reduced to 5.4,
220 the maximal cereulide production of the BtB2-4 strain was 45.23 ± 29.56 ng/ml, with
221 5.53 ± 1.11 ng/ml of cereulide production for MC67 (Fig 2). The maximal production occurred
222 after 3 days and declined thereafter. However, cereulide production by BtB2-4 at pH 5.4 and
223 25 °C was still significantly higher than that of MC67 at pH 7.0 and 25 °C ($p < 0.05$) (Fig 1
224 and 2).

225

226 4 Discussion

227 In our study, the growth phase corresponding to the maximal cereulide production varied
228 with the growth temperature and pH. At 25 °C, cereulide production increased during the
229 stationary growth phase as previously observed for various strains of *B. cereus* at 25-30°C
230 (Dommel et al., 2011; Thorsen et al., 2009a). For *B. weihenstephanensis* BtB2-4, between
231 8 °C and 15 °C, maximum cereulide production occurred at the beginning of the stationary
232 phase and then decreased. At these low temperatures, it is possible that the expression of the
233 cereulide synthase genes (*ces*) is initially induced at the end of the exponential phase and
234 thereafter repressed during the stationary phase. The expression of the *ces* genes is activated
235 by Spo0A and repressed by CodY (Ehling-Schulz et al., 2015). Activation by Spo0A occurs
236 transiently at an early stage of its phosphorylation (Ehling-Schulz et al., 2015) which could
237 explain an initial production of cereulide at the end of the exponential phase. Then, it is
238 possible that at low temperatures repression by CodY was not lifted, explaining the

239 interruption of cereulide production during the stationary phase, in contrast to the
240 accumulation observed at higher temperatures. Indeed, cold has been shown to increase
241 production of CodY in *Listeria monocytogenes* (Durack et al., 2013), but the effect of cold on
242 CodY in *B. weihenstephanensis* is not known. Alternatively, temperatures of 8-10 °C might
243 impact on cereulide production by *B. weihenstephanensis* at the post transcriptional level, as
244 proposed for mesophilic strains of *B. cereus* (Kranzler et al., 2016). In such case, after an
245 initial production at the end of the exponential phase, it is possible that the cereulide synthase
246 loses its activity, due to e.g. a lower stability and a poor translation at low temperatures. At pH
247 5.4, cereulide production occurred during the exponential growth phase. However, the impact
248 of pH on the expression of the *ces* gene cluster, on its effectors, and on the activity of the
249 cereulide synthase, is unknown. Initiation of cereulide production during the exponential
250 growth phase was previously shown by Ronning et al. (2015), in food samples naturally
251 contaminated with a very low number of *B. cereus*.

252 When MC67 grew at 25 °C, cereulide production did not exceed 13 ng/ml in the extract,
253 corresponding to 22 ng/plate, which was low compared to previous results showing that
254 MC67 produced up to 14 µg/cm² after 24 h at 25 °C (Thorsen et al., 2009a). This may be due
255 to differences in the protocols used, such as preparation of the inoculum and/or growth media
256 (Rossvoll et al., 2014), as previously suggested by (Thorsen et al., 2009a) for *B.*
257 *weihenstephanensis* and (Dommel et al., 2010) for mesophilic *B. cereus*. In particular, after a
258 pre-incubation of the inoculated PCA plates at 8 °C for 2 or 3 weeks, the MC67 cereulide
259 production at 25 °C was approximately 10,000 times lower (0.001 µg/cm²) than without pre-
260 incubation (Thorsen et al., 2009a). In the present study, the pre-incubation at 4 °C during the
261 18-21 h before growth experiments could explain the low cereulide production. The
262 expression of the *ces* genes is activated by Spo0A and repressed by CodY, which linked
263 cereulide production to the early stage of sporulation and to the nutritional and energy status

264 of the cells, respectively (Ehling-Schulz et al., 2015). Although cold stress has been shown to
265 increase production of CodY in *Listeria monocytogenes* (Durack et al., 2013), the potential
266 influence of pre-incubation at low temperatures on *B. weihenstephanensis* CodY production
267 remains unknown. Nevertheless, this possible effect of pre-incubation is strain dependent,
268 because in the same conditions, *B. weihenstephanensis* BtB2-4 produced much higher levels
269 of cereulide than MC67.

270 In MC67, the cereulide gene cluster is located on a plasmid and is part of a composite
271 transposon (Mei et al., 2014). Whether this peculiar genetic location of *ces* genes in MC67
272 could explain important variations in their expression is not known.

273 Environmental conditions of growth influenced the amount of cereulide produced by
274 *B. weihenstephanensis*. Lowering the medium pH reduced cereulide production, particularly
275 at pH 5.4, while *B. weihenstephanensis* reached a maximal population of approximately
276 10^9 CFU/plate as at pH 7.0. Cereulide production at various pH levels has been previously
277 reported for mesophilic *B. cereus* strains in food matrices (Dommel et al., 2010). For instance,
278 cereulide production by *B. cereus* F4810/72 was higher in béarnaise sauce at pH 5.8
279 (8.0 $\mu\text{g/g}$) than in cooked rice at pH 7.0 (1.8 $\mu\text{g/g}$) or in camembert cheese at pH 7.9
280 (0.6 $\mu\text{g/g}$). However, one could hypothesize that the food matrix may also have an influence
281 on cereulide production (Ehling-Schulz et al., 2015).

282 In our study, lowering the growth temperature markedly reduced cereulide production for
283 both *B. weihenstephanensis* strains. This is particularly clear for strain BtB2-4, which
284 produced 1,000 ng/ml of cereulide at 25 °C, 10 ng/ml at 15 °C or 10 °C, and 2 ng/ml at 8 °C.
285 Less cereulide was produced by a mesophilic *B. cereus* on media rich in proteinaceous
286 ingredients such as TSA (Dommel et al., 2010) and NaCl, which is absent in PCA, inhibited
287 cereulide production (Dommel et al., 2011). The same low production of cereulide at low

288 temperatures by *B. weihenstephanensis* was measured at 10 °C on the two media (TSA and
289 PCA), indicating that the effect of cold presumably prevails over the effect of growth media.
290 Storing foods at 4 °C prevents the growth of *B. weihenstephanensis* (Guinebretière et al.,
291 2008). Yet, at low temperatures allowing the growth of *B. weihenstephanensis*, it is important
292 to assess if the amount of cereulide produced can cause food poisoning. It has been shown
293 that 10 to 1,280 ng/g of cereulide were present in 13 samples from foods identified as being
294 the source of vomiting-type food poisoning due to cereulide produced by *B. cereus* (Agata et
295 al., 2002). Based on these observations and the results reported in the present study, the risk
296 of an emetic poisoning from food stored below 8 °C seems very low. In contrast, the amount
297 of cereulide produced by *B. weihenstephanensis* BtB2-4 at 10 °C-15 °C and 25 °C and pH
298 7.0, or at 25 °C and pH 5.7, could represent a risk for consumers. However, the low
299 prevalence of cereulide producing *B. weihenstephanensis* must be taken into account. Only
300 three of these strains were found among approximately 2000 strains from *B. cereus sensu lato*
301 (Hoton et al., 2009) and no cereulide producers were found among 143 strains of *B.*
302 *weihenstephanensis* (Guinebretière et al., 2008). This may explain why so far no *B.*
303 *weihenstephanensis* has been linked to emetic foodborne poisoning. However, more strains of
304 *B. weihenstephanensis* might acquire the cereulide genes by horizontal transfer in the future
305 (Mei et al., 2014).

306 The present study and previous works showed that the environmental conditions of growth
307 (temperature, pH, oxygen, NaCl concentration), the inoculum history (temperature,
308 sporulation), the food composition (proteins versus carbohydrates) and the strain strongly
309 influence cereulide production by mesophilic *B. cereus* or psychrotolerant
310 *B. weihenstephanensis*. Therefore, it is important to consider this information to assess
311 cereulide production in food products.

312

313 **Acknowledgements**

314 This study is part of the OPTIFEL project that received funding from the European Union's

315 Seventh Framework Program for research, technological development and demonstration

316 under grant agreement n° 311754. AG's PhD grant is also from the EU Optifel project.

317

ACCEPTED MANUSCRIPT

318 **References**

319

320 Agata, N., Ohta, M., Mori, M., Isobe, M., 1995. A novel dodecadepsipeptide, cereulide, is an
321 emetic toxin of *Bacillus cereus*. FEMS Microbiol. Lett. 129, 17-19.

322 Agata, N., Ohta, M., Yokoyama, K., 2002. Production of *Bacillus cereus* emetic toxin
323 (cereulide) in various foods. Int. J. Food Microbiol. 73, 23-27.

324 Biesta-Peters, E.G., Reij, M.W., Blaauw, R.H., in't Veld, P.H., Rajkovic, A., Ehling-Schulz,
325 M., Abee, T., 2010. Quantification of the Emetic Toxin Cereulide in Food Products by Liquid
326 Chromatography-Mass Spectrometry Using Synthetic Cereulide as a Standard. Appl. Environ.
327 Microbiol. 76, 7466-7472.

328 Carlin, F., Fricker, M., Pielaat, A., Heisterkamp, S., Shaheen, R., Salkinoja Salonen, M.,
329 Svensson, B., Nguyen-the, C., Ehling-Schulz, M., 2006. Emetic toxin-producing strains of
330 *Bacillus cereus* show distinct characteristics within the *Bacillus cereus* group. Int. J. Food
331 Microbiol. 109, 132-138.

332 Dierick, K., Van Coillie, E., Swiecicka, I., Meyfroidt, G., Devlieger, H., Meulemans, A.,
333 Hoedemaekers, G., Fourie, L., Heyndrickx, M., Mahillon, J., 2005. Fatal family outbreak of
334 *Bacillus cereus*-associated food poisoning. J. Clin. Microbiol. 43, 4277 - 4279.

335 Dommel, M.K., Frenzel, E., Strasser, B., Blochinger, C., Scherer, S., Ehling-Schulz, M.,
336 2010. Identification of the Main Promoter Directing Cereulide Biosynthesis in Emetic
337 *Bacillus cereus* and Its Application for Real-Time Monitoring of *ces* Gene Expression in
338 Foods. Appl. Environ. Microbiol. 76, 1232-1240.

339 Dommel, M.K., Lucking, G., Scherer, S., Ehling-Schulz, M., 2011. Transcriptional kinetic
340 analyses of cereulide synthetase genes with respect to growth, sporulation and emetic toxin
341 production in *Bacillus cereus*. Food Microbiol. 28, 284-290.

342 Durack, J., Ross, T., Bowman, J.P., 2013. Characterisation of the Transcriptomes of
343 Genetically Diverse *Listeria monocytogenes* Exposed to Hyperosmotic and Low Temperature
344 Conditions Reveal Global Stress-Adaptation Mechanisms. PLoS One 8.

345 Ehling-Schulz, M., Frenzel, E., Gohar, M., 2015. Food-bacteria interplay: pathometabolism of
346 emetic *Bacillus cereus*. Front. Microbiol. 6:704.

- 347 Ehling-Schulz, M., Fricker, M., Grallert, H., Rieck, P., Wagner, M., Scherer, S., 2006a.
348 Cereulide synthetase gene cluster from emetic *Bacillus cereus*: Structure and location on a
349 mega virulence plasmid related to *Bacillus anthracis* toxin plasmid pXOI. BMC Microbiol.
350 6:20.
- 351 Ehling-Schulz, M., Fricker, M., Scherer, S., 2004. *Bacillus cereus*, the causative agent of an
352 emetic type of food-borne illness. Mol. Nutr. Food Res. 48, 479-487.
- 353 Ehling-Schulz, M., Vukov, N., Schulz, A., Shaheen, R., Andersson, M., Martlbauer, E.,
354 Scherer, S., 2005. Identification and partial characterization of the nonribosomal peptide
355 synthetase gene responsible for cereulide production in emetic *Bacillus cereus*. Appl.
356 Environ. Microbiol. 71, 105-113.
- 357 Guérin, A., Dargaignaratz, C., Broussolle, V., Clavel, T., Nguyen-the, C., 2016. Combined
358 effect of anaerobiosis, low pH and cold temperatures on the growth capacities of
359 psychrotrophic *Bacillus cereus*. Food Microbiol. 59, 119-123.
- 360 Guinebretière, M.H., Thompson, F.L., Sorokin, A., Normand, P., Dawyndt, P., Ehling-Schulz,
361 M., Svensson, B., Sanchis, V., Nguyen-The, C., Heyndrickx, M., De Vos, P., 2008.
362 Ecological diversification in the *Bacillus cereus* Group. Env. Microbiol. 10, 851-865.
- 363 Guinebretiere, M.H., Velge, P., Couvert, O., Carlin, F., Debuyser, M.L., Nguyen-The, C.,
364 2010. Ability of *Bacillus cereus* group strains to cause food poisoning varies according to
365 phylogenetic affiliation (groups I to VII) rather than species affiliation. J. Clin. Microbiol. 48,
366 3388-3391.
- 367 Hoton, F., Andrup, L., Swiecicka, I., Mahillon, J., 2005. The cereulide genetic determinants
368 of emetic *Bacillus cereus* are plasmid-borne. Microbiology 151, 2121-2124.
- 369 Hoton, F., Fornelos, N., N'Guessan, E., Hu, X., Swiecicka, I., Dierick, K., Jaaskelainen, E.,
370 Salkinoja-Salonen, M., Mahillon, J., 2009. Family portrait of *Bacillus cereus* and *Bacillus*
371 *weihenstephanensis* cereulide-producing strains. Environ. Microbiol. Rep. 1, 177-183.
- 372 Kranzler, M., Stollewerk, K., Rouzeau-Szynalski, K., Blayo, L., Sulyok, M., Ehling-Schulz,
373 M., 2016. Temperature Exerts Control of *Bacillus cereus* Emetic Toxin Production on Post-
374 transcriptional Levels. Front. Microbiol. 7.
- 375 Mahler, H., Pasi, A., Kramer, J., Schulte, P., Scoging, A., Bar, W., Krahenbuhl, S., 1997.
376 Fulminant liver failure in association with the emetic toxin of *Bacillus cereus*. N. Engl. J.
377 Med. 336, 1142-1148.

- 378 Mei, X., Xu, K., Yang, L., Yuan, Z., Mahillon, J., Hu, X., 2014. The genetic diversity of
379 cereulide biosynthesis gene cluster indicates a composite transposon Tnces in emetic *Bacillus*
380 *weihenstephanensis*. BMC Microbiol. 14, 149.
- 381 Mikkola, R., Saris, N., Grigoriev, P., Andersson, M., Salkinoja-Salonen, M., 1999.
382 Ionophoretic properties and mitochondrial effects of cereulide - the emetic toxin of *B. cereus*.
383 Eur. J. Biochem. 263, 112-117.
- 384 Naranjo, M., Denayer, S., Botteldoorn, N., Delbrassinne, L., Veys, J., Waegenaere, J.,
385 Sirtaine, N., Driesen, R., Sipido, K., Mahillon, J., Dierick, K., 2011. Sudden death of a young
386 adult associated with *Bacillus cereus* food poisoning. J. Clin. Microbiol. 49, 4379-4381.
- 387 Paananen, A., Mikkola, R., Sareneva, T., Matikainen, S., Hess, M., Andersson, M., Julkunen,
388 I., Salkinoja-Salonen, M., Timonen, T., 2002. Inhibition of human natural killer cell activity
389 by cereulide, an emetic toxin from *Bacillus cereus*. Clin. Exp. Immunol. 129, 420-428.
- 390 Ronning, H.T., Asp, T.N., Granum, P.E., 2015. Determination and quantification of the
391 emetic toxin cereulide from *Bacillus cereus* in pasta, rice and cream with liquid
392 chromatography-tandem mass spectrometry. Food Add. Contam.: Part A 32, 911-921.
- 393 Rossvoll, E., Ronning, H.T., Granum, P.E., Moretro, T., Hjerpekkjon, M.R., Langsrud, S.,
394 2014. Toxin production and growth of pathogens subjected to temperature fluctuations
395 simulating consumer handling of cold cuts. Int. J. Food Microbiol. 185, 82-92.
- 396 Sorokin, A., Candelon, B., Guilloux, K., Galleron, N., Wackerow-Kouzova, N., Ehrlich, S.D.,
397 Bourguet, D., Sanchis, V., 2006. Multiple-locus sequence typing analysis of *Bacillus cereus*
398 and *Bacillus thuringiensis* reveals separate clustering and a distinct population structure of
399 psychrotrophic strains. Appl. Env. Microbiol. 72, 1569-1578.
- 400 Stenfors Arnesen, L.P., Fagerlund, A., Granum, P.E., 2008. From soil to gut: *Bacillus cereus*
401 and its food poisoning toxins. Fems Microbiol. Rev. 32, 579-606.
- 402 Teplova, V.V., Mikkola, R., Tonshin, A.A., Saris, N.E.L., Salkinoja-Salonen, M.S., 2006. The
403 higher toxicity of cereulide relative to valinomycin is due to its higher affinity for potassium
404 at physiological plasma concentration. Toxicol. Appl. Pharm. 210, 39-46.
- 405 Thorsen, L., Budde, B.B., Henrichsen, L., Martinussen, T., Jakobsen, M., 2009a. Cereulide
406 formation by *Bacillus weihenstephanensis* and mesophilic emetic *Bacillus cereus* at
407 temperature abuse depends on pre-incubation conditions. Int. J. Food Microbiol. 134, 133-
408 139.

- 409 Thorsen, L., Budde, B.B., Koch, A.G., Klingberg, T.D., 2009b. Effect of modified
410 atmosphere and temperature abuse on the growth from spores and cereulide production of
411 *Bacillus weihenstephanensis* in a cooked chilled meat sausage. *Int. J. Food Microbiol.* 130,
412 172-178.
- 413 Thorsen, L., Hansen, B.M., Nielsen, K.F., Hendriksen, N.B., Phipps, R.K., Budde, B.B.,
414 2006. Characterization of emetic *Bacillus weihenstephanensis*, a new cereulide-producing
415 bacterium. *Appl. Environ. Microbiol.* 72, 5118-5121.
- 416 Vilas-Boas, G., Sanchis, V., Lereclus, D., Lemos, M.V.F., Bourguet, D., 2002. Genetic
417 differentiation between sympatric populations of *Bacillus cereus* and *Bacillus thuringiensis*.
418 *Appl. Env. Microbiol.* 68, 1414-1424.
- 419
- 420
- 421

422 **Figure Legends**

423 **Figure 1.** Growth curves of *B. weihenstephanensis* BtB2-4 and MC67 (log CFU/plate) (◆)
424 and cereulide (crl) production (ng/ml) (○) at 8 °C, 10 °C, 15 °C and 25 °C on PCA plates and
425 on TSA plates at pH 7.0. Error bars indicate standard deviations of results obtained with
426 independent triplicates and can be smaller than symbols.

427


428

429 **Figure 2.** Growth curves of *B. weihenstephanensis* BtB2-4 and MC67 (log CFU/plate) (◆)
430 and cereulide (crl) production (ng/ml) (○) on PCA plates at pH 6.0 or at pH 5.4 at 25 °C.
431 Error bars indicate standard deviations of results obtained with independent triplicates and can
432 be smaller than symbols.


433

434


Guérin et al. R1 Figure 1



Guérin et al. R1 Figure 1



Guérin et al. R1 Figure 2



Highlights

- *B. weihenstephanensis* produced quantifiable cereulide at 8°C and pH7
- *B. weihenstephanensis* produced cereulide at pH 5.4 and 25°C
- Cereulide production increased 5-fold between 8°C and 10°C
- Amount of cereulide produced was markedly different between the strains tested