



**HAL**  
open science

## Diet-induced hyperglycaemia alters the hepatic microRNA profile in rainbow trout, *O. mykiss*

Jan A. Mennigen, Stéphane Panserat, Lucie Marandel

### ► To cite this version:

Jan A. Mennigen, Stéphane Panserat, Lucie Marandel. Diet-induced hyperglycaemia alters the hepatic microRNA profile in rainbow trout, *O. mykiss*. PWH Symposium (satellite symposium preceding the 56th annual meeting of the Canadian Society of Zoologists), May 2017, Winnipeg, Canada. , pp.27, 2017, PWH Symposium schedule and abstracts. hal-01602644

**HAL Id: hal-01602644**

**<https://hal.science/hal-01602644>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

### 3.3 Diet-induced hyperglycemia alters the hepatic microRNA profile in rainbow trout, *O. mykiss*.

Jan A. Mennigen<sup>1</sup>, Stephane Panserat<sup>2</sup> and Lucie Marandel<sup>2</sup>

<sup>1</sup>*Department of Biology, University of Ottawa, K1N6A5, Ottawa, Ontario, Canada.*

<sup>2</sup>*Institut National de la Recherche Agronomique, Aquapôle, 64310 Saint-Pée-sur-Nivelle, Aquitaine, France*

MicroRNAs are small, non-protein coding RNAs, which have emerged as potent regulators of glucose metabolism in mammalian model species through destabilization of specific mRNAs and inhibition of protein translation. While many miRNAs are highly conserved in evolution, teleost fish possess additional lineage- and species-specific miRNAs, targeting a potentially vastly different set of mRNA targets. To gain insight into the role of miRNA networks in the 'glucose-intolerant' phenotype in rainbow trout, we profiled hepatic miRNA abundance in the liver of rainbow trout exhibiting diet-induced normo- and hyperglycemia. Our study identified a total of 26 differentially regulated miRNAs, among them several previously identified as being regulated in a hyperglycemic environment in mammalian models. While this suggests at least some regulatory similarities between miRNA regulation in hyperglycemic rainbow trout and mammalian models, predicted mRNA targets suggest a large degree of divergence. Using our identified miRNA targets, future studies will focus on possible causative roles for select hepatic miRNAs towards the glucose-intolerant phenotype in rainbow trout.