

Temporal differentiation of soil biota communities and of trophic networks in response to arable crop management strategies

Valérie Coudrain, Mickael Hedde, Matthieu Chauvat, Pierre-Alain Maron, Emilie Bourgeois, Bruno Mary, Joël Léonard, Flemming Ekelund, Cécile Villenave, Elisa Thébault, et al.

▶ To cite this version:

Valérie Coudrain, Mickael Hedde, Matthieu Chauvat, Pierre-Alain Maron, Emilie Bourgeois, et al.. Temporal differentiation of soil biota communities and of trophic networks in response to arable crop management strategies. Sfécologie-2016, International Conference of Ecological Sciences, Société Française d'Ecologie (SFE). FRA., Oct 2016, Marseille, France. hal-01602967

HAL Id: hal-01602967

https://hal.science/hal-01602967

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Temporal differentiation of soil biota communities and of trophic networks in response to arable crop management strategies

Valérie Coudrain*^{1,2,3}, Mickael Hedde^{†2}, Matthieu Chauvat³, Pierre-Alain Maron⁴, Emilie Bourgeois⁴, Bruno Mary⁵, Joel Léonard⁵, Flemming Ekelund⁶, Cécile Villenave⁷, Elisa Thébault⁸, and Sylvie Recous¹

 $^1\mathrm{UMR614}$ FARE – Institut national de la recherche agronomique (INRA) – F-51100 Reims, France, France

²UMR1402 ECOSYS – Institut national de la recherche agronomique (INRA) – F-78850 Thiverval-Grignon, France, France

³Ecodiv – Université de Rouen : EA1293 – F-76821 Mont Saint Aignan, France, France
⁴UMR1347 Agroécologie – Institut national de la recherche agronomique (INRA) – F-21034 Dijon, France, France

⁵UR1158 AgroImpact – Institut national de la recherche agronomique (INRA) – F-02000 Barenton-Bugny, France, France

⁶Department of Biology – University of Copenhagen, DK-2100 Copenhagen, Denmark, Denmark
⁷ELISOL environnement – Entreprise privée – F-30111 Congénies, France, France
⁸IEES – CNRS : UMR7618 – UPMC - iEES Paris - Bât. A, 7e étage 7, quai St Bernard 75252 PARIS cedex 05, France

Abstract

Promoting diverse and functional biological communities is an important objective of agroecology, with increasing attention given to the important role of soil biodiversity. In an experimental study conducted in actual field conditions, we followed over four years the dynamic of soil organisms from microorganisms to micro-, meso- and macro-fauna in a cropping system submitted to four different treatments that varied in tillage, residue management and N fertilization rate. Differentiation in taxonomic diversity, in the abundance and diversity of the multiple trophic groups and in food-web structure was evaluated. A co-statis analysis was used to investigate the joint dynamics of food-web structure and soil functions linked to C and N dynamics.

Our study demonstrated a delayed but effective differentiation in soil biota diversity following implementation of the agricultural practices. Soil biodiversity varied throughout time with some groups responding more readily than others. While tillage appeared a main factor of influence, surprisingly little impact of residue management and nitrogen fertilization could be observed. Significant interactive effect of year and treatment was found on the mean trophic level and the bacterial to fungal path ratio, but not on the number of trophic groups and Shannon's link diversity. Finally, co-statis analysis revealed significant differences in

^{*}Corresponding author: vcoudrain@gmail.com

[†]Speaker

co-dynamics between food-web structure and soil functions over four years, mainly drove by conventionally tilled vs reduced tilled treatments.

Through its multi-taxonomic approach, the present study increases our understanding of the dynamic of soil communities in agricultural cropping systems and helps identify possible consequences for soil functioning.

Keywords: Trophic groups, Invertebrates, Microorganisms, Tillage, Plant residues, Fertilization, Functional traits