



The Ephrin receptor: a potential actor in polerovirus transmission by aphids

Michaël Mulot, Sylvaine Boissinot, Baptiste Monsion, Sophie Meyer, Nicole Bochet, Véronique Brault

► To cite this version:

Michaël Mulot, Sylvaine Boissinot, Baptiste Monsion, Sophie Meyer, Nicole Bochet, et al.. The Ephrin receptor: a potential actor in polerovirus transmission by aphids. 3. Hemipteran-Plant Interactions Symposium. HPIS 2017, Jun 2017, Madrid, Spain. 234 p., 2017, HPIS 2017 3rd Hemipteran-Plant Interactions Symposium: book of abstracts. hal-01603875

HAL Id: hal-01603875

<https://hal.science/hal-01603875>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Ephrin receptor: a potential actor in polerovirus transmission by aphids


Michaël Mulot, Sylvaine Boissinot, Baptiste Monsion, Sophie Meyer, Nicole Bochet, Véronique Brault
michael.mulot@inra.fr

Poleroviruses are phloem-limited RNA viruses strictly transmitted by aphids in a circulative and non-propagative mode. Virus particles, acquired by aphids on infected plants, successively cross intestinal and accessory salivary gland cells by transcytosis before being released, together with saliva, into a plant during aphid feeding. Virus transport through the epithelia relies on the presence of specific virus receptors. The Ephrin receptor (*Ephr*) from *Myzus persicae* has been identified as a potential partner of the structural proteins of the polerovirus *Turnip yellows virus* (TuYV). In order to address its function in TuYV transmission by *M. persicae*, we developed an RNA interference-based strategy to inhibit *Ephr* expression in *M. persicae*. When fed on a viral source, the *Ephr*-silenced aphids showed a lower internalization of viral genomes into their body and a reduced ability to transmit the virus. Taken together, these experiments strongly suggest implication of *Ephr* in TuYV acquisition by *M. persicae*.

Oral aquisition by aphids of *in vitro* synthesized dsRNA targeting *Ephr*


LacZ used as a control

dsRNA-*Ephr*
dsRNA-*LacZ*



In vitro synthesized dsRNA
(agarose gel)

dsRNA-*Ephr* or dsRNA-*LacZ*




5 days acquisition
by aphids

Ephr-mRNA accumulation in aphids after dsRNA-*Ephr* acquisition





Gut dissection before Q-RT-PCR
analysis



	Exp.1	Exp.2	Exp.3	Exp.4
Relative expression of <i>Ephr</i> (% inhibition)	-54%	-9%	+40%	-57%


Inhibition of *Ephr*-mRNA accumulation
after dsRNA acquisition is not
reproducibly observed



Ephr expression is unstable in
aphid's guts after *in vitro* feeding

May be responsible for the
lack of reproducible
silencing of *Ephr*?

Aphid fecondity is not affected by dsRNA-*Ephr* acquisition



Nb larvae daily produced by aphid
during 5 days after dsRNA acquisition

Aphid feeding is positively affected by dsRNA-*Ephr* acquisition


Artificial feeding medium

Ephr-silenced aphids


pH indicator paper




Honeydew collect during 24h
image analysis with ImageJ software




Acquisition of dsRNA-*Ephr* reduces virus internalization in aphids and virus transmission to plants



Virus transmission is decreased after acquisition of mamalian *Ephr* inhibitor*



*Max Tognolini, Dipartimento di scienze degli alimenti e del farmaco, Parma, Italy



Conclusions

A 5 day-acquisition of dsRNA-*Ephr* by aphids reproducibly induced a lower internalization of TuYV genome into aphid's body, although reduction of *Ephr*-mRNA accumulation could not be systematically confirmed. Aphids fed on dsRNA-*Ephr* reproducibly transmitted TuYV with a lower efficiency while their feeding behavior was not reduced and their fecondity not altered. Feeding aphids on *Ephr* inhibitor before virus acquisition also reduces virus transmission. Taken together, these experiments strongly suggest implication of *Ephr* in polerovirus transmission. Whether *Ephr* encodes a true virus receptor or a co-receptor still needs to be elucidated.