



**HAL**  
open science

## In search of a conservation limit for the Allier A. salmon population: how and what for ?

Etienne Prévost

### ► To cite this version:

Etienne Prévost. In search of a conservation limit for the Allier A. salmon population: how and what for?. Scientific Committee for Diadromous Fish of the Loire, Nov 2016, Orléans, France. hal-01603880

**HAL Id: hal-01603880**

**<https://hal.science/hal-01603880>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# In search of a Conservation Limit for the Allier A. Salmon population: How and what for ?

**Etienne Prévost**

*INRA - UMR Ecobiop*

*Saint-Pée-sur-Nivelle - France*






# NASCO Conservation Limit

- Conservation Limit is a Biological Reference Point
  - BRP: Abundance benchmark against which population status can be assessed
- Conservation Limit (CL): abundance level above which population should be maintained
  - If falling below, remedial management action should be taken
- NASCO CL: spawning escapement producing MSY
  - Derived from Stock-Recruitment analysis
- An illustration using the Nivelles R. data series

# The Nivelle R.: the southernmost Index River in Europe



Uxondoa trapping station



Redd counts




Olha trapping station


Adult returns estimation




# The egg-to-egg Stock/Recruitment series of the Nivelles R.




# The egg-to-egg Stock/Recruitment series of the Nivelles R.



# The egg-to-egg Stock/Recruitment series of the Nivelles R.




# The egg-to-egg Stock/Recruitment series of the Nivelles R.







# The egg-to-egg Stock/Recruitment series of the Nivelles R.




# The egg-to-egg Stock/Recruitment series of the Nivelles R.



# The egg-to-egg Stock/Recruitment series of the Nivelles R.




# The egg-to-egg Stock/Recruitment series of the Nivelles R.



# Troubles with NASCO CL definition


- Assume catches (at sea and in rivers) are well known or negligible
  - Generally not true
    - E.g. in the Nivelle: significant but unknown coastal exploitation
- Spawning stock producing MSY: is this relevant to conservation?
  - Rather aims at fisheries exploitation optimization
- What if survival (e.g. at sea) drops?

# Evolution of NASCO CL if sea survival drops



Recruitment curve flattens  
→ NACSO CL decreases

Sea survival drops by 25%



# CL definition: time for a change?


- NASCO CL: based on fisheries optimization principle
  - when times become harder for salmon CL should be revised downward
  - for populations below replacement (i.e. non viable; e.g. Allier): NASCO CL is not defined
- Fisheries optimization  $\neq$  Conservation
- But how can we define Conservation?
- An operational (and partial) proposal: conservation should ensure that wild juvenile recruitment does not fall too low
- Associated CL definition: spawning escapement threshold that ensure wild juvenile recruitment does not fall too low

# The Nivelles R.: the southernmost Index River in Europe

Uxondoa  
trapping  
station



Spawning



Juvenile  
(0+ parr in the fall)  
recruitment



Redd  
counts



escapement




Electrofishing

Olha  
trapping  
station







# The egg-to-0+ parr Stock/Recruitment series of the Nivelles R.



# The egg-to-0+ parr Stock/Recruitment series of the Nivelles R.



# The egg-to-0+ parr Stock/Recruitment series of the Nivelles R.




# CL definition: time for a change?

- Acknowledge recruitment stochasticity (uncertainty)
- Modified CL definition: spawning escapement threshold that ensure a low probability of wild juvenile recruitment falling too low
  - Conservation aims at controlling the risk of low recruitment
- But what is low recruitment?
  - A management decision
  - An operational proposal: set as a proportion of the carrying capacity ( $R_{max}$ )
- And what is low probability?
  - A management decision
- Scientific advice: produce (risk) plots that help management decision


# Risk plot for 3 options of low recruitment level

Risk:  $\text{Proba}(\text{Recruitment} < X\% \text{ of } R_{\text{max}})$



# Risk plot: an example of management decision


Risk:  $\text{Proba}(\text{Recruitment} < X\% \text{ of } R_{\text{max}})$



***New CL definition adopted by DFO for A. Salmon stocks in Canada (apart from Québec)***

# Risk plot: being too risk averse is not realistic

Risk:  $\text{Proba}(\text{Recruitment} < X\% \text{ of } R_{\text{max}})$



# New CL approach: pros & cons

- PROs

- Based on a conservation principle/definition in a risk control context
- Identify the risk of low recruitment level as a management decision
  - Easier to communicate to management stakeholders (?)
- Uses scientific and fisheries independent data only
- Not sensitive to changes in density independent mortality (e.g. at sea)

- CONs

- CL estimation sensitive to recruitment rate at low spawning escapement
  - A quantity usually difficult to estimate from SR data
- Do not take into account potential density dependence of survival after the juvenile recruitment stage considered





# CL for the Allier A. Salmon population ?

- NASCO CL: not an option
  - Population currently below replacement
- Definition of a new CL based based on a low risk of low recruitment: can be done
  - The Allier salmon population model (INRA – LOGRAMI) includes a density dependent SR sub-model

# A CL for the Allier A. Salmon population ?

wild 0+ density dependence relationship



# CL for the Allier A. Salmon population ?


- NASCO CL: not an option
  - Population currently below replacement
- Definition of a new CL based based on a low risk of low recruitment: can be done
  - The Allier salmon population model (INRA – LOGRAMI) includes a density dependent SR sub-model
  - Plots of risk of low recruitment vs number of spawners reaching Vichy can be produced
  - Management (stakeholders) should decide on the definition of low recruitment and low risk (probability)

# CL for the Allier A. Salmon population: so what?

- CL is a useful benchmark for assessing conservation status of the population
  - **Caution:** CL is purely demographic → makes no consideration of genetics
  - CL does not “generate” management measures: it helps assessing consequences of management measures
  - Given the model available for the Allier salmon, a CL is not a necessity: we can directly assess conservation achievement (provided we agree on an operational definition of conservation)
- Retrospective (up to present) assessment:
  - What is current conservation status of the population?
  - Have past management measures improve conservation?
- Prospective assessment:
  - Predicted evolution of conservation according to various management options (habitat improvement and/or stocking)
  - Beyond viability → identification of exploitable surplus (if any)


# A (fictitious) illustration of the use of CL for the Allier A. Salmon population ?

Habitat: status quo  
Stocking: none



Conservation not achieved

Habitat: major improvement  
Stocking: none



Conservation achieved  
& surplus available for exploitation

***Thanks for your attention***

