



**HAL**  
open science

## **Sociodemographic and economic factors are essential determinants of weight gain between before and after cancer diagnosis: Results from the prospective population-based NutriNet-Sante cohort**

Philippine Fassier, Laurent Zelek, Valentin Partula, Lucie Lecuyer, Bernard Srour, P. Bachmann, M. Touillaud, Nathalie Pecollo, Maria del Pilar Galan Hercberg, Serge Hercberg, et al.

### ► To cite this version:

Philippine Fassier, Laurent Zelek, Valentin Partula, Lucie Lecuyer, Bernard Srour, et al.. Sociodemographic and economic factors are essential determinants of weight gain between before and after cancer diagnosis: Results from the prospective population-based NutriNet-Sante cohort. San Antonio Breast Cancer Symposium, Dec 2016, San Antonio, United States. pp.2, 10.1158/1538-7445.SABCS16-P5-13-01 . hal-01603941

**HAL Id: hal-01603941**

**<https://hal.science/hal-01603941>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 2016 San Antonio Breast Cancer Symposium

**Publication Number:** P5-13-01

**Title:** Sociodemographic and economic factors are essential determinants of weight gain between before and after cancer diagnosis: Results from the prospective population-based NutriNet-Santé cohort

Fassier P, Zelek L, Partula V, Lecuyer L, Srour B, Bachmann P, Touillaud M, Druetne-Pecollo N, Galan P, Hercberg S, Cohen P, Hoarau H, Latino-Martel P, Gonzalez R, Deschasaux M and Touvier M. Nutritional Epidemiology Research Team (EREN): Inserm U1153, Inra U1125, Cnam, Paris 5, 7 and 13 Universities, Sorbonne Paris Cité Epidemiology and Statistics Research Center, Bobigny, France; French Network for Nutrition and Cancer Research (NACRe Network), France; Avicenne Hospital, APHP, Bobigny, France; Cancer, Environment and Nutrition Unit, Léon Bérard Center, Lyon, France; Avicenne Hospital, APHP, Bobigny, France and University of Rouen, DySola, EA 4701, Rouen, France.

**Body: Background:** While many cancer patients are affected by weight loss, others tend to gain weight, which may impact prognosis and risk of recurrence and of second cancer. The aim of this prospective study was to investigate weight variation between before and after cancer diagnosis and socio-demographic, economic, lifestyle and clinical factors associated with moderate-to-severe weight gain.

**Methods:** 1051 incident cases of first primary cancer were diagnosed in the NutriNet-Santé cohort between 2009 and 2015. Weight was prospectively collected every 6 months since subjects' inclusion (i.e. an average of 2y before diagnosis). Mean weights before and after cancer diagnosis were compared with paired Student's t-test. Factors associated with moderate-to-severe weight gain ( $\geq 5\%$  of initial weight) were investigated by multivariable logistic regression.

**Results:** Weight loss was observed in men (-3.54kg in those who lost weight,  $p=0.0002$ ) and in colorectal cancer patients (-3.94kg,  $p=0.0012$ ). Weight gain was observed in breast and skin cancers (2.83kg,  $p=0.047$ , and 2.96kg,  $p=0.03$  respectively). Women (OR=1.99[1.18-3.35]), younger patients (OR=1.78[1.05-3.03]), those with lower education (OR=2.17[1.07-4.37]), those with excess weight before diagnosis (OR=1.53[1.02-2.30]) and those who stopped smoking after diagnosis (OR=4.60[2.06-10.25]) were more likely to experience moderate-to-severe weight gain. In breast cancer patients, induced menopause was associated with weight gain (OR=4.12[1.76-9.67]), but no association was detected for tumor characteristics or treatments.

**Conclusion:** This large prospective cohort provided original results on weight variation between before and after cancer diagnosis, highlighting different weight trajectories. Socio-demographic and economic factors appeared to strongly influence the risk of weight gain, illustrating social inequalities in health.