



A protein quantity loci approach combined with a genome-wide association study revealed regulators of protein accumulation in legume seeds

Christine Le Signor, Delphine Aime, N.D. Young, Jean-Marie Prosperi, Richard Thompson, Julia Buitink, Judith Burstin, Karine Gallardo-Guerrero

► To cite this version:

Christine Le Signor, Delphine Aime, N.D. Young, Jean-Marie Prosperi, Richard Thompson, et al.. A protein quantity loci approach combined with a genome-wide association study revealed regulators of protein accumulation in legume seeds. 2. International Legume Society Conference - Legumes for a Sustainable World, Legume Society., Oct 2016, Lisbonne, Portugal. 358 p. hal-01604090

HAL Id: hal-01604090

<https://hal.science/hal-01604090>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



Legumes for a Sustainable World

ILS2 | Second International Legume Society Conference

11th - 14th OCT 2016
TRÓIA RESORT | **PORTUGAL**

itob UNIVERSIDADE NOVA DE LISBOA



Oral Communications

14:30-14:40 Oral – S10

A protein quantity loci approach combined with a genome-wide association study revealed regulators of protein accumulation in legume seeds

Le Signor C.¹, Aime D.¹, Sanchez M.¹, Young N.D.², Prosperi J-M.³, Thompson R.D.¹, Buitink J.⁴, Burstin J.¹, Gallardo K.¹

1 - Agroécologie, AgroSup Dijon, INRA, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France

2 - Department of Plant Biology, University of Minnesota, St. Paul, Minnesota, 55108, USA

3 - UMR AGAP, Equipe Génomique évolutive et gestion des populations, Institut national de Recherche Agronomique (INRA), Montpellier, 34060, France

4 - IRHS, INRA, AGROCAMPUS-Ouest, Université d'Angers, SFR 4207 QUASAV, 7 49071 Beaucozéd cedex, France

The seeds of grain legumes provide proteins (20-40% protein content depending on species) for human nutrition and animal feed. The nutritional and technological quality of legume seeds is mainly determined by the composition and functional properties of the seed protein fractions. By accounting for up to 70% of total proteins in mature seeds, the globulins are major determinants of seed quality. In pea, which is one of the grain legumes most cultivated in Europe, the variability in the abundance of globulin polypeptides across lines is significant, which might enable a wide range of food application. To provide knowledge about the molecular determinants underlying differences in seed protein composition, we have used the model legume species *M. truncatula* for which extensive genetics and genomics resources were available (Young and Bharti, 2012), along with comprehensive transcriptomics data related to seed development (Benedito et al., 2008; Thompson et al., 2009). A PQL approach identified the genomic regions controlling variations in seed protein composition, and a translational approach exploiting the resources developed for *M. truncatula* and pea provided a set of candidate genes underlying PQLs conserved between the two species. The candidate gene selection was then refined by genome-wide association studies (GWAS), which enabled the identification of nucleotide variations associated with variations in globulin synthesis and/or maturation.