



Adaptation of Pea to contrasted French regions : simulation of pea varieties with the AZODYN-Pea crop

Maud Benezit, Annabelle Larmure, Nathalie G. Munier-Jolain, Marie-Helene
Jeuffroy

► To cite this version:

Maud Benezit, Annabelle Larmure, Nathalie G. Munier-Jolain, Marie-Helene Jeuffroy. Adaptation of Pea to contrasted French regions: simulation of pea varieties with the AZODYN-Pea crop. 2. International Legume Society Conference - Legumes for a Sustainable World, Legume Society., Oct 2016, Lisbonne, Portugal. 358 p. hal-01604092

HAL Id: hal-01604092

<https://hal.science/hal-01604092>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



Legumes for a Sustainable World

ILS2 | Second International Legume Society Conference

11th - 14th OCT 2016
TRÓIA RESORT | **PORTUGAL**



UNIVERSIDADE
NOVA
DE LISBOA



P123- S13

**Adaptation of Pea to contrasted French regions:
simulation of pea varieties with the AZODYN-Pea crop**

Bénézit M.¹, Larmure A.², Munier-Jolain N.², Jeuffroy M.-H.¹

1 - UMR Agronomie INRA, AgroParisTech, Université Paris Saclay, 78850 Thiverval-Grignon, France

2 - UMR agroécologie, Dijon

Pea (*Pisum sativum* L.) is a particularly sensitive crop regarding abiotic stress throughout its cycle (Schneider and Huyghe 2015). Climate change results in the increase of the unpredictability of both the frequency and the intensity of these stresses (Stocker et al. 2013). To avoid abiotic stress, breeders have been developing, for the last fifteen years, winter pea varieties sown during autumn, more resistant to frost with an earlier flowering date than spring pea, as well as varieties called "Hr", photoperiod-sensitive, that can be sown even earlier than regular winter types. However, despite the progress of winter type breeding, winter pea surfaces seldom reached more than 25000ha since 2000. Our aim was to study the performances of the different types of pea in contrasted French regions, for various climates, in order to adapt the choice of the pea type to the frequency and intensity of abiotic stress. For this purpose, the crop model Azodyn-Pea (Jeuffroy et al. 2012) was adapted to simulate various types of pea. The model was then used for the evaluation of the three types over the past ten years in different French regions in order to identify which type is most suited to each environment (climate, region). Here we show the potential of winter pea for various French climates. Azodyn-Pea will also allow, in a second phase, the exploration of novel combinations of plant traits in order to design ideotypes and contribute to the development of new varieties within each type.