



HAL
open science

Genome-Wide association mapping of frost tolerance in *Pisum sativum*

Sana Beji, Véronique Fontaine, R. Devaux, Sandra Silvia Negro, Nasser Bahrman, Gregoire Aubert, Bruno Delbreil, Isabelle Lejeune-Henaut

► To cite this version:

Sana Beji, Véronique Fontaine, R. Devaux, Sandra Silvia Negro, Nasser Bahrman, et al.. Genome-Wide association mapping of frost tolerance in *Pisum sativum*. 2. International Legume Society Conference - Legumes for a Sustainable World, Legume Society., Oct 2016, Lisbonne, Portugal. 358 p. hal-01604094

HAL Id: hal-01604094

<https://hal.science/hal-01604094>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



Legumes for a Sustainable World

ILS2 | Second International Legume Society Conference

11th - 14th OCT 2016
TRÓIA RESORT | **PORTUGAL**



UNIVERSIDADE
NOVA
DE LISBOA



17:00-17:10 Oral – S18

**Genome-Wide association mapping of frost tolerance in
*Pisum sativum***

**Beji S.¹, Fontaine V.², Devaux R.³, Negro S.^{5,6}, Bahrman N.², Aubert G.⁴,
Delbreil B.¹, Lejeune-Hénaut I.²**

1 - USC ICV-AFP, Université Lille1, 59655, Villeneuve d'Ascq, France

2 - USC ICV-AFP, INRA, 80203, Péronne, France

3 - UE GCIE, INRA, 80203, Péronne, France

4 - UMR Agroécologie, INRA, 21065, Dijon, France

5 - UMR GQE, INRA, 91190, Gif sur Yvette, France

6 - UMR MIA-Paris, AgroParisTech, 75005, Paris, France

Genome Wide Association Mapping was performed in pea. Accessions from the pea reference collection were phenotyped for frost in field and controlled conditions, and genotyped using Infinium®BeadChip 15K SNPs (Tayeh et al., 2015). After applying filters of quality control, we obtained 363 accessions and 10739 loci for GWA study. Association analyses were conducted with FaST-LMM software using a mixed model that included a relatedness kinship matrix (K) and a population structure matrix (Q) to control for false positives. The K matrix was generated using two approaches. In the first one, the kinship matrix was estimated with all the 10739 markers. In the second approach, we estimated the kinship, called "K-chr", with all the markers other than those located on the same chromosome as the marker being tested (Rincent et al., 2014). Simulations revealed that Rincent's approach was more powerful than the mixed model taking into account a general kinship (estimated for all chromosomes). The GWA study identified 8 loci distributed over different chromosomes comprising 61 SNPs significantly associated with frost tolerance. Results confirmed 3 QTLs that were previously mapped using bi-parental populations and identified 3 novel tolerance loci. Several potential-candidate genes were found corresponding to these SNPs. Additionally, the analyses allowed to identify haplotypes with increased frost tolerance and accessions with favourable alleles for this trait.