



**HAL**  
open science

## Review : Microbial Volatile Organic Compounds (mVOCs) emissions by soil

Letizia Abis, Raluca Ciuraru, Florence Lafouge, Benjamin Loubet,  
Pierre-Alain Maron, Sophie Sadet-Bourgeteau

### ► To cite this version:

Letizia Abis, Raluca Ciuraru, Florence Lafouge, Benjamin Loubet, Pierre-Alain Maron, et al.. Review : Microbial Volatile Organic Compounds (mVOCs) emissions by soil. SOMmic, Nov 2016, Leipzig, Germany. , 2016. hal-01604106

**HAL Id: hal-01604106**

**<https://hal.science/hal-01604106>**

Submitted on 3 Feb 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License




## ABSTRACT

Volatiles Organic Compounds (VOCs) have a central role in the chemistry and composition of the troposphere. VOCs are precursors of atmospheric aerosols (PM 2.5) and tropospheric ozone (O<sub>3</sub>). Biogenic VOCs (bVOCs) are estimated to contribute up to 90% of the total VOCs emissions globally. One of the sources of bVOC is soil microorganisms, especially through sugar degradation but also through secondary metabolism processes. Microbial VOCs production in soils depends also on: nutrients, physiological state of the microorganisms, pH, temperature and oxygen availability. The aim of this poster shows an overview of the existing knowledge on VOC emissions by microorganisms in soil, the reasons why VOCs are emitted and which are the metabolic pathway involved. A focus on the most used techniques to detect VOCs in soils is given and in particular advantages of PTR-MS technique are shown.

## METABOLIC PATHWAY FOR THE VOCs PRODUCTION

### SUGAR DEGRADATION




PYRUVATE, GLYCERALDEHYDE-3-PHOSPHATE, LACTATE, ACETATE


PRECURSORS OF THE VOCs FORMATION

**SECONDARY METABOLITES:** play a central role in inter-organismic interactions. Innumerable cases volatile and non-volatile secondary metabolites have been implicated in defense and communication among organisms.

### FATTY ACID DEGRADATION




### SULPHUR REDUCTION




#### MOST IMPORTANT VOLATILE SULPHUR COMPOUNDS

- H<sub>2</sub>S,
- methanethiol,
- dimethyl sulphide (DMS),
- dimethyl disulphide (DMDS)
- dimethyl trisulphide (DMTS)

### ALCOHOLIC FERMENTATION



### AMINOACID BIOSYNTHESIS



2-PHENYLETHANOL: ONE OF THE MOST WIDESPREAD VOLATILE AROMATIC COMPOUNDS (synthesized by phenolopyruvate)

## WHAT CAN AFFECT MICROBIAL VOCs PRODUCTION

Microbial VOCs production in soils is strongly depending on:

1. NUTRIENTS
2. OXYGEN AVAILABILITY
3. PHYSIOLOGICAL STATE OF THE MICROORGANISMS

they also depend on

- SOIL MOISTURE
- SOIL TEXTURE
- MICROBIAL ACTIVITY
- pH
- TEMPERATURE

**MOST INFLUENT PARAMETER**

**O<sub>2</sub> HIGHLY EFFECTIVE RESPIRATION**


MICRO- AEROBIC OR ANAEROBIC CONDITION:

- I. C-org ARE THE END PRODUCT OF HETERO AND HOMO FERMENTATIVE PROCESS;
- II. DIVERSITY AND AMOUNT OF VOCs EMITTED INCREASE

## VOCs DETECTION TECHNIQUES

### PTR-MS



#### HOW IT WORKS




#### ADVANTAGES

- SIMULTANEOUS ON-LINE MONITORING OF VOLATILE ORGANIC COMPOUNDS;
- LIMIT OF DETECTION <10 pptv
- TIME OF FLIGHT DETECTOR:
- SPEED
- HIGH RESOLUTION


### GAS CHROMATOGRAPHY



### MASS SPECTROMETRY



### FLAME IONIZATION DETECTOR



### FLAME PHOTOMETRIC DETECTOR

THESE TECHNIQUES NEED A METHOD FOR THE VOCs EXTRACTION FROM SAMPLES (e.g. SOLID PHASE MICRO-EXTRACTION (SPME))