



A transcriptomic approach identifies candidate genes for drought tolerance during the reproductive phase in pea

Vanessa Vernoud, Nadia Rossin, Marion Prudent, Christine Le Signor, Myriam Sanchez, Stephanie Pateyron, Sandrine Balzergue, Gregoire Aubert, Judith Burstin, Karine Gallardo-Guerrero, et al.

► To cite this version:

Vanessa Vernoud, Nadia Rossin, Marion Prudent, Christine Le Signor, Myriam Sanchez, et al.. A transcriptomic approach identifies candidate genes for drought tolerance during the reproductive phase in pea. 2. International Legume Society Conference - Legumes for a Sustainable World, Legume Society., Oct 2016, Lisbonne, Portugal. 358 p. hal-01604112

HAL Id: hal-01604112

<https://hal.science/hal-01604112>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



Legumes for a Sustainable World

ILS2 | Second International Legume Society Conference

11th - 14th OCT 2016
TRÓIA RESORT | PORTUGAL



P156 – S18

A transcriptomic approach identifies candidate genes for drought tolerance during the reproductive phase in pea.

Vernoud V.¹, Rossin N.¹, Prudent M.¹, Le Signor C.¹, Sanchez, M.¹,
Pateyron S.², Balzergues S.², Aubert G.¹, Burstin J.¹, Gallardo K.¹,
Thompson R.¹.

1 - Agroécologie, AgroSup Dijon, CNRS, INRA, Univ. Bourgogne Franche-Comté, F-21000
Dijon, France.

2 - POPS Transcriptomic Platform, IPS2, CNRS, INRA, Univ 12 Paris Sud, Univ Evry, Univ
Paris-Diderot, Sorbonne Paris-Cite, Univ Paris-Saclay, 91405 Orsay, France

Drought is a major environmental factor limiting the productivity of crop plants. In pea drought stress occurring during the reproductive phase can greatly affect seed yield and quality. We investigated the response of pea plants (var. Caméor) subjected to water stress during the seed filling period, a phase when massive remobilization from the vegetative organs occurs to sustain seed high-nitrogen demand. Pea plants were subjected to drought stress at the beginning of the seed-filling period of the first two nodes for 8 days. Total and one-seed biomass decreased by 35% and 20% respectively by this limited water stress. Nitrogen allocation to the different plant compartments was also affected, with an increased N allocation to the leaves from the vegetative nodes and a decrease to the root and seed compartments. Transcriptomic changes in water-stressed leaves from the vegetative nodes were analysed by hybridization of a 40k pea micro-array: 178 genes were at least two fold up-regulated in water-stressed samples compared to well-watered samples, whereas 55 genes were down-regulated. Among the most strongly up-regulated genes were those encoding a glutamine amidotransferase, a sucrose transporter from the SWEET family, a phosphoenolpyruvate carboxykinase and a carotenoid cleavage dioxygenase involved in strigolactone biosynthesis.

Acknowledgement: this project was supported by the Burgundy Region (AGRALE6, FABER program) and the FP7 LEGATO project.