



**HAL**  
open science

## Soil, the orphan hydrological compartment

Christophe Hissler, Arnaud Legout, François Barnich, Laurent Pfister

► **To cite this version:**

Christophe Hissler, Arnaud Legout, François Barnich, Laurent Pfister. Soil, the orphan hydrological compartment: Evidence from O and H stable isotopes?. 2015. European Geosciences Union General Assembly (EGU), Apr 2015, Vienne, Austria. , 17, 2015. hal-01604184

**HAL Id: hal-01604184**

**<https://hal.science/hal-01604184>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



## **Soil, the orphan hydrological compartment: evidence from O and H stable isotopes?**

Christophe Hissler (1), Arnaud Legout (2), François Barnich (1), and Laurent Pfister (1)

(1) ERIN/LIST, Belvaux, Luxembourg (christophe.hissler@list.lu), (2) BEF/INRA - UR 1138, Champenoux, France (legout@nancy.inra.fr)

O and H stable isotopes have been successfully used for decades for studying the exchange of waters between the hydrosphere, the pedosphere and the biosphere. They greatly contribute to improve our understanding of soil-water-plant interactions. In particular, the recent hydrological concept of “two water worlds” (separation of meteoric water that infiltrates the soil as (i) mobile water, which can reach the groundwater and can enter the stream, and as (ii) tightly bound water, which is trapped in the soil microporosity and used by plants) calls for a substantial revision of our perceptual models of runoff generation. Nevertheless, there is a need for testing the applicability of this concept over a large range of ecosystemic contexts (i.e. and vegetation types).

To date, many investigations have focused on the relationship between the various processes triggering isotope fractionation within soils. So far, the dominating perception is that the isotope profile of water observed in soils is solely due to evaporative fractionation and its shape is dependent on climate and soil parameters. However, as of today the influence of biogeochemical processes on the spatio-temporal variability of  $\delta^{18}\text{O}$  and  $\delta\text{D}$  of the soil solutions has been rarely quantified. O and H exchanges between soil water and other soil compartments (living organisms, minerals, exchange capacity, organic matter) remain poorly known and require deeper investigations. Eventually, we need to better understand the distribution of O and H isotopes throughout the soil matrix.

In order to address these issues, we have designed and carried out two complementary isotope experiments that use one liter soil columns of a 2mm-sieved and air-dried soil. Our objectives were (1) to observe the temporal evolution of the water O and H isotopic composition starting from the field capacity to the complete drying of the soil and (2) to determine the impact of soil biogeochemical properties on the isotopic composition of different water types in soil (weakly-, moderately- and tightly-bound).

Our results show that mobile and tightly bound water may have different hydrogen isotopic signatures and that their respective isotopic signatures may vary between horizons and soil types. However, it is not yet possible to quantify the contribution of different bio-physico-chemical processes to the oxygen and hydrogen isotopic composition of the soil water because the techniques at hand for water separation are not yet reliable enough. Prior to this type of quantifications, we need to focus in a next step at the improvement of water extraction methods.