



**HAL**  
open science

## Effects of climate data aggregation on regional NPP modelling

Matthias Kuhnert, Jagadeesh Yeluripati, Pete Smith, Holger Hoffmann, Marcel van Oijen, Julie Constantin, Elsa Coucheney, Rene Dechow, Henrik Eckersten, Thomas Gaiser, et al.

### ► To cite this version:

Matthias Kuhnert, Jagadeesh Yeluripati, Pete Smith, Holger Hoffmann, Marcel van Oijen, et al.. Effects of climate data aggregation on regional NPP modelling. 8. International Congress on Environmental Modelling and Software (iEMSs 2016), Jul 2016, Toulouse, France. Sauvage, S., Sánchez-Pérez, J.M., Rizzoli, A.E., 3, 2016, iEMSs 2016 Conference. Environmental modelling and software for supporting a sustainable future. hal-01604211

**HAL Id: hal-01604211**

**<https://hal.science/hal-01604211v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Jul 11th, 5:30 PM - 5:50 PM

# Effects of climate data aggregation on regional net primary production (NPP) modelling

Matthias Kuhnert

*Institute of Biological and Environmental Sciences, University of Aberdeen, matthias.kuhnert@abdn.ac.uk*

Jagadeesh Yeluripati

*Institute of Biological and Environmental Sciences, University of Aberdeen, The James Hutton Institute*

Pete Smith

*Institute of Biological and Environmental Sciences, University of Aberdeen*

Holger Hoffmann

*Crop Science Group, INRES, University of Bonn*

Marcel van Oijen

*Centre for Ecology and Hydrology, CEH-Edinburgh*

*See next page for additional authors*

Follow this and additional works at: <http://scholarsarchive.byu.edu/iemssconference>

 Part of the [Civil Engineering Commons](#), [Data Storage Systems Commons](#), [Environmental Engineering Commons](#), [Hydraulic Engineering Commons](#), and the [Other Civil and Environmental Engineering Commons](#)

Matthias Kuhnert, Jagadeesh Yeluripati, Pete Smith, Holger Hoffmann, Marcel van Oijen, Julie Constantin, Elsa Coucheney, Rene Dechow, Henrik Eckersten, Thomas Gaiser, Balász Grosz, Edwin Haas, Kurt-Christian Kersebaum, Ralf Kiese, Steffen Klatt, Elisabet Lewan, Claas Nendel, Helene Raynal, Carmen Sosa, Xenia Specka, Edmar Teixeira, Enli Wang, Lutz Weihermüller, Gang Zhao, Zhigan Zhao, Stephen Ogle, and Frank Ewert, "Effects of climate data aggregation on regional net primary production (NPP) modelling" (July 11, 2016). *International Congress on Environmental Modelling and Software*. Paper 51.  
<http://scholarsarchive.byu.edu/iemssconference/2016/Stream-B/51>

This Event is brought to you for free and open access by the Civil and Environmental Engineering at BYU ScholarsArchive. It has been accepted for inclusion in International Congress on Environmental Modelling and Software by an authorized administrator of BYU ScholarsArchive. For more information, please contact [scholarsarchive@byu.edu](mailto:scholarsarchive@byu.edu).

---

**Presenter/Author Information**

Matthias Kuhnert, Jagadeesh Yeluripati, Pete Smith, Holger Hoffmann, Marcel van Oijen, Julie Constantin, Elsa Coucheney, Rene Dechow, Henrik Eckersten, Thomas Gaiser, Balász Grosz, Edwin Haas, Kurt-Christian Kersebaum, Ralf Kiese, Steffen Klatt, Elisabet Lewan, Claas Nendel, Helene Raynal, Carmen Sosa, Xenia Specka, Edmar Teixeira, Enli Wang, Lutz Weihermüller, Gang Zhao, Zhigan Zhao, Stephen Ogle, and Frank Ewert

# Effects of climate data aggregation on regional net primary production (NPP) modelling

**Kuhnert, Matthias<sup>a,\*</sup>; Yeluripati, Jagadeesh<sup>a,m</sup>; Smith, Pete<sup>a</sup>; Hoffmann, Holger<sup>b</sup>; van Oijen, Marcel<sup>c</sup>; Constantin, Julie<sup>d</sup>; Coucheney, Elsa<sup>e</sup>; Dechow, René<sup>f</sup>; Eckersten, Henrik<sup>g</sup>; Gaiser, Thomas<sup>b</sup>; Grosz, Balázs<sup>f</sup>; Haas, Edwin<sup>h</sup>; Kersebaum, Kurt-Christian<sup>i</sup>; Kiese, Ralf<sup>h</sup>; Klatt, Steffen<sup>h</sup>; Lewan, Elisabet<sup>e</sup>; Nendel, Claas<sup>i</sup>; Raynal, Helene<sup>d</sup>; Sosa, Carmen<sup>e</sup>; Specka, Xenia<sup>i</sup>; Teixeira, Edmar<sup>j</sup>; Wang, Enli<sup>k</sup>; Weihermüller, Lutz<sup>i</sup>; Zhao, Gang<sup>b</sup>; Zhao, Zhigan<sup>k</sup>; Ogle, Stephen<sup>n</sup>; Ewert, Frank<sup>b</sup>**

<sup>a</sup> Institute of Biological and Environmental Sciences, University of Aberdeen, 23 St Machar Drive, Aberdeen AB24 3UU, UK

<sup>b</sup> Crop Science Group, INRES, University of Bonn, Katzenburgweg 5, 53115 Bonn, DE

<sup>c</sup> Centre for Ecology and Hydrology, CEH-Edinburgh, Bush Estate, Penicuik EH26 0QB, UK

<sup>d</sup> INRA, UMR 1248 AGIR & UR0875 MIA-T, F-31326 Auzeville, France

<sup>e</sup> Department of Soil and Environment, Swedish University of Agricultural Sciences, Lennart Hjelms väg 9, 750 07 Uppsala, SE

<sup>f</sup> Thünen-Institute of Climate-Smart-Agriculture, Bundesallee 50, 38116 Braunschweig, DE

<sup>g</sup> Department of Crop Production Ecology, Swedish University of Agricultural Sciences, Ulls väg 16, 750 07 Uppsala, SE

<sup>h</sup> Institute of Meteorology and Climate Research – Atmospheric Environmental Research, Karlsruhe Institute of Technology, Kreuzeckbahnstraße 19, 82467 Garmisch-Partenkirchen, DE

<sup>i</sup> Institute of Landscape Systems Analysis, Leibniz Centre for Agricultural Landscape Research, 15374 Müncheberg, DE

<sup>j</sup> Systems Modelling Team (Sustainable Production Group), The New Zealand Institute for Plant and Food Research Limited, Canterbury Agriculture & Science Centre, Gerald St, Lincoln 7608, NZ

<sup>k</sup> CSIRO Land and Water, Clunies Ross Street, Canberra, ACT, AU

<sup>l</sup> Institute of Bio- & Geosciences Agrosphere (IBG-3), Forschungszentrum Jülich, 52425 Jülich, DE

<sup>m</sup> The James Hutton Institute, Craigiebuckler, Aberdeen, AB15 8 QH, UK

<sup>n</sup> Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO 80523, USA

\*corresponding author: [matthias.kuhnert@abdn.ac.uk](mailto:matthias.kuhnert@abdn.ac.uk)

## Abstract:

In ecosystem modelling studies, data are often collected at a small scale but are used in models to predict ecosystem responses e.g. net primary production (NPP) at coarser scale, using data aggregation. The data aggregation causes errors that are difficult to predict, because of model complexity and our limited knowledge of the spatial heterogeneity underlying our aggregated input values. Input aggregation thus adds to the uncertainty associated with model predictions. One way of reducing uncertainty related to input aggregation error is to evaluate models at different resolutions for areas where high-resolution input data are available. The objective of this study was the quantification of the aggregation error on modelled NPP introduced by climate data aggregation. Therefore, climate data at 1 km x 1 km resolution (>30 000 grid cells) were used as baseline to simulate NPP with 11 different crop and biogeochemical models for the federal state North Rhine-Westphalia (Germany). These results, for 29 year monocultures of wheat and maize cropping systems, respectively, were compared with simulation results using aggregated climate data for four resolutions (10, 25, 50, 100 km grid cell side length). The aggregation effect is represented by the maximum differences between the NPP, averaged for wheat and maize cropping systems over the growing season, simulated for the five different resolutions. Input data aggregation had little impact on NPP of 29 year averages (0.5 – 7.8 % for wheat and 0.3 – 10 % for maize), while the climate data aggregation effect was higher for single years; up to 9 % and 13 % for wheat and maize, respectively, gradually decreasing to low effects for averages over 10 year periods or longer. The scale effect differed among models and shows only a minor impact (2%) for an ensemble run.

**Keywords:** net primary production, NPP, scaling, extreme events, crop modelling, climate, data aggregation