



**HAL**  
open science

## Analysing data aggregation effects on large-scale yield simulations

Holger Hoffmann, Gang Zhao, Senthold Asseng, Marco Bindi, Davide Cammarano, Julie Constantin, Elsa Coucheney, Rene Dechow, Helene Raynal, Frank Ewert

### ► To cite this version:

Holger Hoffmann, Gang Zhao, Senthold Asseng, Marco Bindi, Davide Cammarano, et al.. Analysing data aggregation effects on large-scale yield simulations. iCROP 2016 International Crop Modelling Symposium "Crop Modelling for Agriculture and Food Security under Global Change", Mar 2016, Berlin, Germany. , 441 p., 2016, International Crop Modelling Symposium "Crop Modelling for Agriculture and Food Security under Global Change" 2016. hal-01604215

**HAL Id: hal-01604215**

**<https://hal.science/hal-01604215v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

15-17 March 2016, Berlin, Germany

iCROP M<sub>2016</sub>



MACSUR and AgMIP jointly present the International Crop Modelling Symposium  
**Crop Modelling for Agriculture and Food Security  
under Global Change**



**ABSTRACTS**

## **Editors**

**Frank Ewert**  
**Kenneth J. Boote**  
**Reimund P. Rötter**  
**Peter Thorburn**  
**Claas Nendel**

<b>Welcome Note .....</b>	<b>1</b>
<b>Abstracts of Keynote Presentations.....</b>	<b>4</b>
<b>Abstracts of Oral Presentations.....</b>	<b>17</b>
<b>Abstracts of Poster Presentations .....</b>	<b>184</b>

## **Welcome**

It is with pleasure that we welcome the international crop and agro-ecosystem modelling community and scientists from related disciplines to the International Crop Modelling Symposium 2016, in Berlin.

The past decade has seen a number of research initiatives launched to advance crop modelling and related research. Among these initiatives, The European Knowledge Hub MACSUR (Modelling European Agriculture with Climate Change for Food Security, <http://macsur.eu/>) and the international AgMIP project (Agricultural Model Intercomparison and Improvement Project, <http://www.agmip.org/>) stand out in terms of the breadth of their research scope. A large and important part of the activities in both projects is comprised of the improvement, comparison and application of crop models for climate change impact and risk assessment for food security in Europe (MACSUR) and further (AgMIP). These projects have brought together a large number of scientists from around the world and produced a substantial body of novel results. The international MACSUR symposium on crop modelling in Oslo in 2014 and the annual Global Workshops of AgMIP have provided forums to exchange some of these results and have been initial and important events towards this symposium. The increasing interest from within and beyond the crop modelling community for a more comprehensive forum for the exchange of results ultimately motivated representatives of MACSUR and AgMIP to organise this symposium, reflecting the successful and joint work of both projects including successful interaction with other international networks.


The overwhelming interest in participation in this symposium has exceeded original expectations. From the large number of submitted papers, it was possible to develop what we, the Symposium Chairs, hope is an exciting programme of oral and poster presentations combined with a range of internationally recognised keynote speakers. The workshop structure follows the main activities related to model improvement and model application, as well as anticipating improvements in genetics, and links between crop and related modelling fields such as grassland and vegetation modelling, and

functional structural plant modelling. Accordingly, four sessions have been organised:

- Session 1: Improvement of crop models and modelling approaches
- Session 2: Linking crop models and genetics
- Session 3: Crop modelling for risk/impact assessment
- Session 4: Expanding and supporting modelling activities

The organisation of this symposium was only possible due to the help of several people. Special thanks go to the Session Chairs and the Scientific Committee Members for supporting the development of the symposium programme. We are particularly grateful for the effort of the local host ZALF (Centre for Agricultural Landscape Research) for organising the venue, registration, website and logistics of the programme. The financial and in-kind support from the Research Council of Norway through MACSUR, CSIRO, AgMIP, University of Bonn, Luke and the University of Florida are likewise gratefully acknowledged.

We wish all participants a very fruitful and inspiring symposium and we look forward to the many interesting keynotes, oral and poster presentations. We also hope to have the chance to interact with many of you during the course of the symposium and that the symposium may help to support ongoing and initiate new collaborations to further advance research on crop modelling.



Frank Ewert

On behalf of the Symposium Chairs, Kenneth J Boote, Peter Thorburn and Reimund Rötter, and the local host at ZALF, Claas Nendel.

International Crop Modelling Symposium **iCROPM<sub>2016</sub>** 15-17 March 2016, Berlin

# **Abstracts of Keynote Presentations**

### **Analysing data aggregation effects on large-scale yield simulations**

*H. Hoffmann*<sup>1\*</sup> – G. Zhao<sup>1</sup> – S. Asseng<sup>2</sup> – M. Bindi<sup>3</sup> – D. Cammarano<sup>4</sup> – J. Constantin<sup>5</sup> – E. Coucheney<sup>6</sup> – R. Dechow<sup>7</sup> – L. Doro<sup>8</sup> – H. Eckersten<sup>9</sup> – T. Gaiser<sup>1</sup> – B. Grosz<sup>7</sup> – E. Haas<sup>10</sup> – B. Kassie<sup>2</sup> – K. C. Kersebaum<sup>11</sup> – R. Kiese<sup>10</sup> – S. Klatt<sup>10</sup> – M. Kuhnert<sup>12</sup> – E. Lewan<sup>6</sup> – M. Moriondo<sup>13</sup> – C. Nendel<sup>11</sup> – H. Raynal<sup>5</sup> – P. P. Roggero<sup>8</sup> – R. Rötter<sup>14</sup> – S. Siebert<sup>1</sup> – C. Sosa<sup>6</sup> – X. Specka<sup>11</sup> – F. Tao<sup>14</sup> – E. Teixeira<sup>5</sup> – G. Trombi<sup>3</sup> – J. Yeluripati<sup>16</sup> – E. Vanuytrecht<sup>17</sup> – D. Wallach<sup>5</sup> – E. Wang<sup>18</sup> – L. Weihermüller<sup>19</sup> – Z. Zhao<sup>18</sup> – F. Ewert<sup>1</sup>

<sup>1</sup> Crop Science Group, INRES, University of Bonn, Katzenburgweg 5, 53115 Bonn, DE

<sup>2</sup> Agricultural & Biological Engineering Department, University of Florida, Frazier Rogers Hall, Gainesville, FL 32611, USA

<sup>3</sup> Department of Agri-food Production and Environmental Sciences - University of Florence, Piazzale delle Cascine 18, 50144 Firenze, IT

<sup>4</sup> The James Hutton Institute, Invergowrie Dundee DD2 5DA, Scotland UK

<sup>5</sup> INRA, UMR 1248 AGIR & UR0875 MIA-T, F-31326 Castanet-Tolosan, FR

<sup>6</sup> Biogeophysics and water quality, Department of Soil and Environment, Swedish University of Agricultural Sciences, Lennart Hjelm's väg 9, 750 07 Uppsala, SE

<sup>7</sup> Thünen-Institute of Climate-Smart-Agriculture, Bundesallee 50, 38116 Braunschweig, DE

<sup>8</sup> Desertification Research Group, Università degli Studi di Sassari, Viale Italia 39, 07100 Sassari, IT

<sup>9</sup> Department of Crop Production Ecology, Swedish University of Agricultural Sciences, Ulls väg 16, 750 07 Uppsala, SE

<sup>10</sup> Institute of Meteorology and Climate Research – Atmospheric Environmental Research, Karlsruhe Institute of Technology, Kreuzackbahnstraße 19, 82467 Garmisch-Partenkirchen, DE

<sup>11</sup> Institute of Landscape Systems Analysis, Leibniz Centre for Agricultural Landscape Research, 15374 Müncheberg, DE

<sup>12</sup> Biological and Environmental Sciences, School of Biological Sciences, University of Aberdeen, 23 St Machar Drive, Aberdeen AB24 3 UU, Scotland, UK

<sup>13</sup> Marco Moriondo, CNR-Ibimet, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy

<sup>14</sup> Climate Impacts Group, Natural Resources Institute Finland (Luke), 00790 Helsinki, FI

<sup>15</sup> Systems Modelling Team (Sustainable Production Group), The New Zealand Institute for Plant and Food Research Limited, Canterbury Agriculture & Science Centre, Gerald St, Lincoln 7608, NZ

<sup>16</sup> The James Hutton Institute, Craigiebuckler, Aberdeen, AB15 8 QH, UK

<sup>17</sup> Division Soil & Water Management, KU Leuven, Celestijnenlaan 200E, PO 2411, 3001 Heverlee, BE

<sup>18</sup> CSIRO Land and Water, Clunies Ross Street, Canberra, ACT, AU

<sup>19</sup> Institute of Bio- & Geosciences Agrosphere (IBG-3), Forschungszentrum Jülich, 52425 Jülich, DE

\* hhoffmann@uni-bonn.de

### **Introduction**

Large-scale yield simulations often use data of coarse spatial resolution as input for process-based models (Ewert et al., 2015; Zhao et al., 2015). However, using aggregated data as input for process-based models entails the risks of introducing errors linked to aggregation effects (AE) such as: i) data modification; ii) missing the valid range of the model; iii) data inconsistencies between data types. While the regional crop yield bias is usually <5 % on average over all years, it may increase to more than 10 % under specific conditions, e.g. in single years (Hoffmann et al., 2015), depending on the model. In order to assess these differences in detail, we present a model intercomparison on AE for a range of environmental conditions differing in climate and soil for two crops grown under three different production situations.



### **Materials and Methods**

Multi-model ensemble runs were conducted with soil and climate input data at resolutions from 1 to 100 km for the state of North Rhine-Westphalia, Germany. Climate data was spatially averaged. Soil data was aggregated by area majority. Winter wheat and silage maize yields of 1982-2011 were simulated with 11 models for potential, water-limited and water-nitrogen-limited production after calibration to average regional sowing date, harvest date and crop yield.

### **Results and Discussion**

Regional yields were reproduced by the models on average, regardless of input data type and resolution. However, *AE* were observed in dry years as well as due to soil aggregation. Large positive *AE* between coarser and 1 km resolutions were associated with low soil water holding capacity (SWHC) and low climatic water balance (CWB) at 1 km resolution in combination with increases in SWHC and CWB when aggregating to coarser resolutions. Consistently, the opposite was found for large negative *AE*. Large *AE* (larger/lower mean +/-2 standard deviations of *AE*) were due to changes in soil type. However, the lower 50 % of all *AE* also showed differences in the aggregation method: soil aggregation by majority led to about 20 % of grid cells with no *AE* at 100 km resolution whereas climate data aggregation by averaging always led to *AE*. Still, soil aggregation by majority led to a larger fraction of *AE* larger than 10 % as compared to climate averaging. Notably, this was further increased by the combined use of aggregated soil and climate data. Finally, models differed considerably in *AE*.

### **Conclusions**

The results highlight the interactions between model, data and aggregation method with *AE*, emphasizing the importance of models intercomparison analyses.

### **Acknowledgements**

The study was conducted in the frame of the FACCE JPI MACSUR and received funding from several national institutions. Please see Hoffmann et al., (2015) and Zhao et al., (2015) for further details on funding.

### **References**

- Hoffman, H., G. Zhao, L.G.J. Van Bussel et al., (2015). *Climate Research*, doi: 10.3354/cr01326.
- Ewert F., L.G.J. Van Bussel, G. Zhao et al., (2015). In: Rosenzweig, C. and Hillel, D: *Handbook of Climate Change and Agroecosystems*. Imperial College Press, London, UK, 261-277.
- Zhao, G., H. Hoffmann, L.G.J. Van Bussel et al., (2015). *Climate Research*, doi: 10.3354/cr01301.