



Impact of embryonic heat conditioning on bird epigenome

Vincent Coustham, Nathalie Couroussé, Sabine Crochet, Sarah-Anne David, Thierry Bordeau, Estelle Godet, Benoit Piegu, Celine Noiro, Yves Bigot, Frederique Pitel, et al.

► To cite this version:

Vincent Coustham, Nathalie Couroussé, Sabine Crochet, Sarah-Anne David, Thierry Bordeau, et al.. Impact of embryonic heat conditioning on bird epigenome. EMBO Conference: Chromatin and Epigenetics, May 2017, Heidelberg, Germany. hal-01604366

HAL Id: hal-01604366

<https://hal.science/hal-01604366>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Impact of embryonic heat conditioning on bird epigenome

Coustham Vincent¹, Couroussé Nathalie¹, Crochet Sabine¹, David Sarah-Anne¹, Bordeau Thierry¹, Godet Estelle¹, Piégu Benoît², Noirot Céline³, Bigot Yves², Pitel Frédérique⁴, Collin Anne¹.

¹INRA – UR83 Recherches Avicoles – 37380 NOUZILLY

²INRA – UMR85 PRC – 37380 NOUZILLY

³INRA - UR875 MIAT – 31326 CASTANET-TOLOSAN

⁴INRA – UMR1388 GenPhySE – 31326 CASTANET-TOLOSAN

Corresponding author:
vincent.coustham@inra.fr

Changes in gene activity through epigenetic alterations induced by early environmental challenges during the embryogenesis are known to impact the phenotype, health and disease risk of animals. The epigenome is therefore an essential contributor to phenotypic plasticity, and learning how environmental exposures translate into persisting epigenetic changes may open new doors to improve robustness and resilience of developing animals.

Birds are agronomic species of choice to directly manipulate the embryonic environment and study its consequences on the developing animal. We previously showed that the heat tolerance of male commercial chickens was improved by cyclically elevating the egg incubation temperature. This treatment named embryonic thermal manipulation (TM) was associated with changes in gene expression that persisted during the development of chickens and enhanced gene expression response in case of heat challenge at slaughter age, 35 days post-hatch.

To further explore the molecular basis of heat acclimation, we took advantage of an inbred line of Japanese quails (*Coturnix japonica*) to investigate the impact of TM on bird methylome. Among other advantages, quail generation cycle is 3-4 time faster than chicken and the use of an inbred genotype should reduce the phenotype variations associated with genetic variability. We performed a characterization of TM on quail phenotype in interaction with a post-hatch heat stress at 35 days of age. Several parameters such as weight of the animals, temperature, egg laying for females, hormone levels and blood gas characteristics were measured. Concomitantly we investigated the impact of TM on quail transcriptome by RNA-seq and methylome by a whole genome bisulfite sequencing (WGBS) on brain tissues sampled at hatch. Analysis of the results is ongoing and preliminary data will be presented.