



HAL
open science

Genetic dissection of tomato fruit quality in the genome era: new tools for in depth QTL characterization

Elise Albert, Laura Pascual-Banuls, Christopher Sauvage, Janejira Duangjit, Jean-Paul Bouchet, Frederique Bitton, Isidore Diouf, Yolande Carretero, Mathilde Causse

► To cite this version:

Elise Albert, Laura Pascual-Banuls, Christopher Sauvage, Janejira Duangjit, Jean-Paul Bouchet, et al.. Genetic dissection of tomato fruit quality in the genome era: new tools for in depth QTL characterization. 3. International Symposium on Horticulture in Europe, SHE2016, Oct 2016, Chania, Greece. 2016. hal-01604736

HAL Id: hal-01604736

<https://hal.science/hal-01604736>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ISHS Conference



III International Symposium on Horticulture in Europe



Growing Health and Life

Programme & Book of Abstracts

CHANIA, CRETE, GREECE, OCTOBER 17-21, 2016

III International Symposium
on Horticulture in Europe
SHE2016

Programme
and
Book of Abstracts

Chania, Greece
17 - 21 October, 2016

Conveners
Dr. Panagiotis Kalaitzis, Dr. George Manganaris

GENETIC DISSECTION OF TOMATO FRUIT QUALITY IN THE GENOME ERA: NEW TOOLS FOR IN DEPTH QTL CHARACTERIZATION

Elise Albert, Laura Pascual, Christopher Sauvage, Janejira Duangjit, Jean-Paul Bouchet, Frédérique Bitton, Isidore Diouf, Yolande Carretero, Mathilde Causse

INRA, UR1052, Centre de Recherche PACA, 67 Allée des Chênes CS60094, 84143 Montfavet Cedex, France

Tomato fruit quality is characterized by a large number of components influenced by the genotypes and the environment. To better understand the genetic control of these components, quantitative trait loci (QTL) have been mapped for years. Today the genome sequence availability changes the paradigm of genetic approaches. QTL have been first identified using linkage mapping populations and positional cloning identified a few QTLs. However linkage mapping is limited to the analysis of traits differing between two parental lines. Genome-wide association (GWA) has then been proposed to assess a large range of variability. In tomato, we have shown that GWA study is possible, using the admixed nature of cherry tomato genomes that reduces the impact of population structure. Nevertheless, GWA success is limited in some regions due to the low decay of linkage disequilibrium, which varies along the genome. Rare alleles are also difficult to detect with GWA studies. Multi-parent advanced generation intercross (MAGIC) populations offer an alternative to traditional linkage and GWAs by increasing the precision of QTL mapping but with equilibrated allelic frequencies. We have developed a MAGIC population by crossing eight tomato lines whose genomes were resequenced. We showed the potential of the MAGIC population when coupled with whole genome sequencing to detect candidate single nucleotide polymorphisms (SNP) underlying the QTLs. QTLs for fruit quality traits were mapped and related to the variations detected at the genome sequence and expression levels. The advantages and limitations of the three types of population, in the context of the available genome sequence and resequencing facilities, will be discussed.