

#### Contamination of durum wheat by cadmium: lessons from the field

Christophe Nguyen, Emmanuelle Gourdain, Guénolé Grignon, Bruno Barrier-Guillot, Benoit Meleard

#### ▶ To cite this version:

Christophe Nguyen, Emmanuelle Gourdain, Guénolé Grignon, Bruno Barrier-Guillot, Benoit Meleard. Contamination of durum wheat by cadmium: lessons from the field. ICOBTE 2017, Jul 2017, Zurich, Switzerland. 1 p. hal-01605302

#### HAL Id: hal-01605302 https://hal.science/hal-01605302v1

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Contamination of Durum Wheat by Cadmium: Lessons from the Field



Christophe Nguyen<sup>a</sup>, E. Gourdain<sup>b</sup>, M. Jacquard<sup>b</sup>, G. Grignon<sup>b</sup>, B. Barrier-Guillot<sup>b</sup> and B. Méléard<sup>b</sup>







<sup>a</sup> Interactions soil-plant-atmosphere INRA - BSA, France - <sup>b</sup> Arvalis, Institut du Végétal, France christophe.nguyen@inra.fr

#### Cadmium (Cd)


Very toxic (carcinogenic) - Accumulated by crops due to root uptake from soil - European regulatory limits for contents in some foodstuffs (EC 1881/2006) - Dietary exposure to Cd too high in Europe (Efsa, 2009) - Wheat = major contributor to dietary exposure to Cd Project of lowering regulatory limit for Cd in wheat (DGSanco, 2011)

### Cd in French durum wheat compared to current and possible new regulatory limits






### Cd in durum wheat grain : Phytoavailability of Cd in soil > cultivar differences




Contrary to Canada, in France, cultivars are not yet selected for the low Cd accumulation allele (Zimmerl et al., 2014)


## N fertilization increases Cd in durum wheat grain Spliting N dose can help limiting contamination




Increase proteins but not Cd in grains = Split N quantity NH4<sup>+</sup> assumed to desorb Cd and/or acidify soil solution

#### Risk assessment for the contamination of durum wheat grain by Cd







# Probability (grain Cd > limit) = f(Cdtot : total soil Cd, pH : soil pH, Corg : soil organic carbon, Calc : soil calcareous, Clay : soil clay, Silk\_C, Silk\_F : coarse & fine silk) Field data : n=408


**Logitistic models** 

(2008-2016, French production of durum wheat)


# Model performances for field applications

The models predict correctly more than **91%** of the cases

Weak




13 Cdtot + pHeau + Silt\_F + Corg




24 Cdtot + pHeau + Clay + Silt\_F + Silt\_C + Corg

25 Cdtot + pHeau + Clay + Silt\_F + Silt\_C + Calc

26 Cdtot + pHeau + Clay + Silt F + Silt C + Corg + Calc





Conform (-)



Models are sensitive & accurate

Trade-off between detection and reliability

More data are needed to improve models

Non conform (+)

Good

Increase grain Cd: Cdtot, Clay and Silk (buffer power)

Decrease grain Cd: pH (solubility, speciation), Corg (sorption),

Calc (sorption, precipitation)