



**HAL**  
open science

## Are sown flower strips an efficient tool to limit aphid colonization and viral epidemics in melon crops?

Alexandra Schoeny, Nathalie Boissot, Catherine Wipf-Scheibel, Karine Nozeran, Pascale Mistral, Patrick Gognalons, Pauline Millot, Hervé Lecoq

### ► To cite this version:

Alexandra Schoeny, Nathalie Boissot, Catherine Wipf-Scheibel, Karine Nozeran, Pascale Mistral, et al.. Are sown flower strips an efficient tool to limit aphid colonization and viral epidemics in melon crops?. 10. Conférence de la Société française de phytopathologie, 12. European foundation for plant pathology conference, May 2017, Malo-les-Bains, France. , 2017, Deepen knowledge in plant pathology for innovative agro-ecology. hal-01605708

**HAL Id: hal-01605708**

**<https://hal.science/hal-01605708v1>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

# Are sown flower strips an efficient tool to limit aphid colonization and viral epidemics in melon crops?



**Schoeny Alexandra<sup>1</sup>**  
**Boissot Nathalie<sup>2</sup>**  
**Wipf-Scheibel Catherine<sup>1</sup>**  
**Nozeran Karine<sup>1</sup>**  
**Mistral Pascale<sup>2</sup>**  
**Gognalons Patrick<sup>1</sup>**  
**Millot Pauline<sup>1</sup>**  
**Lecoq Hervé<sup>1</sup>**

<sup>1</sup>Pathologie Végétale  
 INRA, 84140, Montfavet, France  
<sup>2</sup>Génétique et Amélioration des  
 Fruits et Légumes  
 INRA, 84140, Montfavet, France

alexandra.schoeny@inra.fr

The melon aphid *Aphis gossypii* is a serious pest on melon crops causing leaf-curling, stunting and even plant death when colonization is intense. It is also an efficient vector of viruses frequently observed on melon crops in France: *Cucurbit aphid-borne yellows virus* (CABYV, persistent), *Cucumber mosaic virus* (CMV, non-persistent) and *Watermelon mosaic virus* (WMV, non-persistent).

The management of aphids and viruses is all the more challenging that the evolution of the regulation (plan Ecophyto 2018) imposes a progressive reduction of the usage of the phytosanitary products. Innovative strategies are needed to control these bioaggressors.

The bibliography suggests that the management of field margins could contribute to regulate the populations of aphids and/or their viral load (1,2). Indeed, flower strips can participate in aphid biological control by favouring natural enemies and strips of non-host plants can protect crops from non-persistently aphid-transmitted viruses by allowing aphids to probe on healthy plants and thus to lose their virus load before reaching the crops.

**The objective was to evaluate the potential role of sown flower strips to reduce the risk of aphid colonization and the risk of viral epidemics in melon crops.**




*Aphis gossypii* infestation on melon



WMV infection on melon




## RESULTS



Entomofauna was highly stimulated by flower strips. In particular, aphid predators such as Coccinellidae and Syrphidae were significantly more abundant near flower strips.

Aphid colonization was consistently reduced (up to 47%) in melon crops with flower strips; still the aphid index remained above the action threshold in 4/5 trials.

Indeed, except in 2012 where no quadrat reached score 2 in FS modality, the management of the field margins wouldn't have avoided an insecticide treatment « in real life ».




The impact of flower strips varied according to the virus: mainly neutral on CABYV, sometimes deleterious on CMV, usually favourable on WMV. Underlying mechanisms remain to be elucidated.

**Sown flower strips alone are not sufficient to control aphids and viruses in melon crops but could positively complement the efficiency of a genetic resistance in an IPM programme (3).**

## METHODS

- Five field experiments conducted in Avignon (2011-2015).
- Experimental design with two modalities of field margins: bare soil (BS) and flower strip (FS) sown with an ad hoc mix (40% sainfoin, 30% grass pea, 20% salad burnet, 5% cornflower, 5% marjoram) two months before melon planting.



- Flying insects sampled with tent traps placed near field margins (2014-2015).
- Aphid colonization assessed with a 0-4 rating scale (3-5 dates per trial)  
 → Aphid Index = area under the pest progress curve standardized by the duration of the monitoring
- Virus detection by double antibody sandwich enzyme-linked immunosorbent assay (DAS-ELISA) with specific polyclonal antisera (7-9 dates per trial)  
 → Mean incidence over the epidemic = area under the disease progress curve standardized by the duration of the monitoring

SCORE	DESCRIPTION
0	No aphids
1	Less than 10 isolated aphids
2	Small colonies (<50 aphids)
3	Large colonies + honeydew
4	Very large colonies (>500 aphids)

## REFERENCES

- 1) Pfiffner L. and Wyss E. (2004) In: Ecological engineering for pest management. Advances in habitat manipulation for arthropods. CSIRO Publishing, Australia, pp. 167-188
- 2) Hooks C.R.E. and Fereres A. (2006) Virus Res. 120, 1-16
- 3) Boissot N., Schoeny A. and Vanlerberghe-Masutti F. (2016) Frontiers Plant Sci. 7(1420), 1-18

## ACKNOWLEDGEMENTS

We thank Joël Béraud, Michel Pascal, Frédéric Pascal and Antoine Lauvernay for their efficient help in field experiments.

This study was funded by INRA (AAP2010 Gestion durable des résistances) and Ministère de l'Agriculture, de l'Alimentation et de l'Environnement (CTPS 25-C-2011-09 and CASDAR 12-1278 Agath).

12<sup>th</sup> European Foundation for Plant Pathology (EFPP)  
 10<sup>th</sup> French Society for Plant Pathology (SFP) Conference

Dunkirk-Malo-les-Bains, France  
 29 May-2 June 2017