

## Wheat sourdoughs: a new model to link ecology and evolution

Belen Carbonetto, Elisa Michel, Yoann Robert, Johan Ramsayer, Charlotte Urien, Judith Legrand, Delphine Sicard

## ► To cite this version:

Belen Carbonetto, Elisa Michel, Yoann Robert, Johan Ramsayer, Charlotte Urien, et al.. Wheat sourdoughs: a new model to link ecology and evolution. EMBO conference New Model Systems for Linking Evolution and Ecology, May 2016, EMBL Heidelberg, Germany. hal-01606190

## HAL Id: hal-01606190 https://hal.science/hal-01606190

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

## Wheat sourdoughs: a new model to link ecology and evolution

Belen Carbonetto<sup>1</sup>, Elisa Michel<sup>2</sup>, Yoann Robert<sup>1</sup>, Charlotte Urien, Johann Ramsayer<sup>1, 3</sup>, Judith Legrand<sup>3</sup>, Delphine Sicard<sup>1</sup>

<sup>1</sup> INRA/Supagro, UMR 1083 Sciences pour l'oenologie, 34060 Montpellier

<sup>2</sup> Oniris, Laboratoire de Microbiologie Alimentaire et Industrielle, CS 82225, 44322 Nantes Cedex 3, France

<sup>3</sup> Univ Paris-Sud, UMR 0320/UMR8120 G´en´etique Quantitative et Evolution—Le Moulon, Universit ´e Paris-Saclay,

F-91190 Gif-sur-Yvette, France

We are studying wheat sourdough as a new ecosystem model for linking ecology and evolution. The sourdough consists of a mixture of flour and water that is fermented by yeasts and lactic acid bacteria (LAB). The baker can make it in a natural way by mixing floor and water in its bakehouse (natural sourdough) or on the opposite can purchase it from outside. Like in batch culture, bakers refresh natural sourdough along and between bread-making process by adding fresh flour and water. The microbial dynamic and the microbial genetic and phenotypic evolution of these semi-opened sink/source ecosystem can be studied, when including bakers in participatory research action. The species diversity of wheat sourdough has been intensively described in European artisan bakers. Most sourdough contains *S. cerevisiae* as dominant yeast species or a species belonging to the Saccharomyces most-closely related clade, Kazachstania. Using metabarcoding analysis (pyrosequencing and Illumina) and identification of over 1000 yeast isolated strains, we have analyzed the species composition of sourdoughs coming from 36 French bakers with diverse socio-cultural practices (farmer-bakers, artisan bakers, companies). We found that geography did not explain the sourdough species composition but bakery practices did. While companies mostly carry *S. cerevisiae* as dominant sourdough species, farmer-bakers harbor much higher species diversity with new yeast species of the Kazachstania clade. The analysis of the metabolic phenotype (i.e. fermentation kinetics, glucose, maltose, acetate, lactate, glycerol and ethanol consumption/production) and population size of sourdough isolated strains and type strains of six Kazachstania species and Saccharomyces cerevisiae revealed phenotypic convergence of sourdoughs strains

belonging to different yeast species. The genome of the *Kazachstania* species are currently sequenced and annotated and will be used to identified the genes that might be associated to the phenotypic convergence, using RNAseq analysis.