

Optimization of control strategies of plant epidemics through a simulation model validated by experiments

Loup Rimbaud, Emmanuel Jacquot, Samuel Soubeyrand, Gael Thébaud

▶ To cite this version:

Loup Rimbaud, Emmanuel Jacquot, Samuel Soubeyrand, Gael Thébaud. Optimization of control strategies of plant epidemics through a simulation model validated by experiments. 1. Rencontres du GdR Ecologie Statistique, Mar 2015, Lyon, France. hal-01606295

HAL Id: hal-01606295

https://hal.science/hal-01606295

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Optimization of control strategies of plant epidemics through a simulation model validated by experiments

Loup Rimbaud¹, Emmanuel Jacquot², Samuel Soubeyrand³, Gaël Thébaud²

² INRA, UMR BGPI, Cirad TA A-54/K, F-34398 Montpellier Cedex 5

loup.rimbaud@supagro.inra.fr

Management strategies of epidemics are often based on expert opinions rather than on the formal demonstration that they are, at least in theory, effective. This mainly results from the complexity of taking into account the biological processes and the human interventions that both impact disease spread. A promising approach to overcome these obstacles consists in identifying optimized management strategy through modeling both the epidemic process and the control measures.

This approach, transposable to many epidemic diseases, is applied on sharka, the most damaging disease of stone fruit trees (especially apricot, peach and plum). It is caused by *Plum pox virus* (PPV, genus *Potyvirus*), a quarantine pathogen in the European Union. A spatiotemporal model simulating sharka spread in a heterogeneous landscape has been developed. One key hypothesis of this model is the synchrony between incubation (the time between inoculation and symptom expression) and latency (the time between inoculation and infectiousness of the host) for PPV-infected peach trees. We developed a generic approach to estimate the mismatch between incubation and latency from repeated transmission experiments. Then, sensitivity analyses are performed in order to identify key parameters of both the epidemic process and the control strategy.

Key words: infectious potential, latent period, *Potyviridae*, SEIR, stochastic spatiotemporal simulation model

¹ Montpellier SupAgro, UMR BGPI, Cirad TA A-54/K, F-34398 Montpellier Cedex 5

³ INRA, UR 546 Biostatistics and Spatial Processes, F-84914 Avignon Cedex 9