



French genomic experience: genomics for all ruminant species

Eric Venot, Didier Boichard, Vincent Ducrocq, Pascal Croiseau, S. Fritz, A. Baur, R. Saintilan, Thierry Tribout, Marie-Pierre Sanchez, Pascal Boulesteix, et al.

► To cite this version:

Eric Venot, Didier Boichard, Vincent Ducrocq, Pascal Croiseau, S. Fritz, et al.. French genomic experience: genomics for all ruminant species. 40. Biennal session of ICAR, Oct 2016, Puerto Varas, Chile. hal-01606324

HAL Id: hal-01606324

<https://hal.science/hal-01606324>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License




French genomic experience: genomic for all ruminant species

E. Venot¹, D. Boichard¹, V. Ducrocq¹, P. Croiseau¹, S. Fritz³,
A. Baur³, R. Saintilan³, T. Tribout¹, M-P Sanchez¹,
P. Boulesteix², JM. Astruc², A. Legarra¹, C. Robert-Granié¹,
C. Carillier¹, I. Palhière¹, F. Tortereau¹, R. Rupp¹,
H. Larroque¹, V. Loywyck², S. Mattalia²

¹ INRA, ² IDELE, ³ ALLICE



French specificities




Source: www.CartesFrance.fr

French specificities



Large number of breeds



Small ruminant breeds





France
Génétique
Elevage

French national and collective organization

- ▶ Inter-professional association for genetic improvement of all ruminants
- ▶ Common management quality system for all on-field organizations
- ▶ **Centralized national database**
- ▶ Close relationship between Industry partners and R & D
 - ▶ **Win – Win situation**
- ▶ Share of R & D **between species**
- ▶ Example with the share of tools:
 - in 2016 a common pipeline to impute & phase Dairy & Beef genomic information from different chips
 - implementation of dairy sheep in the same pipeline in progress



French genomic experience:
genomics for all ruminant species
27th October 2016



French genomic developments led by Dairy cattle pioneers

IDEAL SITUATION

- ▶ large number of tested bulls and large use of AI
- ▶ large generation interval
- ▶ high cost of progeny testing
- ▶ small genetic size populations
- ▶ relative small genotyping cost / breeding animal price
- ▶ international breeds (Interbull evaluation and possible consortia)



ALL TOOLS TO IMPROVE:

$$\Delta G = \frac{i.R.\sigma_g}{T}$$

Reference population size & Accuracy (R)

Reference population per breed and reliability expected for candidates

| Breed | Nb Bulls | Nb Dams | Young candidate REL |
|--------------|----------|---------|---------------------|
| HOLSTEIN | 33000 | - | [0.55-0.70] |
| BROWN | 6000 | - | [0.45-0.70] |
| MONTBELIARDE | 2800 | 31000 | [0.55-0.70] |
| NORMANDE | 2400 | 16000 | [0.50-0.65] |

intergenomics

December 2015



Reference population size & Accuracy (R)

Reference population per breed and reliability expected for candidates


| Breed | Nb Bulls | Nb Dams | Young candidate REL |
|--------------|----------|---------|---------------------|
| HOLSTEIN | 33000 | - | [0.55-0.70] |
| BROWN | 6000 | - | [0.45-0.70] |
| MONTBELIARDE | 2800 | 31000 | [0.55-0.70] |
| NORMANDE | 2400 | 16000 | [0.50-0.65] |



December 2015


Genomic selection started in 2009 (QTL-GS)
6 times / year with QTL and marker effects estimation
 +
Weekly (without QTL effect estimation)

Reduction of generation interval and more efficient selection on functional traits




Quick adoption of GS by breeders: use of young genomic bulls

AI % in France




Quick adoption of GS by breeders: more balanced selection

Genetic profile of the AI bulls used /year Prim Holstein breed




Large increase of genotyping



Larger panel of bulls used

% of AI done with the 5 most used bulls each year Montbéliarde breed



Implementation in other breeds and species?

- ▶ Less favorable situation for Beef cattle breeds
and Dairy regional breeds
 - genetic population size larger
 - reference population more difficult to generate
 - smaller population size
 - less AI for beef breeds
 - phenotype less precise / DYD



Multi-breed option?

Multi-breed genomic evaluation for cattle?

HD genotyping of bulls from 18 cattle breeds in selection in France



Unsuccessful

=> very few common QTL found between breeds

=> no multi-breed genomic evaluation possible



Good basis

to develop genomic selection for several breeds

Genomic selection for regional dairy cattle breeds

Reference population per breed and reliability expected for candidates

| Breed | Nb Bulls | Nb Dams | Young candidate REL |
|------------|----------|---------|---------------------|
| ABONDANCE | 350 | 1900 | [0.35-0.55] |
| TARENTAISE | 300 | 1300 | [0.30-0.50] |
| SIMMENTAL | 300 | 200 | [0.20-0.50] |
| VOSGIENNE | 60 | 1100 | [0.20-0.50] |



December 2015



Official in 2015

Reference population for beef cattle breeds

Maternal effects

| | | Charolais | Limousine | Blonde |
|-------------------------|---------|---------------|---------------|---------------|
| Direct genetic effect | Birth | 15 000 / 1800 | 6 600 / 2 700 | 5 600 / 1 200 |
| | Weaning | 12 500 / 2400 | 5 600 / 2 400 | 3 900 / 1 000 |
| | Carcass | 2 400 / 600 | 1 750 / 400 | 720 / 300 |
| Maternal genetic effect | Birth | 5 200 / 500 | 3 300 / 200 | 3 000 / 200 |
| | Weaning | 4 500 / 450 | 2 400 / 500 | 2 100 / 200 |



Blending of polygenic and genomic results (VanRaden et al., 2009)



Example of reliability gain for young Charolais calf

| EBV | Reliability gain |
|----------------------------|------------------|
| Birth | + 0.10 |
| Growth Total Merit Index | + 0.06 |
| Maternal Total Merit Index | + 0.09 |

Reference population for beef cattle breeds

Maternal effects

| | | Charolais | Limousine | Blonde |
|-------------------------|---------|---------------|---------------|---------------|
| Direct genetic effect | Birth | 15 000 / 1800 | 6 600 / 2 700 | 5 600 / 1 200 |
| | Weaning | 12 500 / 2400 | 5 600 / 2 400 | 3 900 / 1 000 |
| | Carcass | 2 400 / 600 | 1 750 / 400 | 720 / 300 |
| Maternal genetic effect | Birth | 5 200 / 500 | 3 300 / 200 | 3 000 / 200 |
| | Weaning | 4 500 / 450 | 2 400 / 500 | 2 100 / 200 |



Blending of polygenic and genomic results (VanRaden et al., 2009)



Official since end 2015 for BLA, CHA and LIM

2 national runs / year

+

Weekly computation for new candidates
(without re-estimation of marker effects)

What about Small ruminants?

► Less favorable situation for small ruminants

- cost of genotyping relatively high
- already short generation interval
- use of fresh semen for AI in sheep
- AI demand highly concentrated in time



► Different cases:

| | Number of milk recorded ewes | Genotyped rams with daughters |
|-----------------|------------------------------|-------------------------------|
| Lacaune | 156 491 | 4 716 |
| Manech TR | 66 020 | 1 879 |
| Basco-Béarnaise | 21 620 | 562 |
| Manech TN | 9 302 | 475 |
| Corse | 14 028 | 197 |

(2015)

Results for dairy sheep

- QTL detection studies: no big QTL found (limited power detection)
- ssGBLUP with phantom group


=> reliability gain: [0.13 ; 0.20]

- Lacaune experimental study:

- different genomic selection pressures on young rams modeled


=> with 1/3 selection pressure, + 15 % genetic gain without extra cost

=> cost of genotyping balanced by reduction of number of rams in AI Center.



► GS started in 2015 (7 GS /year) + end of progeny testing

Candidate selection
(born in 2011)



Comparison of Total Merit Index after progeny testing in 2013

Progeny testing


$$TMI_{\text{genomic}} - TMI_{\text{classic}} = 0.5 \sigma_{TMI}$$

Meat sheep

- ▶ limited or no progeny testing and limited AI => No GS planned
- ▶ Genotyping first for parentage verification and assignation
- ▶ Research to include major gene in genetic evaluation (prolificacy...)

Goats

- ▶ Goat chip available in 2010
- ▶ Research in progress on:
 - combination of the 2 breeds (Alpine and Saanen)
 - international collaboration
 - major gene inclusion in genetic evaluation



Goal: start of Genomic Selection in 2017

CONCLUSION

- ▶ It has been a **real genomic revolution** these last years
- ▶ **large impact** on selection scheme organization
=> early use of breeding animals for higher genetic gain on all traits
- ▶ thanks to the French collective and efficient organization with Industry partners closely linked to R & D, genomic selection have been adapted to the different specificities of ruminant species and breeds



Genomic evaluation in place in 14 cattle and 1 sheep breeds.

In the very next future for 4 sheep and 2 goat breeds

END OF THE STORY? ...

... NO! JUST THE BEGINNING!

► several challenges to come:

- Organizational reforms
- Reference population maintenance
- Competition for new phenotypes
- Integration of sequence information

...



