



Tick-Pathogen ensembles: Do molecular interactions lead ecological innovation?

Alejandro Cabezas Cruz, Agustin Estrada-Pena, Ryan O. M. Rego, Jose de La Fuente

► To cite this version:

Alejandro Cabezas Cruz, Agustin Estrada-Pena, Ryan O. M. Rego, Jose de La Fuente. Tick-Pathogen ensembles: Do molecular interactions lead ecological innovation?. *Frontiers in Cellular and Infection Microbiology*, 2017, 7, 10.3389/fcimb.2017.00074 . hal-01607006

HAL Id: hal-01607006

<https://hal.science/hal-01607006>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Tick-Pathogen Ensembles: Do Molecular Interactions Lead Ecological Innovation?

Alejandro Cabezas-Cruz^{1,2,3*}, Agustín Estrada-Peña⁴, Ryan O. M. Rego^{2,3} and José De la Fuente^{5,6}

¹ UMR BIPAR, Animal Health Laboratory, ANSES, Institut National de la Recherche Agronomique, ENVA, Maisons Alfort, France, ² Department of Parasitology, Faculty of Science, University of South Bohemia, České Budějovice, Czechia, ³ Biology Center, Institute of Parasitology, Czech Academy of Sciences, České Budějovice, Czechia, ⁴ Faculty of Veterinary Medicine, University of Zaragoza, Zaragoza, Spain, ⁵ SaBio. Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-UCLM-JCCM), Ciudad Real, Spain, ⁶ Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State University, Stillwater, OK, USA

Keywords: tick-pathogen interactions, transcriptional reprogramming, epigenetics, ecological adaptation, *Anaplasma phagocytophilum*

OPEN ACCESS

Edited by:

Yasuko Rikihisa,
Ohio State University at Columbus,
USA

Reviewed by:

Jason A. Carlyon,
Virginia Commonwealth University
School of Medicine, USA
Girish Neelakanta,
Old Dominion University, USA

*Correspondence:

Alejandro Cabezas-Cruz
cabezasalejandrocruz@gmail.com

Received: 16 November 2016

Accepted: 27 February 2017

Published: 13 March 2017

Citation:

Cabezas-Cruz A, Estrada-Peña A,
Rego ROM and De la Fuente J (2017)
Tick-Pathogen Ensembles: Do
Molecular Interactions Lead Ecological
Innovation?
Front. Cell. Infect. Microbiol. 7:74.
doi: 10.3389/fcimb.2017.00074

Ticks are arthropods distributed worldwide that constitute the most important vectors of diseases to animals, and second to mosquitoes regarding pathogens of public health importance. Ticks are remarkably plastic and can colonize diverse ecological niches of the planet, from tropics to polar areas (de la Fuente et al., 2008). In the last decade, the reports of tick-borne pathogens have increased sharply, motivating vigorous research programs that addressed major questions on the epidemiology of tick-borne diseases, vector-host-pathogen interactions, tick ecology, and tick genomics. Notably, the first tick genome was released this year (Gulia-Nuss et al., 2016), opening new possibilities to explore tick-host-pathogen interactions (de la Fuente et al., 2016a). In contrast, the evolutionary and ecological implications of tick-pathogen associations have received comparatively less attention. Herein, we hypothesized that tick-pathogen associations evolved to form “intimate epigenetic relationships” similar to those described for *Theileria* spp. and its vertebrate host (Cheeseman and Weitzman, 2015) in which the pathogen induces transcriptional reprogramming in infected ticks. This will ultimately favor pathogen propagation, but will also select for the most suitable ecological adaptations in the tick vector. These phenotypic and genetic changes may have the potential to be transmitted to the next generation of ticks. As a result, the ecological associations between tick, vertebrates, and pathogens would evolve to maximize pathogen circulation in these communities (Estrada-Peña et al., 2015, 2016).

Our hypothesis was based on the following evidences: (i) tick-borne pathogens induce transcriptional reprogramming in infected tick (Ayllón et al., 2015; Villar et al., 2015; Weisheit et al., 2015) and vertebrate cells (Lee et al., 2008; Bouquet et al., 2016); (ii) tick-borne pathogens produce and secrete effector proteins, nucleomodulins, which constitute a family of proteins produced by bacterial pathogens to control host transcription and other nuclear processes (Bierne and Cossart, 2012), that interact with host epigenetic machinery and induce transcriptional reprogramming (Garcia-Garcia et al., 2009a,b; Rennoll-Bankert et al., 2015; Sinclair et al., 2015; Lina et al., 2016), and (iii) tick-pathogen interactions increase tick fitness (Neelakanta et al., 2010; Belova et al., 2012; Herrmann and Gern, 2015; de la Fuente et al., 2016b).

TICK-BORNE PATHOGENS INDUCE TRANSCRIPTIONAL REPROGRAMMING IN HOST CELLS

Several studies using “omics” technologies have revealed that a common pattern in the infection by tick-borne pathogens is the transcriptional reprogramming of the host cells. These pathogens include obligate intracellular bacterial such as *Anaplasma phagocytophilum* (Carlyon et al., 2002; Borjesson et al., 2005; Pedra et al., 2005; Sukumaran et al., 2005; Lee et al., 2008; Ayllón et al., 2015) and *Ehrlichia chaffeensis* (Miura and Rikihisa, 2009), the extracellular bacterial pathogen *Borrelia burgdorferi* (Bouquet et al., 2016) and viruses such as TBEV (Weisheit et al., 2015). This transcriptional reprogramming not only affect gene expression but also impact protein abundance (Lin et al., 2011; Ayllón et al., 2015). Among the cellular components and processes affected in ticks by pathogen infection are the cytoskeleton, cell immunity, apoptosis, metabolism, and potentially the posttranslational modification of histone tails (Ayllón et al., 2015; Villar et al., 2015; Cabezas-Cruz et al., 2016). Notably, gene expression regulation by tick-borne pathogens occurs in a tissue-specific manner. For example, to establish an infection in ticks, *A. phagocytophilum* inhibits the apoptosis in infected midgut and salivary glands. However, in tick midgut, *A. phagocytophilum* inhibits the apoptosis by upregulating the Janus kinase (JAK)-signaling transducer activator of transcription (JAK-STAT) pathway, but in salivary glands this bacterium down-regulates the expression of porin, which results in the inhibition of cytochrome c release and the intrinsic apoptosis pathway (Ayllón et al., 2015; Alberdi et al., 2016). Taken together, these findings reveal that during evolution tick-borne pathogens have developed specific mechanisms to manipulate gene expression in host cells.

MOLECULAR MESSENGERS OF PATHOGEN MANIPULATION

To manipulate gene expression, pathogens activate signaling pathways or hijack the epigenetic machinery of host cells. Both mechanisms have been described during *A. phagocytophilum* infection in ticks. For example, *A. phagocytophilum* infection triggers expression of antimicrobial peptides in salivary glands that control bacterial load. The expression of this family of antimicrobial peptides is mediated by the activation of the JAK-STAT pathway (Liu et al., 2012). It has also been shown that *A. phagocytophilum* induces the activation of the PI3K signaling pathway leading to actin phosphorylation to increase the expression of the gene *salp16* coding for a tick salivary protein crucial for *A. phagocytophilum* survival (Sultana et al., 2010). However, while signaling pathways activation can explain the regulation of some genes, (Sultana et al., 2010; Liu et al., 2012), it does not explain the massive gene regulation induced by *A. phagocytophilum* infection in ticks (Ayllón et al., 2015). In fact, *A. phagocytophilum* induces the differential expression of 8,516 (from 16,083 gene transcripts identified), 5,394 (12,651) and 2,487 (11,105) genes in *Ixodes scapularis* tick nymphs, adult

midguts, and salivary gland, respectively (Ayllón et al., 2015; de la Fuente et al., 2016a).

A. phagocytophilum produces a family of proteins called nucleomodulins that control host gene expression at the epigenetic level (Sinclair et al., 2015). In particular, the ankyrin repeat effector protein ankyrin A (AnkA) was reported to be secreted by *A. phagocytophilum* through the bacterial type IV secretion system (T4SS) in infected neutrophils (Garcia-Garcia et al., 2009a,b; Rennoll-Bankert et al., 2015). AnkA enters the granulocyte nucleus, binds stretches of AT-rich DNA and alters transcription of antimicrobial defense genes, including down-regulation of *CYBB*, which codes for a NADPH oxidase 2 (Nox2). This enzyme is involved in the production of reactive oxygen species (ROS), which is crucial in the neutrophil immune response against intracellular bacteria. To achieve this regulatory process, AnkA recruits host histone deacetylase 1 (HDAC1) and decreases histone H3 acetylation in infected cells (Garcia-Garcia et al., 2009a,b). This results in chromatin changes that down-regulate the expression of target genes (e.g., *CYBB*). Remarkably, 50 proteins were identified in the genome of *A. phagocytophilum* that may have a function similar to that of AnkA (Sinclair et al., 2015). In addition, genome wide evidence showed that AnkA not only binds to *CYBB* promoter regions, but broadly throughout all chromosomes and correlates with infection-induced differential gene expression (Dumler et al., 2016). Whether *A. phagocytophilum* AnkA is expressed during tick infection is not known. However, it was recently shown that *I. scapularis* has a homolog of the HDAC1 protein that is over-represented in salivary glands in response to *A. phagocytophilum* infection (Cabezas-Cruz et al., 2016). In addition, pharmacological inhibition of tick HDAC1 reduced the load of *A. phagocytophilum* in ISE6 tick cells (Cabezas-Cruz et al., 2016). This result suggests that *A. phagocytophilum* uses similar strategies to manipulate tick and vertebrate host cells (de la Fuente et al., 2016c). The role of *A. phagocytophilum* nucleomodulins during infection provides the molecular basis for specific and genome wide manipulation of host gene expression.

TICK-PATHOGEN INTERACTIONS INCREASE TICK FITNESS

Pathogens must overcome many barriers in order to establish an infection in the tick. Increasing tick fitness by pathogen infection so as to survive would be a *win-win strategy* (de la Fuente et al., 2016b). There are remarkable examples in which pathogens manipulate tick protective responses to facilitate infection but preserving tick feeding and vector capacity to guarantee the survival of both the pathogens and ticks. For example, Neelakanta et al. (2010) demonstrated that *I. scapularis* ticks infected with *A. phagocytophilum* show enhanced fitness against freezing injury due to the induced expression of a tick antifreeze glycoprotein. They further showed that improved survival of infected ticks correlated with higher bacterial infection, therefore providing a direct link between pathogen infection and tick fitness in unfavorable ecological conditions. *A. phagocytophilum* may also affect tick questing behavior by increasing the levels of Heat

Shock Proteins (HSP), which also prevent blood-feeding stress and desiccation at high temperatures (Busby et al., 2012; Villar et al., 2015). Tick questing behavior is essential to find new hosts and survive in nature. Similarly, *A. phagocytophilum* does not manipulate the levels of Subolesin, a protein involved in the tick innate immune response, because it affects infection, tick feeding, and reproduction (de la Fuente et al., 2016b). In contrast, Porin levels are down-regulated by *A. phagocytophilum* infection as a mechanism to inhibit apoptosis, but without affecting tick fitness (Ayllón et al., 2015; Alberdi et al., 2016; de la Fuente et al., 2016b). These results support that *A. phagocytophilum*-induced transcriptional reprogramming selectively manipulates the expression of tick genes that increase tick fitness and therefore pathogen circulation.

Although similar molecular mechanisms have not been described for *Borrelia* spp. and TBEV infections, they also appear to increase tick fitness (Herrmann and Gern, 2015). *Borrelia* and TBEV-infected *I. scapularis* and *I. persulcatus* ticks were found at higher questing heights when compared to uninfected ticks. Higher questing height increases the chances of a tick to find a larger host that could accommodate more ticks increasing their feeding possibilities, but at the same time exposes ticks to more desiccating conditions (Lefcort and Durden, 1996; Romashchenko et al., 2012). Low relative humidity is detrimental for ticks because they spend their energy reserves quicker than at higher relative humidity (Randolph and Storey, 1999). The

fact that *Borrelia* and TBEV-infected ticks choose higher questing height suggests that these pathogens help ticks to survive under dry conditions. In agreement with this hypothesis, *I. ricinus* infected by *B. burgdorferi* move less toward a humid environment and their survival is higher in highly desiccating conditions (Herrmann and Gern, 2010, 2012).

TICK-BORNE PATHOGENS HAVE THE POTENTIAL TO LEAD ECOLOGICAL ADAPTATION DURING TICK EVOLUTION

It was generally assumed that DNA changes are the only way information can be passed from parents to the offspring, and that some phenotypic changes acquired during the life span cannot be transmitted to the following generations. Accumulating evidence, however, indicates that both genetic and epigenetic (defined as changes in gene expression due to processes that arise independent of changes in the underlying DNA sequence) have important effects on evolutionary outcomes (Danchin et al., 2011; Gómez-Díaz et al., 2012). While host physiology manipulation by pathogens is a widely accepted phenomenon, we lack evidence of the heritable character of host phenotypes induced by pathogens (Gómez-Díaz et al., 2012; Poulin and Maure, 2015). It was previously proposed that all trans-generational effects on host offspring phenotype that are induced by parasites must involve




FIGURE 1 | Model of tick ecological adaptation induced by *A. phagocytophilum*. A model is proposed for the general mechanism of tick manipulation by tick-borne pathogens and induction of ecological adaptation. The intracellular bacterium *A. phagocytophilum* (*Ap*) is used as a model. Upon contact with the host membrane or once inside the parasitophorous vacuole, *A. phagocytophilum* secretes nucleomodulins that will enter the tick cell nucleus and recruit histone modifying enzymes (i.e., HDAC1) to modify the expression of target genes. Some of these genes are involved in traits that favor adaptive phenotypes (red ticks) to abiotic factors (e.g., environmental conditions) or biotic factors (e.g., interactions with microorganisms that may be harmful for the ticks). Histone tail modifications (deacetylation/acetylation, methylation/demethylation, etc) resulting from histone modifying enzymes recruitment, will be passed to the next generation. The ticks able to establish this “intimate epigenetic relationships” (Cheeseman and Weitzman, 2015) with the pathogen will have higher fitness compared to the ticks that are not infected (gray ticks). During evolution, this process will lead to tick ecological adaptation and innovation.

a strong epigenetic component (Poulin and Thomas, 2008). Given the demonstrated propensity of tick-borne pathogens to modulate tick gene expression, to do so epigenetically, and the increase in tick fitness that can result, we propose that: pathogen-induced effects on tick phenotype have the potential to be transmitted across generations, therefore accelerating the ecological adaptation of ticks to natural environments (Figure 1). The tick antifreeze glycoprotein triggered by *A. phagocytophilum* infection in *I. scapularis* offers a good model to study this phenomenon (Neelakanta et al., 2010). Natural populations of *I. scapularis* in North America inhabit regions where the temperature reaches freezing conditions for much of the winter (Eisen et al., 2016). Ticks infected by *A. phagocytophilum* will be better adapted to cold temperatures in natural environments. It is reasonable to hypothesize that the up-regulation of the antifreeze glycoprotein expression

is associated with specific histones or DNA modifications. Inheritance of these epigenetic modifications may transmit the cold-survival phenotype to tick offspring. If this phenotype is advantageous, it may be fixed in the tick population even in the absence of the initial stimulus (i.e., *A. phagocytophilum* infection).

AUTHOR CONTRIBUTIONS

All authors listed, have made substantial, direct and intellectual contribution to the work, and approved it for publication.

ACKNOWLEDGMENTS

We thank Professor Kayla King at the University of Oxford for her revision and insightful comments on the present manuscript.

REFERENCES

- Alberdi, P., Espinosa, P. J., Cabezas-Cruz, A., and de la Fuente, J. (2016). *Anaplasma phagocytophilum* manipulates host cell apoptosis by different mechanisms to establish infection. *Vet. Sci.* 3:15. doi: 10.3390/vetsci3030015
- Ayllón, N., Villar, M., Galindo, R. C., Kocan, K. M., Šíma, R., López, J. A., et al. (2015). Systems biology of tissue-specific response to *Anaplasma phagocytophilum* reveals differentiated apoptosis in the tick vector *Ixodes scapularis*. *PLoS Genet.* 11:e1005120. doi: 10.1371/journal.pgen.1005120
- Belova, O. A., Burenkova, L. A., and Karganova, G. G. (2012). Different tick-borne encephalitis virus (TBEV) prevalences in unfed versus partially engorged ixodid ticks - evidence of virus replication and changes in tick behavior. *Ticks Tick Borne Dis.* 3, 240–246. doi: 10.1016/j.ttbdis.2012.05.005
- Bierne, H., and Cossart, P. (2012). When bacteria target the nucleus: the emerging family of nucleomodulins. *Cell. Microbiol.* 14, 622–633. doi: 10.1111/j.1462-5822.2012.01758.x
- Borjesson, D. L., Kobayashi, S. D., Whitney, A. R., Voyich, J. M., Argue, C. M., and Deleo, F. R. (2005). Insights into pathogen immune evasion mechanisms: *Anaplasma phagocytophilum* fails to induce an apoptosis differentiation program in human neutrophils. *J. Immunol.* 174, 6364–6372. doi: 10.4049/jimmunol.174.10.6364
- Bouquet, J., Soloski, M. J., Swei, A., Cheadle, C., Federman, S., Billaud, J. N., et al. (2016). Longitudinal transcriptome analysis reveals a sustained differential gene expression signature in patients treated for acute lyme disease. *MBio* 7, e00100–e00116. doi: 10.1128/mBio.00100-16
- Busby, A. T., Ayllón, N., Kocan, K. M., Blouin, E. F., de la Fuente, G., Galindo, R. C., et al. (2012). Expression of heat shock proteins and subolesin affects stress responses, *Anaplasma phagocytophilum* infection and questing behaviour in the tick, *Ixodes scapularis*. *Med. Vet. Entomol.* 26, 92–102. doi: 10.1111/j.1365-2915.2011.00973.x
- Cabezas-Cruz, A., Alberdi, P., Ayllón, N., Valdés, J. J., Pierce, R., Villar, M., et al. (2016). *Anaplasma phagocytophilum* increases the levels of histone modifying enzymes to inhibit cell apoptosis and facilitate pathogen infection in the tick vector *Ixodes scapularis*. *Epigenetics* 11, 303–319. doi: 10.1080/15592294.2016.1163460
- Carlyon, J. A., Chan, W. T., Galán, J., Roos, D., and Fikrig, E. (2002). Repression of *rac2* mRNA expression by *Anaplasma phagocytophilum* is essential to the inhibition of superoxide production and bacterial proliferation. *J. Immunol.* 169, 7009–7018. doi: 10.4049/jimmunol.169.12.7009
- Cheeseman, K., and Weitzman, J. B. (2015). Host-parasite interactions: an intimate epigenetic relationship. *Cell. Microbiol.* 17, 1121–1132. doi: 10.1111/cmi.12471
- Danchin, É., Charmantier, A., Champagne, F. A., Mesoudi, A., Pujol, B., and Blanchet, S. (2011). Beyond DNA: integrating inclusive inheritance into an extended theory of evolution. *Nat. Rev. Genet.* 12, 475–486. doi: 10.1038/nrg3028
- de la Fuente, J., Estrada-Peña, A., Cabezas-Cruz, A., and Kocan, K. M. (2016c). *Anaplasma phagocytophilum* uses common strategies for infection of ticks and vertebrate hosts. *Trends Microbiol.* 24, 173–180. doi: 10.1016/j.tim.2015.12.001
- de la Fuente, J., Estrada-Peña, A., Venzal, J. M., Kocan, K. M., and Sonenshine, D. E. (2008). Overview: ticks as vectors of pathogens that cause disease in humans and animals. *Front. Biosci.* 13, 6938–6946. doi: 10.2741/3200
- de la Fuente, J., Villar, M., Cabezas-Cruz, A., Estrada-Peña, A., Ayllón, N., and Alberdi, P. (2016b). Tick-host-pathogen interactions: conflict and cooperation. *PLoS Pathog.* 12:e1005488. doi: 10.1371/journal.ppat.1005488
- de la Fuente, J., Waterhouse, R. M., Sonenshine, D. E., Roe, R. M., Ribeiro, J. M., Sattelle, D. B., et al. (2016a). Tick genome assembled: new opportunities for research on tick-host-pathogen interactions. *Front. Cell. Infect. Microbiol.* 6:103. doi: 10.3389/fcimb.2016.00103
- Dumler, J. S., Sinclair, S. H., Pappas-Brown, V., and Shetty, A. C. (2016). Genome-Wide *Anaplasma phagocytophilum* Anka-DNA interactions are enriched in intergenic regions and gene promoters and correlate with infection-induced differential gene expression. *Front. Cell. Infect. Microbiol.* 6:97. doi: 10.3389/fcimb.2016.00097
- Eisen, R. J., Eisen, L., and Beard, C. B. (2016). County-Scale Distribution of *Ixodes scapularis* and *Ixodes pacificus* (Acari: Ixodidae) in the Continental United States. *J. Med. Entomol.* 53, 349–386. doi: 10.1093/jme/tjv237
- Estrada-Peña, A., de la Fuente, J., Ostfeld, R. S., and Cabezas-Cruz, A. (2015). Interactions between tick and transmitted pathogens evolved to minimise competition through nested and coherent networks. *Sci. Rep.* 5:10361. doi: 10.1038/srep10361
- Estrada-Peña, A., Sprong, H., Cabezas-Cruz, A., de la Fuente, J., Ramo, A., and Coipan, E. C. (2016). Nested coevolutionary networks shape the ecological relationships of ticks, hosts, and the Lyme disease bacteria of the *Borrelia burgdorferi* (s.l.) complex. *Parasit. Vectors* 9, 517. doi: 10.1186/s13071-016-1803-z
- García-García, J. C., Barat, N. C., Trembley, S. J., and Dumler, J. S. (2009a). Epigenetic silencing of host cell defense genes enhances intracellular survival of the rickettsial pathogen *Anaplasma phagocytophilum*. *PLoS Pathog.* 5:e1000488. doi: 10.1371/journal.ppat.1000488
- García-García, J. C., Rennoll-Bankert, K. E., Pelly, S., Milstone, A. M., and Dumler, J. S. (2009b). Silencing of host cell CYBB gene expression by the nuclear effector Anka of the intracellular pathogen *Anaplasma phagocytophilum*. *Infect. Immun.* 77, 2385–2391. doi: 10.1128/IAI.00023-09
- Gómez-Díaz, E., Jordà M., Peinado, M. A., and Rivero, A. (2012). Epigenetics of host-pathogen interactions: the road ahead and the road behind. *PLoS Pathog.* 8:e1003007. doi: 10.1371/journal.ppat.1003007
- Gulia-Nuss, M., Nuss, A. B., Meyer, J. M., Sonenshine, D. E., Roe, R. M., Waterhouse, R. M., et al. (2016). Genomic insights into the *Ixodes scapularis* tick vector of Lyme disease. *Nat. Commun.* 7:10507. doi: 10.1038/ncomms10507

- Herrmann, C., and Gern, L. (2010). Survival of *Ixodes ricinus* (Acari: Ixodidae) under challenging conditions of temperature and humidity is influenced by *Borrelia burgdorferi* sensu lato infection. *J. Med. Entomol.* 47, 1196–1204. doi: 10.1603/ME10111
- Herrmann, C., and Gern, L. (2012). Do the level of energy reserves, hydration status and *Borrelia* infection influence walking by *Ixodes ricinus* (Acari: Ixodidae) ticks? *Parasitology* 139, 330–337. doi: 10.1017/S0031182011002095
- Herrmann, C., and Gern, L. (2015). Search for blood or water is influenced by *Borrelia burgdorferi* in *Ixodes ricinus*. *Parasit. Vectors* 8:6. doi: 10.1186/s13071-014-0526-2
- Lee, H. C., Kioi, M., Han, J., Puri, R. K., and Goodman, J. L. (2008). *Anaplasma phagocytophilum*-induced gene expression in both human neutrophils and HL-60 cells. *Genomics* 92, 144–151. doi: 10.1016/j.ygeno.2008.05.005
- Lefcort, H., and Durden, L. A. (1996). The effect of infection with Lyme disease spirochetes (*Borrelia burgdorferi*) on the phototaxis, activity, and questing height of the tick vector *Ixodes scapularis*. *Parasitology* 113, 97–103.
- Lin, M., Kikuchi, T., Brewer, H. M., Norbeck, A. D., and Rikihisa, Y. (2011). Global proteomic analysis of two tick-borne emerging zoonotic agents: *Anaplasma phagocytophilum* and *Ehrlichia chaffeensis*. *Front. Microbiol.* 2:24. doi: 10.3389/fmicb.2011.00024
- Lina, T. T., Farris, T., Luo, T., Mitra, S., Zhu, B., and McBride, J. W. (2016). Hacker within! *Ehrlichia chaffeensis* effector driven phagocyte reprogramming strategy. *Front. Cell. Infect. Microbiol.* 6:58. doi: 10.3389/fcimb.2016.00058
- Liu, L., Dai, J., Zhao, Y. O., Narasimhan, S., Yang, Y., Zhang, L., et al. (2012). *Ixodes scapularis* JAK-STAT pathway regulates tick antimicrobial peptides, thereby controlling the agent of human granulocytic anaplasmosis. *J. Infect. Dis.* 206, 1233–1241. doi: 10.1093/infdis/jis484
- Miura, K., and Rikihisa, Y. (2009). Liver transcriptome profiles associated with strain-specific *Ehrlichia chaffeensis*-induced hepatitis in SCID mice. *Infect. Immun.* 77, 245–254. doi: 10.1128/IAI.00979-08
- Neelakanta, G., Sultana, H., Fish, D., Anderson, J. F., and Fikrig, E. (2010). *Anaplasma phagocytophilum* induces *Ixodes scapularis* ticks to express an antifreeze glycoprotein gene that enhances their survival in the cold. *J. Clin. Invest.* 120, 3179–3190. doi: 10.1172/JCI42868
- Pedra, J. H., Sukumaran, B., Carlyon, J. A., Berliner, N., and Fikrig, E. (2005). Modulation of NB4 promyelocytic leukemic cell machinery by *Anaplasma phagocytophilum*. *Genomics* 86, 365–377. doi: 10.1016/j.ygeno.2005.05.008
- Poulin, R., and Maure, F. (2015). Host manipulation by parasites: a look back before moving forward. *Trends. Parasitol.* 31, 563–570. doi: 10.1016/j.pt.2015.07.002
- Poulin, R., and Thomas, F. (2008). Epigenetic effects of infection on the phenotype of host offspring: parasites reaching across host generations. *Oikos* 117, 331–335. doi: 10.1111/j.2007.0030-1299.16435.x
- Randolph, S. E., and Storey, K. (1999). Impact of microclimate on immature tick-rodent host interactions (Acari: Ixodidae): implications for parasite transmission. *J. Med. Entomol.* 36, 741–748.
- Rennoll-Bankert, K. E., Garcia-Garcia, J. C., Sinclair, S. H., and Dumler, J. S. (2015). Chromatin-bound bacterial effector ankyrin A recruits histone deacetylase 1 and modifies host gene expression. *Cell. Microbiol.* 17, 1640–1652. doi: 10.1111/cmi.12461
- Romashchenko, A. V., Ratushnyak, A. S., Zapara, T. A., Tkachev, S. E., and Moshkin, M. P. (2012). The correlation between tick (*Ixodes persulcatus* Sch.) questing behaviour and synganglion neuronal responses to odours. *J. Insect. Physiol.* 58, 903–910. doi: 10.1016/j.jinsphys.2012.04.004
- Sinclair, S. H., Garcia-Garcia, J. C., and Dumler, J. S. (2015). Bioinformatic and mass spectrometry identification of *Anaplasma phagocytophilum* proteins translocated into host cell nuclei. *Front. Microbiol.* 6:55. doi: 10.3389/fmicb.2015.00055
- Sukumaran, B., Carlyon, J. A., Cai, J. L., Berliner, N., and Fikrig, E. (2005). Early transcriptional response of human neutrophils to *Anaplasma phagocytophilum* infection. *Infect. Immun.* 73, 8089–8099. doi: 10.1128/IAI.73.12.8089-8099.2005
- Sultana, H., Neelakanta, G., Kantor, F. S., Malawista, S. E., Fish, D., Montgomery, R. R., et al. (2010). *Anaplasma phagocytophilum* induces actin phosphorylation to selectively regulate gene transcription in *Ixodes scapularis* ticks. *J. Exp. Med.* 207, 1727–1743. doi: 10.1084/jem.20100276
- Villar, M., Ayllón, N., Alberdi, P., Moreno, A., Moreno, M., Tobes, R., et al. (2015). Integrated metabolomics, transcriptomics and proteomics identifies metabolic pathways affected by *Anaplasma phagocytophilum* infection in tick cells. *Mol. Cell. Proteomics* 14, 3154–3172. doi: 10.1074/mcp.M115.051938
- Weisheit, S., Villar, M., Tykalová, H., Popara, M., Loecherbach, J., Watson, M., et al. (2015). *Ixodes scapularis* and *Ixodes ricinus* tick cell lines respond to infection with tick-borne encephalitis virus: transcriptomic and proteomic analysis. *Parasit. Vectors* 8:599. doi: 10.1186/s13071-015-1210-x

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Cabezas-Cruz, Estrada-Peña, Rego and De la Fuente. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.