



HAL
open science

Stress proteins in cull cows: relationships with transport and lairage durations but not with meat tenderness

Mohammed Gagaoua, Claudia Terlouw, Valérie Monteils, Sebastien Couvreur, Brigitte Picard

► To cite this version:

Mohammed Gagaoua, Claudia Terlouw, Valérie Monteils, Sebastien Couvreur, Brigitte Picard. Stress proteins in cull cows: relationships with transport and lairage durations but not with meat tenderness. 63. International Congress of Meat Science and Technology (ICoMST), Aug 2017, Cork, Ireland. 10.3921/978-90-8686-860-5 . hal-01607294

HAL Id: hal-01607294

<https://hal.science/hal-01607294v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

2017
CORK IRELAND
ICoMST
63rd INTERNATIONAL CONGRESS OF MEAT SCIENCE & TECHNOLOGY

Nurturing Locally, Growing Globally



63rd INTERNATIONAL CONGRESS OF
MEAT SCIENCE AND TECHNOLOGY



edited by:
Declan Troy
Ciara M^cDonnell
Laura Hinds
Joseph Kerry

63rd International Congress of Meat Science and Technology

63rd

International Congress of

Meat Science and **Technology**

NURTURING LOCALLY, GROWING GLOBALLY

edited by:

Declan Troy
Ciara McDonnell
Laura Hinds
Joseph Kerry



Wageningen Academic
P u b l i s h e r s

EAN: 9789086863136
e-EAN: 9789086868605
ISBN: 978-90-8686-313-6
e-ISBN: 978-90-8686-860-5
DOI: 10.3921/978-90-8686-860-5

First published, 2017

© Wageningen Academic Publishers
The Netherlands, 2017

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned. Nothing from this publication may be translated, reproduced, stored in a computerised system or published in any form or in any manner, including electronic, mechanical, reprographic or photographic, without prior written permission from the publisher, Wageningen Academic Publishers, P.O. Box 220, 6700 AE Wageningen, the Netherlands,
www.WageningenAcademic.com
copyright@WageningenAcademic.com

The individual contributions in this publication and any liabilities arising from them remain the responsibility of the authors.

The publisher is not responsible for possible damages, which could be a result of content derived from this publication.

STRESS PROTEINS IN CULL COWS: RELATIONSHIP WITH TRANSPORT AND LAIRAGE DURATIONS BUT NOT WITH MEAT TENDERNESS

M. Gagaoua^{1*}, E.M.C. Terlouw, V. Monteils¹, S. Couvreur² and B. Picard¹

¹UMR1213 Herbivores, INRA, VetAgro Sup, Clermont universit , Universit  de Lyon, 63122 Saint-Gen s-Champanelle, France; ²URSE, Universit  Bretagne Loire, Ecole Sup rieure d'Agricultures (ESA), 55 rue Rabelais, BP 30748, 49007 Angers Cedex, France; mohammed.gagaoua@inra.fr

Abstract – This work used *K*-means ($k=3$) after PCA analysis on 109 PDO Maine-Anjou cows according to the abundances of small and large heat shock proteins (HSPs) in the *Longissimus thoracis* muscle. Classes LP ($n=24$) and SP ($n=33$) were characterized by greater abundances of large and small HSPs, respectively, while class IP ($n=52$) was intermediate. The classes were compared for the abundance of other proteins in the same muscle and for meat tenderness. The effects of pre-slaughter factors (transport and lairage durations) were also investigated. Meat tenderness did not differ for any comparison. Certain proteins were influenced by pre-slaughter conditions. Among them, H2AFX and μ -calpain were impacted by both factors. The possible role of H2AFX (H2A histone family, member) in the balance of various processes involved in meat tenderization is discussed.

Key Words – muscle proteome, HSPs proteins, PDO Maine-Anjou cows, meat quality, pre-slaughter handling

INTRODUCTION

HSPs are chaperones expressed constitutively or inductively, which play an important role in regulating cellular homeostasis and promoting cell survival [1]. The abundances of HSPs are believed to play an important role in muscle to meat conversion due to their protective function on structural proteins and their anti-apoptotic properties [2]. However, there are few studies exploring the relationship between HSPs and meat tenderness and the relationships found vary across studies. This study compared three classes of animals varying in their abundances of HSPs and for their abundances of other muscle proteins and tenderness. The effects of pre-slaughter handling (transport and lairage times) on muscle proteome were also investigated.

MATERIALS AND METHODS

109 French PDO Maine-Anjou cull cows of ~67 months old were slaughtered in a commercial abattoir in compliance with the French welfare regulations. Samples from the *Longissimus thoracis* (LT, oxido-glycolytic) muscle were excised from the right side of the carcass of each animal 24 h after slaughter. They served for the quantification by Dot-Blot [3] of the levels of 20 protein biomarkers of tenderness representative of various biological pathways: *heat shock proteins* (α B-crystallin, HSP20, 27, 70-8, 70-1A, 70-1B, 70-Grp75), *oxidative stress* (DJ-1, Prdx6, SOD1), *energy metabolism* (ENO3, PGM1), *structure* (α -actin, MyBP-H, MyHC-IIx, MyLC-1F), *proteolysis* (μ -calpain, m-calpain), *apoptosis* (TP53) and *transcription* (H2AFX) and for electrophoresis of the proportions of myosin heavy chains I, IIa and IIx [4]. Tenderness was assessed by a trained sensory panel and Warner-Bratzler measurements [4]. Statistical analyses comprised first, principal component analysis (PCA) using the small (α B-crystallin, HSP20, 27) and large (HSP70-8, 70-1A, 70-1B) HSPs. Subsequently, a *K*-means cluster analysis ($k=3$) using the variability explained by the axes with eigenvalues >1.0 , allowed the creation of three classes of animals. ANOVA was used to compare muscle proteins other than HSPs and tenderness between classes. ANCOVA was used to study the effect of transport and lairage times on all the proteins. Correlation analyses were performed between proteins to construct a robust correlation network (i.e. correlations present within and across the three classes cf. [3]).

RESULTS AND DISCUSSION

The results show that class LP was characterized by an overabundance of large HSPs and low abundance of small HSPs, while class SP was characterized by the opposite. Class IP had relatively low abundances of both small and large HSPs (Figure 1A and Table 1). These differences may be related to differences in muscle characteristics, as class LP had higher




Figure 1. Loading and score plots (A) of the variables used to discriminate between the animals. The classes LP, SP and IP (1-3) are illustrated by blue, green and red circles, respectively. Robust correlation network (B) constructed between the proteins studied highlighting H2AFX at the crossroad of the network.

glycolytic properties levels than class SP and IP. In rabbits and pigs, glycolytic muscles were reported to contain greater levels of large Hsp70 than oxidative which contain mainly small Hsp [5, 6]. In cattle, no studies have reported this before. No differences were observed between sensory tenderness scores or WBSF values between the three classes. This contrasts with earlier reports and may be related to the multifactorial character of the tenderizing process, which involves not only apoptosis, but also oxidation and proteolysis. Specifically, oxidative (Grp75, Prdx6 and DJ-1) and proteolytic enzymes (μ -calpain) showed different abundances between classes which may have disturbed a straightforward relationship between HSPs and tenderness. Possibly, H2AFX has played a role in the balance between the different processes. This histone binds to DNA and to various enzymes, modulating many cellular pathways [2], and is at a crossroad of the correlation network with 8 connectors (Figure 1B). In addition, H2AFX and μ -calpain were both affected by transport and lairage times. Three other proteins (HSP70-1B, Prdx6, DJ-1) were affected by TT and one (Hsp70-8) by TL (low abundance for long lairage duration).

Table 1. Abundances of muscle proteins (in arbitrary units) in the 3 classes and the effects of transport (TT) and lairage times (LT) (in min) on the abundances.

Variables ¹	C1-LP (n=24)	C2-SP (n=33)	C3-IP (n=52)	SEM	Treatments (P-values) ²		
					Class	TT	LT
α B-crystallin	211 ^b	297^a	185 ^b	7.97	***		
HSP20	160 ^b	208^a	138 ^c	4.32	***		
HSP27	68 ^b	95^a	76 ^b	1.92	***		
HSP40	123 ^b	125 ^b	139 ^a	1.48	***		
HSP70-8	133^a	115 ^b	99 ^c	2.40	***		*
HSP70-1A	143^a	126 ^b	105 ^c	2.50	***		
HSP70-1B	223^a	199 ^b	153 ^c	4.19	***	*	
HSP70-Grp75	151 ^a	157 ^a	133 ^b	2.97	***		
Prdx6	109 ^{ab}	113 ^a	100 ^b	1.67	**	*	
DJ-1	97 ^a	91 ^{ab}	86 ^b	1.24	**	*	
ENO3	147 ^a	155 ^a	137 ^b	3.48	*		
PGM1	119 ^a	98 ^b	96 ^b	2.62	**		
α -actin	162 ^a	112 ^b	111 ^b	3.89	***		
MyBP-H	135 ^a	128 ^a	119 ^b	1.79	***		
MyHC-IIx	101 ^a	91 ^{ab}	85 ^b	2.65	*		
μ -calpain (CAPN1)	167 ^a	156 ^{ab}	142 ^b	3.70	*	*	*
H2AFX	116 ^b	127 ^a	114 ^b	2.14	*	*	*
TP53	97 ^b	107 ^a	95 ^b	1.80	*		
Variables related to animal handling (pre-slaughter conditions)							
Transport time (TT)	700 ^a	512 ^b	510 ^b	56	*		
Lairage time (LT)	328 ^b	631 ^{ab}	921 ^a	105	*		

¹ Only variables that were different are shown.

² *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$.

CONCLUSIONS

This study shows that various proteins are influenced by pre-slaughter stress conditions and suggests that some of these may orient post mortem biochemical processes involved in the tenderizing process.

ACKNOWLEDGEMENTS

The authors would thank 'Pays de Loire Region', the defense Trade Union of PDO Maine-Anjou, Adema and Elivia.

REFERENCES

- Schmitt, E., Gehrman, M., Brunet, M., Multhoff, G., & Garrido, C. (2007). Intracellular and extracellular functions of heat shock proteins: repercussions in cancer therapy. *Journal of leukocyte biology* 81: 15-27.
- Picard, B. & Gagaoua, M. (2017). Chapter 11: Proteomic investigations of beef tenderness. In *Proteomics in Food Science: From Farm to Fork*, Colgrave, M., Ed. Elsevier Science: Netherlands; p 538.
- Gagaoua, M.; Terlou, E. M.; Boudjellal, A. & Picard, B. (2015). Coherent correlation networks among protein biomarkers of beef tenderness: What they reveal. *J Proteomics* 128: 365-74.
- Picard, B.; Gagaoua, M.; et al. (2014). Inverse relationships between biomarkers and beef tenderness according to contractile and metabolic properties of the muscle. *J Agric Food Chem*, 62: 9808-18.
- Xu, Y. J., Jin, M. L., Wang, L. et al. (2009). Differential proteome analysis of porcine skeletal muscles between Meishan and Large White. *J. Anim. Sci.* 87: 2519-2527.
- Neufer, D. & Benjamin I.J. (1996). Differential expression of B-crystallin and Hsp27 in skeletal muscle during continuous contractile activity. Relationship to myogenic regulatory factors. *J. Biol. Chem.* 271: 24089-24095.

M. GAGAOUA^{1*}, E.M.C. TERLOUW¹, V. MONTEILS¹, S. COUVREUR², B. PICARD¹

Background

Heat shock proteins (HSPs) play a pivotal role in muscle to meat conversion due to their

- protective function on structural proteins,
- anti-apoptotic properties,
- protein repair activity after cell damage.

HOWEVER


- few studies explored the relationships between HSPs and meat tenderness;
- those existing showed controversies concerning the relationships and the direction of the associations between HSPs and meat tenderness

Aim


To cluster cows according to the abundances of small and large HSPs in the *Longissimus thoracis*:

- to explore the biological mechanisms and understand better the conversion of muscle into meat,
- to investigate the effect of pre-slaughter conditions (transport and lairage times) on muscle proteome.

Methods



Results



Graph 1: Classes LP (n=24) and SP (n=33) were characterized by greater abundances of **Large** and **Small** HSPs, respectively, while class IP (n=52) was intermediate.

No differences in tenderness between the clusters of Heat Shock Proteins

WHY?

The tenderizing process is multifactorial:

involving not only apoptosis, but also oxidation and proteolysis.

1 Significant abundance differences (Table 1)

- oxidative proteins: Grp75, PRDX6 and DJ-1
- proteolytic enzymes: μ -calpain


This may have disturbed a straightforward relationship between HSPs and tenderness

2 \rightarrow H2AFX

possibly played a role in the balance between the different processes:

- It is at a crossroad of the correlation network with 8 connectors (**Graph 2**)
- It binds to DNA and various enzymes, modulating many cellular pathways (**Graph 3**)

Graph 2: the correlation network built between quantified proteins highlights that H2AFX is at the central crossroad of the network



Graph 3: the functional network of proteins involved in tenderization highlights the central role that H2AFX may play in the regulation of the conversion of muscle into meat




Table 1: abundances of muscle proteins differing according to the three classes (ANOVA), and pre-slaughter conditions (ANCOVA).

Biomarkers	LP (n = 24)	SP (n = 33)	IP (n = 52)	SEM	Class	TT	LT
αB-crystallin	211 ^b	297^a	185 ^b	7.97	***		
HSP20	160 ^b	208^a	138 ^c	4.32	***		
HSP27	68 ^b	95^a	76 ^b	1.92	***		
HSP40	123 ^b	125 ^b	139 ^a	1.48	***		
HSP70-8	133^a	115 ^b	99 ^c	2.40	***		*
HSP70-1A	143^a	126 ^b	105 ^c	2.50	***		
HSP70-1B	223^a	199 ^b	153 ^c	4.19	***		*
HSP70-Grp75	151 ^a	157 ^a	133 ^b	2.97	***		
PRDX6	109 ^{a,b}	113 ^a	100 ^b	1.67	**	*	
DJ-1	97 ^a	91 ^{a,b}	86 ^b	1.24	**	*	
ENO3	147 ^a	155 ^a	137 ^b	3.48	*		
PGM1	119 ^a	98 ^b	96 ^b	2.62	**		
α -actin	162 ^a	112 ^b	111 ^b	3.89	***		
MyBP-H	135 ^a	128 ^a	119 ^b	1.79	***		
MyHC-IIx	101 ^a	91 ^{a,b}	85 ^b	2.65	*		
μ -calpain (CAPN1)	167 ^a	156 ^{a,b}	142 ^b	3.70	*	*	*
H2AFX (histone)	116 ^b	127 ^a	114 ^b	2.14	*	*	*
TP53	97 ^b	107 ^a	95 ^b	1.80	*		

Variables related to pre-slaughter conditions

Variable	LP (n = 24)	SP (n = 33)	IP (n = 52)	SEM	Class	TT	LT
Transport time (TT)	700 ^a	512 ^b	510 ^b	56	*		
Lairage time (LT)	328 ^b	631 ^{a,b}	921 ^a	105	*		

¹Only variables that were different are shown; ² *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$

Pre-slaughter conditions effect:

TT and LT influence H2AFX and μ -calpain; TT influences HSP70-1B, PRDX6 and DJ-1; LT influences HSP70-8.

Conclusion

Various proteins are influenced by pre-slaughter stress conditions and some of these may orient *post-mortem* biochemical processes involved in the determinism of meat tenderness. The possible role of H2AFX (H2A histone family, member) in the regulation of the various processes involved in meat tenderization is of interest and needs further investigations.