



## Impact of UV-C radiations of strawberry plants on its sensitivity to Botrytis cinerea

Marine Forges, Hilarion Vasquez, Florence Charles, Laurent Urban, Yves Lizzi, Marc Bardin, Jawad Aarrouf

### ► To cite this version:

Marine Forges, Hilarion Vasquez, Florence Charles, Laurent Urban, Yves Lizzi, et al.. Impact of UV-C radiations of strawberry plants on its sensitivity to Botrytis cinerea. 10. Conférence de la Société française de phytopathologie - 12. European foundation for plant pathology conference, May 2017, Malo-les-Bains, France. , 2017, Deepen knowledge in plant pathology for innovative agro-ecology. hal-01607498

HAL Id: hal-01607498

<https://hal.science/hal-01607498v1>

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Session 6: Emerging tools for the management of plant diseases: Agroecology and disease management  
Oral presentation

**Impact of UV-C radiations of strawberry plants on its sensitivity to *Botrytis cinerea***

M. Forges<sup>1,2</sup>, H. Vásquez<sup>1</sup>, F. Charles<sup>1</sup>, L. Urban<sup>1</sup>, Y. Lizzi<sup>1</sup>, M. Bardin<sup>2</sup>, J. Aarrouf<sup>1</sup>

1 Unité Mixte de Recherche Qualisud, Laboratoire de Physiologie des fruits et Légumes, Université d'Avignon et des Pays de Vaucluse, 301 Rue Baruch de Spinoza, BP2139 – 84916, Avignon, France

2 Pathologie Végétale, INRA, 84140 Montfavet, France

Increasing the resistance of plants by using physical methods remains poorly studied compared to the use of biological or chemical elicitors. The objective of the study is to test the hypothesis that it is possible to reduce the sensitivity of strawberry plants to *Botrytis cinerea* by the application of non-deleterious dose of UV-C radiation on the plant.

Tests were carried out on three cultivars of strawberry. We first showed that these cultivars have different levels of basal sensitivity to *B. cinerea* on leaves. Various treatments combining different doses of UV-C and frequency of application resulted in the selection of non-deleterious doses for strawberry plants. Measures of chlorophyll fluorescence, leaf color and plant growth revealed that doses comprised between 0.4 and 1.70 kJ/m<sup>2</sup> had not any significant effect on these plant physiological parameters, regardless of the frequency of application and of the cultivar tested.

It also resulted in the identification of hormetic doses that can improve the resistance of leaves against grey mold. UV-C treatments applied on plants at 0.85 and 1.70 kJ/m<sup>2</sup> 4 times every two days, resulted in a systematic decrease in the sensitivity of the cultivar Candiss to the strain Bc1 of *B. cinerea* (leaf protection between 11 and 27%). But the effect observed is different according to the strawberry cultivar and to the strain of *B. cinerea* used. In future experiments, we envisage to test the effect of these treatments on fruit quality and in their post-harvest sensitivity to *B. cinerea*.

**Keywords:** strawberry, *Botrytis cinerea*, UV-C radiations, hormetic dose