

## Multiple screening approach for the selection of efficient biological control agents against Botrytis cinerea on tomato

Yousra Bouaoud, Abdelhamid Foughalia, Claire Troulet, Kamel Aissat, Marc Bardin

## ▶ To cite this version:

Yousra Bouaoud, Abdelhamid Foughalia, Claire Troulet, Kamel Aissat, Marc Bardin. Multiple screening approach for the selection of efficient biological control agents against Botrytis cinerea on tomato. 10. Conférence de la Société française de phytopathologie. 12. European foundation for plant pathology conference, May 2017, Malo-les-Bains, France. , 2017, Deepen knowledge in plant pathology for innovative agro-ecology. hal-01607609

HAL Id: hal-01607609

https://hal.science/hal-01607609

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



10. Conférence de la Société française de phytopathologie, 12. European foundation for plant pathology conference, Malo-les-Bains, FRA (2017/05/29 – 2017/06/02)

## Session 6 - Emerging tools for the management of plant diseases: Agroecology and disease management

**Poster presentation** 

Multiple screening approach for the selection of efficient biological control agents against *Botrytis cinerea* on tomato

Yousra Bouaoud<sup>1,2</sup>, Abdelhamid Foughalia<sup>1</sup>, Claire Troulet<sup>2</sup>, Kamel Aissat<sup>1</sup>, Marc Bardin<sup>2</sup>.

One hundred and twenty one bacterial isolates collected from tomato stems and from soil in the Bejaia region (Algeria) were screened for their potential to control gray mold on tomato. The bacteria were initially tested for their direct *in vitro* effect against two strains of *B. cinerea*. Based on these results, 37 isolates showing *in vitro* inhibition of *B. cinerea* mycelial growth (ranging from 0% to 69% inhibition) were chosen for further analysis. These isolates were identified based on 16S rDNA sequencing. Those identified as potentially pathogens for humans, mammals or plants and able to grow at 37°C were discarded. Twenty five were then selected to evaluate their effectiveness as biological control agents against *B. cinerea* using a tomato plant bioassay. Comparisons of *in vitro* and *in planta* screening methods highlighted the absence of correlation between these methods and confirm that a single screening strategy is not sufficient to select efficient biological control agents.

Among the 25 isolates, 8 exert high and significant antagonistic activity against two strains of B. cinerea. Among them, three bacteria belonging to the Pseudomonas genus were finally selected for their significant and stable efficiency at 3 concentrations ( $10^7$ ,  $10^8$  and  $10^9$ CFU/mL) to reduce the lesion development of B. cinerea on tomato stem (% protection comprised between 60 and 100%), their ability to grow at  $15^\circ$ C and their inability to grow at  $37^\circ$ C. These three isolates are also able to reduce significantly the sporulation of B. cinerea on tomato stem.

This study is a first step in the selection of biological control agents able to protect the tomato against *B. cinerea* and adapted to the greenhouse conditions in Algeria.

Keywords: Botrytis cinerea, tomato, biological control agents, screening, Pseudomonas spp.

<sup>&</sup>lt;sup>1</sup>Microbial Ecology Laboratory. University of Abderrahmane Mira, Bejaia 06000, Algeria.

<sup>&</sup>lt;sup>2</sup>Plant Pathology, INRA, 84140 Monfavet, France.