

Charaterization of new MYB genes potentially regulating anthocyanin synthesis in Vitis vinifera

Camila Gomez, Geneviève Conejero, Veronique Cheynier, Nancy Terrier, Agnes Ageorges

▶ To cite this version:

Camila Gomez, Geneviève Conejero, Veronique Cheynier, Nancy Terrier, Agnes Ageorges. Charaterization of new MYB genes potentially regulating anthocyanin synthesis in Vitis vinifera. 3. International Symposium on MACROWINE, Jun 2010, Torino, Italy. hal-01607640

HAL Id: hal-01607640

https://hal.science/hal-01607640

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





MACROWINE2010

on Macromolecules and Secondary Metabolites of Grapevine and Wines

Torino, Italy, June 16-18 / 2010 - Congress Center Torino Incontra Via Nino Costa 8











BOOK OF ABSTRACTS

9.00 – 9.20 Opening Ceremony

- SESSION I – CHEMISTRY, FUNCTIONALITY AND ANALYSIS OF GRAPE AND WINE MACROMOLECULES AND SECONDARY METABOLITES

Chairperson: Véronique Cheynier, Montpellier, France

9.20 - 10.00 Keynote Lecture

CHEMISTRY AND ANALYSIS OF KEY VOLATILE COMPOUNDS OF WINE AND THEIR PRECURSORS IN GRAPE

Philippe Darriet, University of Bordeaux 2, Villenave d'Ornon, France

10.00 - 10.40 Dalmasso Lecture

WINE OXIDATION: RECENT REVELATIONS

Andrew Waterhouse, University of California, Davis, USA

10.40 - 11.00

Characterization and formation kinetic study of phenolic compounds-volatile thiol adducts by chemical and enzymatic oxidation

Maria Nikolantonaki, Institut des Sciences de la Vigne et du Vin, Villenave d'Ornon, France

11.00 - 11.30 Coffee break

11.30 - 11.50

Preliminary studies on polyphenol assessment by Fourier transform-near infrared spectroscopy (FT-NIR) in berries

Chiara Pagliarani, University of Torino, Torino, Italy

11.50 - 12.10

A metabolomic view of wine micro-oxygenation

Fulvio Mattivi, Fondazione Edmund Mach, San Michele all'Adige, Trento, Italy

12.10 - 12.30

Effects of ionic strength and sulfate upon thermal aggregation of grape chitinases and thaumatin-like proteins in model system

Matteo Marangon, The Australian Wine Research Institute, Glen Osmond, Australia

12.30 - 12.50

Progress in precision viticulture for the assessment of grape anthocyanins at harvest in Australian vineyards

Marine Le Moigne, Centre Universitaire Orsay, Orsay Cedex, France

13.00 - 14.00 Lunch

- **SESSION II** - BERRY DEVELOPMENT AND BIOSYNTHESIS OF GRAPE SECONDARY METABOLITES

Chairperson: Serge Delrot, Bordeaux

14.00 - 14.40 Keynote Lecture

REGULATION OF FLAVONOID BIOSYNTHESIS DURING GRAPE BERRY DEVELOPMENT

Jochen Bogs, University of Heidelberg, Adelaide, Germany

14.40 - 15.00

Grapevine anthocyanin transport involves in vivo vesicle-mediated trafficking and the contribution of anthomate transporters and GST

Agnès Ageorges, INRA - UMR SPO, Montpellier, France

15.00 - 15.20

Cellular localization of leucoanthocyanidin reductase and anthocyanidin reductase in grape skins of Vitis vinifera L. cv. Cabernet-Sauvignon during development

Soizic Lacampagne, Institut des Sciences de la Vigne et du Vin, Villenave d'Ornon, France

15.20 - 15.40

Cloning and characterization of putative anthocyanin acyltransferases from grape and Petunia

Sofia Provenzano, University of Torino, Torino, Italy

15.40 - 16.10 Coffee break

16.10 - 16.30

New MYBS through the glass: a deeper look into the regulatory network of flavonoid synthesis in wine grapes

J. Tomas Matus, Center for Research in Agricultural Genomics, Barcelona, Spain

16.30 - 16.50

Study of genetic determinants of flavonol content in grape berry skin Giulia Malacarne, Fondazione Edmund Mach, San Michele All'Adige, Trento, Italy

16.50 - 17.10

Are VvMYB5a and VvMYB5b regulators of grape flavonoid pathway? Giambattista Tornielli, *University of Verona, Verona, Italy*

17.10 - 17.30

Investigation of VvDXS function and its effects on Muscat flavor levels
Juri Battilana, IASMA Research and Innnovation Centre, San Michele All'Adige, Trento, Italy

17.30 - 17.50

Microclimate influence on gene expression and metabolite accumulation in grape berries Christian Kappel, University of Bordeaux, Bordeaux, France

18.00 - 20.00 Welcome cocktail

GRAPEVINE ANTHOCYANIN TRANSPORT INVOLVES IN VIVO VESICLE-MEDIATED TRAFFICKING AND THE CONTRIBUTION OF ANTHOMATE TRANSPORTERS AND GST

C. Gomez¹, G. Conejero², L. Torregrosa³, V. Cheynier¹, N. Terrier¹, A. Ageorges¹

¹ INRA, UMR1083 Sciences Pour l'Œnologie, F-34060 Montpellier, France

³ Montpellier SupAgro, UMR1083 Sciences Pour l'Œnologie, F-34060 Montpellier, France E-mail: ageorges@supagro.inra.fr

Anthocyanins are plant secondary metabolites responsible for most of the red to purple pigmentation found in flowers, fruits and leaves. In grape berry, these compounds are accumulated during development from *véraison* to ripening, and are responsible for pigmentation found in red skin berries. Grapevine anthocyanins are 3-monoglucosides, which can be acylated by acetic, p-coumaric or caffeic acid. In the cell, anthocyanins are synthesized at the cytoplasmic surface of the endoplasmic reticulum and are then transported and finally accumulated inside the vacuole. In grapevine, two kinds of molecular actors are putatively associated in vacuolar sequestration of anthocyanins: a Glutathion S-transferase (GST) (1) and two MATE-type transporters, anthoMATE1 (AM1) and anthoMATE3 (AM3), which specifically mediate acylated anthocyanin transport *in vitro* (2). However, the molecular suite of mechanisms by which anthocyanins are imported into the vacuole remains unclear.

We used Myb-A1 transformed hairy roots ⁽³⁾ as a grapevine model tissue producing anthocyanins and took advantage of the unique autofluorescence of anthocyanins to study their cellular trafficking. In these tissues, anthocyanins were not only visible in the largest vacuoles but also, and more intensively present in several small vesicles, suggesting that these vesicles could mediate anthocyanins to the vacuole. In the cell, these vesicles actively moved alongside the tonoplast, suggesting a vesicular trafficking.

Subcellular homologous localisation assays in anthocyanin accumulating organs revealed that AMs were not only located to the tonoplast but also closely related with these small vesicles while GST was localised in the cytoplasm, and specifically around the nucleus, suggesting a close association with the endoplastic reticulum.

Furthermore, in hairy roots expressing AM3 antisense, no such anthocyanin containing vesicles were observed, while in hairy roots expressing GST antisense, anthocyanins were accumulated in these vesicles only but not in the vacuole. These observations suggest that in grapevine, AnthoMATE transporters are involved in vesicular transport of anthocyanins while GST could be involved in another yet unknown transport mechanism.

REFERENCE LIST

- [1] Conn et al. (2008) Purification, molecular cloning, and characterization of glutathione S-transferases (GSTs) from pigmented Vitis vinifera L. cell suspension cultures as putative anthocyanin transport proteins. *J Exp Bot* 59: 3621-3634
- [2] Gomez et al. (2009) Grapevine MATEtype proteins act as vacuolar H+-dependent acylated anthocyanin transporters. *Plant Physiol* <u>150</u>: 402-415
- [3] Cutanda-Perez et al. (2009) Ectopic expression of VlmybA1 in grapevine activates a narrow set of genes involved in anthocyanin synthesis and transport. *Plant Mol Biol* <u>69</u>: 633-648

² INRA, UMR5004 Biochimie et Physiologie Moléculaire des Plantes, Plate-forme histocytologie et d'imagerie cellulaire végétale, CIRAD, F-34398 Montpellier, France