

## Quantification of nitrogen fluxes and explanatory plant traits during a two-year legume-cereal rotation

Maé Guinet, Anne-Sophie Voisin, Bernard Nicolardot

## ▶ To cite this version:

Maé Guinet, Anne-Sophie Voisin, Bernard Nicolardot. Quantification of nitrogen fluxes and explanatory plant traits during a two-year legume-cereal rotation. 2. International Legume Society Conference - Legumes for a Sustainable World, Legume Society., Oct 2016, Lisbonne, Portugal. 358 p. hal-01608364

## HAL Id: hal-01608364

https://hal.science/hal-01608364

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.







## Legumes for a Sustainable World

ILS2 | Second International Legume Society Conference

11<sup>th</sup> - 14<sup>th</sup> OCT 2016 TRÓIA RESORT | PORTUGAL



2016-17. Performances variability has been analysed in relation with (a)biotic limiting factors and farmers practices.

14:40-14:50 Oral - \$16

Quantification of nitrogen fluxes and explanatory plant traits during a two-year legume-cereal rotation Guinet M., Voisin A-S., Nicolardot B.

UMR Agroécologie, AgroSup Dijon, INRA, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France

In the context of agroecological transition, the reintroduction of legume crops should play a key role in cropping system sustainability by allowing a reduction of nitrogen (N) inputs. But few references are available concerning the agronomical and ecological services provided by a wide range of legume crops, particularly within crops succession scale. Thus, the main objective of our study is to quantify the N fluxes during and after the legume crops taking into account 10 legume crops (peas, lupin, faba bean, soybean...). Our experiment consists in i) quantifying symbiotic N fixation depending on the amount of soil inorganic N, the mineralisation of N present in legume crop residues after soil incorporation and N losses outside of the soil-plant system (leaching, emission of nitrous oxide), ii) identifying plant biological traits associated to N fluxes. Thus, different N fluxes were quantified during a two-year field experiment, i.e. the first year (2014) legume crops were implanted and followed by wheat the second year (14-15) after incorportation of legume residues. This experiment will be repeated in 2016-2017. The first results concerning legume pre-crop effects will be presented during the congress, considering the effect of preceding legume crops on wheat seed yield and N content. This effect will be related to legume aerial and below ground residues that were characterized by their biomass, C/N content and potential mineralization.