



HAL
open science

Red and processed meat intake and cancer risk: Results from the prospective NutriNet-Santé cohort study

Abou Diallo, Mélanie Deschasaux, Paule Latino-Martel, Serge Hercberg, Maria del Pilar Galan Hercberg, Philippine Fassier, Benjamin Allès, Françoise Guéraud, Fabrice H.F. Pierre, Mathilde Touvier

► **To cite this version:**

Abou Diallo, Mélanie Deschasaux, Paule Latino-Martel, Serge Hercberg, Maria del Pilar Galan Hercberg, et al.. Red and processed meat intake and cancer risk: Results from the prospective NutriNet-Santé cohort study. *International Journal of Cancer*, 2017, in press (2), 10.1002/ijc.31046 . hal-01608688

HAL Id: hal-01608688

<https://hal.science/hal-01608688>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Red and processed meat intake and cancer risk: results from the prospective NutriNet-Santé cohort study

Abou Diallo^{1,2} , Mélanie Deschasaux¹, Paule Latino-Martel¹, Serge Hercberg^{1,2}, Pilar Galan¹, Philippine Fassier¹, Benjamin Allès¹, Françoise Guéraud³, Fabrice H. Pierre³, Mathilde Touvier¹

Affiliations:

¹ Sorbonne Paris Cité Epidemiology and Statistics Research Center (CRESS), Inserm U1153, Inra U1125, Cnam, Paris 13 University, Nutritional Epidemiology Research Team (EREN), Bobigny, France

² Département de Santé Publique, Hôpital Avicenne (AP-HP), F-93017 Bobigny, France

³ Toxalim (Research Centre in Food Toxicology), Université de Toulouse, INRA, ENVT, INP-Purpan, UPS, Toulouse, France

Authors's lastnames: Diallo, Deschasaux, Latino-Martel, Hercberg, Galan, Fassier, Allès, Guéraud, Pierre, Touvier

Corresponding author:

Abou Diallo

Sorbonne Paris Cité Epidemiology and Statistics Research Center (CRESS), Inserm U1153, Inra U1125, Cnam, Paris 13 University, Nutritional Epidemiology Research Team (EREN), rue Marcel Cachin, F-93017, Bobigny Cedex, France

e-mail: a.diallo@eren.smbh.univ-paris13.fr

Telephone number: +33 1 48 38 89 54

Sources of support: The NutriNet-Santé study was supported by the following public institutions: Ministère de la Santé, Institut de Veille Sanitaire (InVS), Institut National de la Prévention et de l'Éducation pour la Santé (INPES), Région Ile-de-France (CORDDIM), Institut National de la Santé et de la Recherche Médicale (INSERM), Institut National de la Recherche Agronomique (INRA), Conservatoire National des Arts et Métiers (CNAM) and Université Paris 13. Mélanie Deschasaux and Philippine Fassier were funded by PhD grants from the Cancéropôle Ile de France / Région Ile de France (public funding).

Keywords: red meat; processed meat; breast cancer; prostate cancer; prospective study

Conflict of interest: The authors have no conflict of interest to disclose.

Novelty and Impact: We observed that red meat intake was associated with increased overall and breast cancer risk, in line with mechanistic hypotheses from experimental studies. These findings suggest that limiting red meat intake may not only be beneficial for colorectal cancer, but also for the prevention of other tumor locations such as breast cancer.

Word count (from Introduction to Discussion/Conclusions): 2831

Tables: 3; **Figure:** 0

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1002/ijc.31046

ABSTRACT

The International Agency for Research on Cancer (WHO-IARC) classified red meat and processed meat as probably carcinogenic and carcinogenic for humans, respectively. These conclusions were mainly based on studies concerning colorectal cancer, but scientific evidence is still limited for other cancer locations. In this study, we investigated the prospective associations between red and processed meat intakes and overall, breast, and prostate cancer risk. This prospective study included 61,476 men and women of the French NutriNet-Santé cohort (2009-2015) aged ≥ 35 y and who completed at least 3 24h dietary records during the first year of follow-up. The risk of developing cancer was compared across sex-specific quintiles of red and processed meat intakes by multivariable Cox models. 1,609 first primary incident cancer cases were diagnosed during follow-up, among which 544 breast cancers and 222 prostate cancers. Red meat intake was associated with increased risk of overall cancers ($HR_{Q5vs.Q1}=1.31$ (1.10–1.55), $P_{trend}=0.01$) and breast cancer ($HR_{Q5vs.Q1}=1.83$ (1.33–2.51), $P_{trend}=0.002$). The latter association was observed in both premenopausal ($HR_{Q5vs.Q1}=2.04$ (1.03–4.06)) and postmenopausal women ($HR_{Q5vs.Q1}=1.79$ (1.26–2.55)). No association was observed between red meat intake and prostate cancer risk. Processed meat intake was relatively low in this study (cut-offs for the 5th quintile=46g/d in men and 29g/d in women) and was not associated with overall, breast or prostate cancer risk. This large cohort study suggested that red meat may be involved in carcinogenesis at several cancer locations (other than colon-rectum), in particular breast cancer. These results are consistent with mechanistic evidence from experimental studies.

INTRODUCTION

The International Agency for Research on Cancer (WHO-IARC) recently classified consumption of processed meat as “carcinogenic to humans” and consumption of red meat as “probably carcinogenic to humans”(1) The World Cancer Research Fund and the American Institute for Cancer Research (WCRF/AICR) recommends consuming less than 500g/week of red meat and less than 50g/d of processed meat (2). These conclusions were mainly based on findings concerning colorectal cancer, for which the weight of evidence is considered as convincing (3;4). Indeed, experimental studies showed that several components of red and/or processed meat act locally on the colorectal mucosa to promote carcinogenesis. Potential carcinogens include heme iron, nitrates and nitrites and mutagenic compounds such as neoformed products generated in red meats and processed meat (heterocyclic amines, polycyclic aromatic hydrocarbons, N-nitroso compounds (3;5;6)). However, these pro-carcinogens may also be involved in more systemic mechanisms (7-10), suggesting that red and processed meat may impact cancer risk for cancer locations other than colon-rectum.

Despite these mechanistic hypotheses, epidemiological evidence regarding red/processed meat and cancer risk is limited for other cancer locations, and notably for breast and prostate cancers, which are the two main cancer sites in many Western countries (11;12). In a previous study performed in the SU.VI.MAX cohort (13), we observed that processed meat intake was associated with increased breast cancer risk. This result is consistent with two recent meta-analyses suggesting positive associations with breast cancer risk (14;15). Since the publication of these meta-analyses, two prospective cohort studies were published. Inoue-Cho et al (16) observed an increased risk of breast cancer in post-menopausal women with high consumption of red or processed meat; and Bertrand et al (17) showed increased breast density in pre-menopausal

women associated with high consumption of red meat. In 2014, the World Cancer Research Fund and the American Institute for Cancer Research (WCRF/AICR) observed null results for their meta-analyses of the associations between red and processed meat and prostate cancer risk (18), consistent with a meta-analysis published in 2015 (19). In contrast, in a pooled analysis of 15 cohort studies published in 2016, Wu et al (20) observed a positive association between red and processed meat and risk of advanced prostate cancer. Thus, the weight of evidence is still considered as “limited” regarding red and processed meat and cancer risk for non-colorectal locations (3;18;21;22). No consensus has been reached so far and additional prospective studies are needed to more thoroughly elucidate the relationship between red and processed meat intakes and breast or prostate cancer risk.

The objective of this prospective study was to investigate the associations between red meat and processed meat intakes and overall, breast and prostate cancer risk, in a large cohort of French adults with accurate and up-to-date dietary intake data.

METHODS

Study population

The NutriNet-Santé study is an ongoing web-based cohort launched in 2009 in France with the objective to study the associations between nutrition and health as well as the determinants of dietary behaviors and nutritional status. This cohort has been previously described in details (23). Participants aged over 18 years with access to the Internet are continuously recruited since May 2009 among the general population by means of vast multimedia campaigns. All questionnaires are completed online using a dedicated website (www.etude-nutrinet-sante.fr). The NutriNet-Santé study is conducted according to the Declaration of Helsinki guidelines and was approved by the Institutional Review Board of the French Institute for Health and Medical Research (IRB

Inserm n°0000388FWA00005831) and the "Commission Nationale de l'Informatique et des Libertés" (CNIL n°908450/n°909216). Electronic informed consent is obtained from each participant (EudraCT no. 2013-000929-31).

Data collection

Dietary data

Dietary intakes were assessed every six months through a series of three non-consecutive validated web-based 24h-dietary records, randomly assigned over a 2-week period (2 weekdays and 1 weekend day) (24-26). Participants used a dedicated interface of the study website to declare all foods and beverages consumed during a 24h-period: three main meals (breakfast, lunch, dinner) or any other eating occasion. Portion sizes were estimated using validated photographs (27). Mean daily energy, alcohol and nutrient intakes were estimated using a published French food composition table (>3300 items) (28). Amounts consumed from composite dishes were estimated using French recipes validated by food and nutrition professionals. Dietary underreporting was identified on the basis of the method proposed by Black (29). Red meat intake was defined as fresh, minced and frozen beef, veal, pork, and lamb. Processed meat intake was defined as mostly pork and beef preserved by methods other than freezing, such as salting, smoking, marinating, air-drying or heating and included ham, bacon, sausages, blood sausages, liver pâté, salami, mortadella, tinned meat and others.

Covariates

At inclusion, participants fulfilled a set of five questionnaires related to socio-demographic and lifestyle characteristics (30) (e.g. sex, date of birth, educational level, smoking status, number of children), anthropometrics (31;32) (e.g. height and weight), dietary intakes (see above), physical

activity (validated IPAQ questionnaire) (33), and health status (e.g. personal and family history of diseases, medication use including hormonal treatment for menopause and oral contraception, menopausal status).

Case ascertainment

Participants self-declared health events through the yearly health status questionnaire, through a specific check-up questionnaire for health events (every three months) or at any time through a specific interface on the study website. Following this declaration, participants were invited to send their medical records (diagnosis, hospitalization, etc.) and, if necessary, the study physicians contacted the participants' treating physician or the medical structures to collect additional information. Then, data were reviewed by an independent physician expert committee for the validation of major health events. Cancer cases were classified using the International Chronic Diseases Classification, 10th Revision, Clinical Modification (ICD-10) (34). In this study, all first primary cancers diagnosed between the inclusion and August 2015 were considered as cases (except basal cell skin carcinoma, which was not considered as cancer).

Statistical analyses

So far, 96716 subjects without cancer at baseline provided at least three valid 24h-dietary records during their first year of follow-up. Participants aged <35y (n=32882) were excluded because of a very low susceptibility to develop cancer and so were subjects with a null follow-up (n=2358). Thus, 61476 subjects were included in the analyses.

Estimated red and processed meat and other dietary intakes were based on the average intake for each subject across all 24h-dietary records available in their first year of follow-up. For all covariates except physical activity, $\leq 5\%$ of values were missing and were imputed to the modal

value. For physical activity (13% of missing values), a “missing class” was introduced into the models.

Baseline characteristics of participants were compared across sex-specific quintiles of red and processed meat intake using χ^2 tests or Fisher tests wherever appropriate. We estimated hazard ratios (HR) and 95% confidence intervals (CI) using Cox proportional hazards models, with age as the primary time variable, to characterize the association between sex-specific quintiles of red meat, processed meat and total red and processed meat intake and incidence of overall, breast or prostate cancer risk (the two main cancer locations in the cohort). We confirmed that the assumptions of proportionality were satisfied through examination of the log-log (survival) vs. log-time plots. Tests for linear trend were performed using the ordinal score on sex-specific quintiles of intake. Participants contributed person-time until the date of cancer diagnosis, the date of last completed questionnaire, the date of death, or August 31 2015, whichever occurred first. For cancer site specific analysis, women who reported a cancer other than breast cancer and men who reported a cancer other than prostate cancer during the study period were censored at the date of diagnosis. Analyses were performed according to menopausal status for breast cancer analyses. For these analyses, women contributed person-time in the Cox model until their date of menopause for premenopausal breast cancer analysis or from their date of menopause for postmenopausal breast cancer analysis. Additionally, models restricted to invasive breast cancer cases (excluding in situ cases) were tested.

Models were adjusted for age (time-scale), sex (for overall cancers only), BMI (kg/m², continuous), height (cm, continuous), physical activity (high, moderate, low, computed following IPAQ recommendations (35)), smoking status (never smokers, former smokers, current smokers), number of 24h-dietary records (continuous), fruits and vegetables intake (g/d, continuous), total lipids intake (g/d, continuous), alcohol intake (g/d, continuous), energy intake (without alcohol,

g/d, continuous), family history of cancer (yes/no), and educational level (<high-school degree, <2 years after high-school degree, ≥ 2 years after high-school degree). Red and processed meat models were mutually adjusted for processed meat and red meat intakes, respectively. For breast cancer analyses, additional adjustments were performed for the number of biological children (continuous), menopausal status at baseline (yes/no), hormonal treatment for menopause at baseline (only for postmenopausal analyses, yes/no) and oral contraception use at baseline (only for premenopausal analyses, yes/no). Since antioxidants may partly counteract lipid peroxidation by heme iron from red and processed meat (i.e. one of the hypothesized mechanisms involved in their potentially procarcinogenic effect) (1), we have tested for a potential interaction between fruit and vegetable intake (as a proxy for antioxidant exposure, according to sex-specific median intake) and red and processed meat intake by introducing the product of the two variables into Cox models for each cancer location. Stratified analyses were performed when appropriate (i.e. p-interaction < 0.1).

All tests were two-sided, and $P < 0.05$ was considered statistically significant. SAS version 9.4 (SAS Institute) was used for the analyses.

RESULTS

Between May 2009 and August 2015 (median follow-up time: 4.1y; 229835 person-years), 1609 incident cancer cases were diagnosed, among which 544 breast cancers (169 premenopausal and 375 postmenopausal; 71.6% ER+/PR+, 13.5% ER-/PR-, 14.6% ER+/PR-, 0.3% ER-/PR+; 80.4% invasive and 19.6% in situ), 222 prostate cancers (88,46% Gleason score < 7, 11,54% Gleason score ≥ 7) and 843 other cancers (169 skin (other than basal cell carcinoma), 120 colorectal, 64 lymphomas, 63 lung, 39 thyroid, 38 cervix, 38 bladder, 37 uterus, 35 leukemia, 30 kidney, and 210 others).

Mean age at diagnosis was $51.68y \pm 10.14$ and mean baseline-to-diagnosis time was $2.43y \pm 1.60$.

Mean number of 24h dietary records per subject over their first year of follow-up was 4.53 ± 1.61 .

Characteristics of the participants according to quintiles of total red and processed meat intakes are described in **Table 1**. Mean daily red meat intake was 42.9 ± 39.0 g/d (0.4 ± 1.9 g/d in the first quintile, 102.3 ± 33.7 g/d in the fifth quintile). Mean daily processed meat intake was 19.1 ± 23.8 g/d (0 g/d in the first quintile, 56.0 ± 25.9 g/d in the fifth quintile) [data not tabulated]. Subjects with higher total red and processed meat intake were more likely to be younger, to have a higher body mass index, to smoke, to have higher energy, lipid and alcohol intakes and lower fruit and vegetable intake, to have a lower educational level and to be less physically active.

Associations between red and processed meat intakes and overall, breast and prostate cancer risk are presented in **Table 2**. Red meat intake was associated with increased overall cancer risk (HR_{Q5vs.Q1}=1.31; 95% CI 1.10, 1.55; $P_{\text{trend}}=0.01$) and increased breast cancer risk (HR_{Q5,Q1}=1.83; 95% CI 1.33, 2.51; $P_{\text{trend}}=0.002$), but not with prostate cancer risk ($P_{\text{trend}}=0.9$). This association between red meat intake and increased breast cancer risk was observed in both premenopausal (HR_{Q5vs.Q1}=2.04 (1.03-4.06)) and postmenopausal women (HR_{Q5vs.Q1}=1.79 (1.26-2.55)) (**Table 3**), and was similarly observed when analyses excluded cases diagnosed during their first year of follow-up (413 cases/40,892 non-cases included; HR_{Q5vs.Q1}=1.82 (1.27, 2.62)) or when analyses were restricted to invasive breast cancers (470 cases/45,386 non-cases; HR_{Q5vs.Q1}=1.78 (1.26, 2.50)) [data not tabulated]. Results with and without BMI adjustment were very similar for overall, breast and prostate cancers models (without BMI adjustment in Supplemental Table 2). No association was detected for processed meat intake ($P_{\text{trend}}=0.5$, 0.4 and 0.3 for overall, breast and prostate cancers, respectively, Table 2). No association was statistically significant for red or

processed meat intake with colorectal or with lung cancers or with lymphomas (Supplemental Table 1). No interaction was detected between red or processed meat intake and fruit and vegetable or individual antioxidant intakes (vitamins C, E, beta-carotene and selenium) regarding overall and site-specific cancer risk (all $P > 0.05$, data not shown).

DISCUSSION

In this large prospective cohort, red meat intake was significantly associated with increased overall and breast cancer risks. No association was observed for prostate cancer. Processed meat intake was not associated with cancer risk in this study.

For red meat, our result of a direct association with breast cancer risk is consistent with two recent meta-analyses: Guo and al. (14) based on 14 cohort studies for red meat and 12 cohort studies for processed meat, and Wu and al (15) based on 12 cohort studies for red meat and 15 cohort studies for processed meat, both showing positive associations with breast cancer risk. The two prospective studies published after this meta-analysis also suggest direct associations between red meat intake and post-menopausal breast cancer risk in the NIH-AARP cohort (16) and increased breast density (17). In a previous study performed on the SU.VI.MAX cohort, we did not observe statistically significant relationships between red meat and breast cancer risk. However, red meat intakes in women of the SU.VI.MAX cohort were relatively low (fourth quartile < 500 g/week), while they were higher in the present NutriNet-Santé cohort, where 19.60% exceeded 500g of red meat per week. In the French general population, about one out of four adults consume more than 500g/week of red meat (3). In Europe the median range of daily red meat intake is 24-57 g/day (36), while mean intake is about 53 g/d in the U.S (37).

Regarding prostate cancer, our null result is consistent with two large and recent meta-analyses of prospective studies, performed by the WCRF/AICR in 2014 (18) and Blyma et al. in 2015 (19). In a pooled analysis of 15 cohort studies, Wu et al. (20) did not observe any association between red meat intake and overall prostate cancer risk, but showed a modest positive association for tumors identified as advanced stage at diagnosis. In our study, our results did not differ according to Gleason score ($<$ or ≥ 7) [data not shown]. However, statistical power was limited for this sub-analysis. In the WCRF/AICR meta-analyses, the summary RR were not statistically significant for the different prostate cancer subtypes, (RR per 100g/d = 0.99 (0.89, 1.11) for advanced/high grade and 1.19 (0.88, 1.59) for fatal cases) (18).

The small number of cancer cases other than breast and prostate locations did not allow us to have enough statistical analysis to conclude for these locations. However, the pro-carcinogenic effect of high red meat intake on colorectal carcinogenesis has been well established in several national and international collective expert evaluations (1;3;4). In 2012, the WCRF/AICR also judged the direct association between red meat intake and pancreatic cancer risk as “suggestive”. Along with the positive association observed for breast cancer, these may contribute to explain the positive association observed in the present study between red meat intake and overall cancer risk. It is also possible that the lack of association with processed meat might be a chance finding or could change with longer follow-up.

While several studies suggested direct associations between processed meat intake and colorectal (1;3;4), breast (13;14;16), stomach (38), or pancreatic (39) cancer risk, no association was detected in the present study. This may be explained by the fact that processed meat intakes were too low to properly investigate any adverse effect. Indeed, the cut-off for quintile 5 of processed meat intake was 45.9 g/d for men and 29.0 g/d for women, i.e. lower than the 50g/d upper dose

recommended by the WCRF/AICR for colorectal cancer prevention (4). In the French general population, more than one out of four adults consume at least 50 g of processed meat per day (3). In Europe the median range of daily processed meat intake is 5-49 g/day (36), while mean intake is about 18 g/d in the U.S (37).

Our epidemiological findings are supported by mechanistic data. Red and processed meat contain pro-carcinogenic components, such as heterocyclic aromatic amines (HAA), polycyclic aromatic hydrocarbons (PAHs) resulting from meat processing or preparation (such as cooking at high-temperature), nitrites (used as additives) and induces N-nitroso compounds (NOCs) formation in the digestive tract (40-44). These chemicals may exert a pro-carcinogenic effect through direct DNA damage and have been associated with mammary tumor development in animal (7;9;41) and human (8;10;45) studies (13;46;47). Most importantly, red meat contains high levels of heme iron, which may contribute to initiate carcinogenesis via several mechanisms, including the production of genotoxic free radicals, NOCs or through lipid peroxidation (5;48-50).

Strengths of this study include its prospective design, its large sample size, and the assessment of usual dietary intakes using repeated 24h-dietary records based on a recent food composition database with a large choice of items (>3300). These repeated 24h-dietary records allowed a better insight into the food products consumed compared to food frequency questionnaires with more aggregated food items. However, some limitations should be acknowledged. First, caution is needed regarding the extrapolation of these results since this study included volunteers involved in a long-term cohort study investigating the association between nutrition and health, with overall more health-conscious behaviors and higher professional and/or educational level compared to the general French population. Thus, unhealthy dietary behaviors may have been

underrepresented in this study, which may have weakened the observed associations and may have prevented us from observing significant associations for processed meat. Second, although the number of overall cancer cases was reasonably large, the number of cancers at any given site was more restricted, which did not allow us to investigate more cancers sites and receptor types for breast cancer. Finally, the observed relationships could be partly affected by unmeasured or residual confounding. However, main potential confounders have been accounted for in this study; thus, it is unlikely that residual confounding entirely explains the observed associations.

In conclusion, this prospective cohort study brings new contribution into the role of red and processed meat intake as cancer risk factors. We observed that red meat intake was associated with increased overall and breast cancer risk, in line with mechanistic hypotheses from experimental studies. If confirmed, these findings suggest that limiting red meat intake may not only be beneficial for colorectal cancer, but also for the prevention of other tumor locations such as breast cancer.

ACKNOWLEDGEMENTS

The authors thank all the volunteers of the NutriNet-Santé cohort. We also thank Véronique Gourlet, Nathalie Arnault, Stephen Besseau, Laurent Bourhis, Than Duong Van, Younes Esseddik, Paul Flanzy, Julien Allègre, Mac Rakotondrazafy, Fabien Szabo, Cédric Agaesse, Claudia Chahine, Anne-Elise Dussoulier, Roland Andrianasolo and Fatoumata Diallo for their technical contribution to the NutriNet-Santé study. This work was conducted in the framework of the French network for Nutrition And Cancer Research (NACRe network), www.inra.fr/nacre.

The author's responsibilities were as follows – AD and MT: designed the research; SH, MT, PG: conducted the research; AD: performed statistical analysis and wrote the paper; MT: supervised the study; MD, PLM, PG, SH, PF, BA, FG, FHP: contributed to the data interpretation and revised each draft for important intellectual content; and MT had primary responsibility for final content. All authors read and approved the final manuscript. The funders had no role in the design, implementation, analysis, or interpretation of the data. The authors have no conflict of interest to disclose.

Reference List

- (1) Bouvard V, Loomis D, Guyton KZ, Grosse Y, Ghissassi FE, Benbrahim-Tallaa L, Guha N, Mattock H, Straif K. Carcinogenicity of consumption of red and processed meat. *Lancet Oncol* 2015 Dec;16(16):1599-600.
- (2) World Cancer Research Fund / American Institute for Cancer Research. Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. 2007.
- (3) Latino-Martel P, Cottet V, Druesne-Pecollo N, Pierre FH, Touillaud M, Touvier M, Vasson MP, Deschasaux M, Le MJ, Barrandon E, Ancellin R. Alcoholic beverages, obesity, physical activity and other nutritional factors, and cancer risk: A review of the evidence. *Crit Rev Oncol Hematol* 2016 Mar;99:308-23.
- (4) WCRF/AICR. Colorectal cancer 2011 report : Food, Nutrition, Physical Activity, and the Prevention of Colorectal Cancer. 2011.
- (5) Bastide NM, Pierre FH, Corpet DE. Heme iron from meat and risk of colorectal cancer: a meta-analysis and a review of the mechanisms involved. *Cancer Prev Res (Phila)* 2011 Feb;4(2):177-84.
- (6) Abid Z, Cross AJ, Sinha R. Meat, dairy, and cancer. *Am J Clin Nutr* 2014 Jul;100 Suppl 1:386S-93S.
- (7) El-Bayoumy K, Chae YH, Upadhyaya P, Rivenson A, Kurtzke C, Reddy B, Hecht SS. Comparative tumorigenicity of benzo[a]pyrene, 1-nitropyrene and 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine administered by gavage to female CD rats. *Carcinogenesis* 1995 Feb;16(2):431-4.
- (8) Gammon MD, Santella RM, Neugut AI, Eng SM, Teitelbaum SL, Paykin A, Levin B, Terry MB, Young TL, Wang LW, Wang Q, Britton JA, et al. Environmental toxins and breast cancer on Long Island. I. Polycyclic aromatic hydrocarbon DNA adducts. *Cancer Epidemiol Biomarkers Prev* 2002 Aug;11(8):677-85.
- (9) Shirai T, Tamano S, Sano M, Masui T, Hasegawa R, Ito N. Carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in rats: dose-response studies. *Princess Takamatsu Symp* 1995;23:232-9.
- (10) Sinha R, Gustafson DR, Kullendorff M, Wen WQ, Cerhan JR, Zheng W. 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine, a carcinogen in high-temperature-cooked meat, and breast cancer risk. *J Natl Cancer Inst* 2000 Aug 16;92(16):1352-4.
- (11) Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015 Mar 1;136(5):E359-E386.
- (12) INCa. Les cancers en France, Les Données, INCa, édition 2015. INCa; 2016. Report No.: ISBN 978-2-37219-066-4.

- (13) Pouchieu C, Deschasaux M, Hercberg S, Druesne-Pecollo N, Latino-Martel P, Touvier M. Prospective association between red and processed meat intakes and breast cancer risk: modulation by an antioxidant supplementation in the SU.VI.MAX randomized controlled trial. *Int J Epidemiol* 2014 Oct;43(5):1583-92.
- (14) Guo J, Wei W, Zhan L. Red and processed meat intake and risk of breast cancer: a meta-analysis of prospective studies. *Breast Cancer Res Treat* 2015 May;151(1):191-8.
- (15) Wu J, Zeng R, Huang J, Li X, Zhang J, Ho JC, Zheng Y. Dietary Protein Sources and Incidence of Breast Cancer: A Dose-Response Meta-Analysis of Prospective Studies. *Nutrients* 2016 Nov 17;8(11).
- (16) Inoue-Choi M, Sinha R, Gierach GL, Ward MH. Red and processed meat, nitrite, and heme iron intakes and postmenopausal breast cancer risk in the NIH-AARP Diet and Health Study. *Int J Cancer* 2016 Apr 1;138(7):1609-18.
- (17) Bertrand KA, Burian RA, Eliassen AH, Willett WC, Tamimi RM. Adolescent intake of animal fat and red meat in relation to premenopausal mammographic density. *Breast Cancer Res Treat* 2016 Jan;155(2):385-93.
- (18) WCRF/AICR. Diet, nutrition, physical activity and prostate cancer. Continuous Update Project . 2014.
- (19) Bylsma LC, Alexander DD. A review and meta-analysis of prospective studies of red and processed meat, meat cooking methods, heme iron, heterocyclic amines and prostate cancer. *Nutr J* 2015;14:125.
- (20) Wu K, Spiegelman D, Hou T, Albanes D, Allen NE, Berndt SI, van den Brandt PA, Giles GG, Giovannucci E, Alexandra GR, Goodman GG, Goodman PJ, et al. Associations between unprocessed red and processed meat, poultry, seafood and egg intake and the risk of prostate cancer: A pooled analysis of 15 prospective cohort studies. *Int J Cancer* 2016 May 15;138(10):2368-82.
- (21) WCRF/AICR. Breast cancer 2010 report : Food, Nutrition, Physical Activity, and the Prevention of Breast Cancer. 2010.
- (22) WCRF/AICR. Cancer prevention and survival : Summary of global evidence on diet, weight, physical activity & what increases or decreases your risk of cancer. 2016.
- (23) Hercberg S, Castetbon K, Czernichow S, Malon A, Mejean C, Kesse E, Touvier M, Galan P. The Nutrinet-Sante Study: a web-based prospective study on the relationship between nutrition and health and determinants of dietary patterns and nutritional status. *BMC Public Health* 2010;10:242.
- (24) Lassale C, Castetbon K, Laporte F, Deschamps V, Vernay M, Camilleri GM, Faure P, Hercberg S, Galan P, Kesse-Guyot E. Correlations between Fruit, Vegetables, Fish, Vitamins, and Fatty Acids Estimated by Web-Based Nonconsecutive Dietary Records and Respective Biomarkers of Nutritional Status. *J Acad Nutr Diet* 2016 Mar;116(3):427-38.

- (25) Lassale C, Castetbon K, Laporte F, Camilleri GM, Deschamps V, Vernay M, Faure P, Hercberg S, Galan P, Kesse-Guyot E. Validation of a Web-based, self-administered, non-consecutive-day dietary record tool against urinary biomarkers. *Br J Nutr* 2015 Mar 28;113(6):953-62.
- (26) Touvier M, Kesse-Guyot E, Mejean C, Pollet C, Malon A, Castetbon K, Hercberg S. Comparison between an interactive web-based self-administered 24 h dietary record and an interview by a dietitian for large-scale epidemiological studies. *Br J Nutr* 2011 Apr;105(7):1055-64.
- (27) Le Moulenc N, Deheeger M, Preziosi P, Montero P, Valeix P, Rolland-Cachera M, Potier de Courcy G, Christides J, Galan P, Hercberg S. Validation du manuel photo utilisé pour l'enquête alimentaire de l'étude SU.VI.MAX. [Validation of the food portion size booklet used in the SU.VI.MAX study] (in French). *Cah Nutr Diet* 1996;31:158-64.
- (28) Arnault N, Caillot L, Castetbon Keal. Table de composition des aliments, étude NutriNet-Santé. [Food composition table, NutriNet-Santé study] (in French). Paris: Les éditions INSERM/Economica. 2013.
- (29) Black AE. Critical evaluation of energy intake using the Goldberg cut-off for energy intake:basal metabolic rate. A practical guide to its calculation, use and limitations. *Int J Obes Relat Metab Disord* 2000 Sep;24(9):1119-30.
- (30) Vergnaud AC, Touvier M, Mejean C, Kesse-Guyot E, Pollet C, Malon A, Castetbon K, Hercberg S. Agreement between web-based and paper versions of a socio-demographic questionnaire in the NutriNet-Sante study. *Int J Public Health* 2011 Aug;56(4):407-17.
- (31) Lassale C, Peneau S, Touvier M, Julia C, Galan P, Hercberg S, Kesse-Guyot E. Validity of web-based self-reported weight and height: results of the Nutrinet-Sante study. *J Med Internet Res* 2013;15(8):e152.
- (32) Touvier M, Mejean C, Kesse-Guyot E, Pollet C, Malon A, Castetbon K, Hercberg S. Comparison between web-based and paper versions of a self-administered anthropometric questionnaire. *Eur J Epidemiol* 2010 May;25(5):287-96.
- (33) Craig CL, Marshall AL, Sjostrom M, Bauman AE, Booth ML, Ainsworth BE, Pratt M, Ekelund U, Yngve A, Sallis JF, Oja P. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003 Aug;35(8):1381-95.
- (34) WHO. ICD-10, International Classification of Diseases and Related Health Problems. 10th Revision. Geneva: World Health Organization. ; 1993. Report No.: ISBN 978-2-37219-066-4.
- (35) IPAQ Group. Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ). ; 2005. Report No.: ISBN 978-2-37219-066-4.
- (36) Linseisen J, Kesse E, Slimani N, Bueno-de-Mesquita HB, Ocke MC, Skeie G, Kumle M, Dorronsoro IM, Morote GP, Janson L, Stattin P, Welch AA, et al. Meat consumption in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts: results from 24-hour dietary recalls. *Public Health Nutr* 2002 Dec;5(6B):1243-58.

- (37) Daniel CR, Cross AJ, Koebnick C, Sinha R. Trends in meat consumption in the USA. *Public Health Nutr* 2011 Apr;14(4):575-83.
- (38) WCRF/AICR. Stomach cancer 2016 report : Diet, nutrition, physical activity and stomach cancer. 2016.
- (39) WCRF/AICR. Pancreatic cancer 2012 report : Food, Nutrition, Physical Activity and the Prevention of Pancreatic Cancer. 2012.
- (40) Genkinger JM, Koushik A. Meat consumption and cancer risk. *PLoS Med* 2007 Dec;4(12):e345.
- (41) Ito N, Hasegawa R, Sano M, Tamano S, Esumi H, Takayama S, Sugimura T. A new colon and mammary carcinogen in cooked food, 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP). *Carcinogenesis* 1991 Aug;12(8):1503-6.
- (42) Joosen AM, Kuhnle GG, Aspinall SM, Barrow TM, Lecommandeur E, Azqueta A, Collins AR, Bingham SA. Effect of processed and red meat on endogenous nitrosation and DNA damage. *Carcinogenesis* 2009 Aug;30(8):1402-7.
- (43) Santarelli RL, Vendevre JL, Naud N, Tache S, Gueraud F, Viau M, Genot C, Corpet DE, Pierre FH. Meat processing and colon carcinogenesis: cooked, nitrite-treated, and oxidized high-heme cured meat promotes mucin-depleted foci in rats. *Cancer Prev Res (Phila)* 2010 Jul;3(7):852-64.
- (44) Tricker AR. N-nitroso compounds and man: sources of exposure, endogenous formation and occurrence in body fluids. *Eur J Cancer Prev* 1997 Jun;6(3):226-68.
- (45) Rundle A, Tang D, Hibshoosh H, Estabrook A, Schnabel F, Cao W, Grumet S, Perera FP. The relationship between genetic damage from polycyclic aromatic hydrocarbons in breast tissue and breast cancer. *Carcinogenesis* 2000 Jul;21(7):1281-9.
- (46) Steck SE, Gaudet MM, Eng SM, Britton JA, Teitelbaum SL, Neugut AI, Santella RM, Gammon MD. Cooked meat and risk of breast cancer--lifetime versus recent dietary intake. *Epidemiology* 2007 May;18(3):373-82.
- (47) Egeberg R, Olsen A, Autrup H, Christensen J, Stripp C, Tetens I, Overvad K, Tjønneland A. Meat consumption, N-acetyl transferase 1 and 2 polymorphism and risk of breast cancer in Danish postmenopausal women. *Eur J Cancer Prev* 2008 Feb;17(1):39-47.
- (48) Tappel A. Heme of consumed red meat can act as a catalyst of oxidative damage and could initiate colon, breast and prostate cancers, heart disease and other diseases. *Med Hypotheses* 2007;68(3):562-4.
- (49) Bastide NM, Chenni F, Audebert M, Santarelli RL, Tache S, Naud N, Baradat M, Jouanin I, Surya R, Hobbs DA, Kuhnle GG, Raymond-Letron I, et al. A central role for heme iron in colon carcinogenesis associated with red meat intake. *Cancer Res* 2015 Mar 1;75(5):870-9.
- (50) Santarelli RL, Pierre F, Corpet DE. Processed meat and colorectal cancer: a review of epidemiologic and experimental evidence. *Nutr Cancer* 2008;60(2):131-44.

Table 1 Baseline characteristics of study participants (n=61476) according to sex-specific quintiles of red and processed meat intake, NutriNet-Santé cohort, France, 2009 – 2016^a

	Quintile 1 (n= 12 292)	Quintile 2 (n= 12 298)	Quintile 3 (n= 12 303)	Quintile 4 (n= 12 287)	Quintile 5 (n= 12 296)
Age, y	51.7 +/- 10.2	52 +/- 10.3	52.2 +/- 10.2	51.9 +/- 10.1	50.6 +/- 9.8
Sex					
Men	3107 (25.28)	3111 (25.29)	3110 (25.28)	3108 (25.30)	3110 (25.29)
Women	9185(74.72)	9187(74.71)	9193(74.72)	9179(74.70)	9186(74.71)
Height, cm	166.5 +/- 8.2	166.4 +/- 8.1	166.4 +/- 8.2	166.6 +/- 8.2	167.1 +/- 8.3
Body mass index, kg/m ²	23.4 +/- 4.2	23.9 +/- 4.2	24.3 +/- 4.4	24.9 +/- 4.6	25.7 +/- 5.2
Family history of cancer ^c , yes	5429 (44.2)	5557 (45.2)	5627 (45.7)	5570 (45.3)	5465 (44.4)
Number of children, n	1.7 +/- 1.2	1.8 +/- 1.2	1.9 +/- 1.1	1.9 +/- 1.2	1.9 +/- 1.1
Higher education					
No	2,718 (22.1)	2,806 (22.8)	3,034 (24.7)	3,316 (27.0)	3,543 (28.8)
Yes, < 2 years	1,885 (15.3)	1,876 (15.3)	1,785 (14.5)	1,955 (15.9)	2,042 (16.6)
Yes, ≥ 2 years	7,689 (62.6)	7,616 (61.9)	7,484 (60.8)	7,016 (57.1)	6,711 (54.6)
Smoking status					
Current	1476 (12.0)	1476 (12.0)	1542 (12.5)	1777 (14.5)	2116 (17.2)
Former	5063 (41.2)	5046 (41.0)	5000 (40.6)	5042 (41.0)	5109 (41.6)
Never	5753 (46.8)	5776 (47.0)	5761 (46.8)	5468 (44.5)	5071 (41.2)
IPAQ Physical activity level ^d					
High	4292 (34.9)	4056 (33)	3881 (31.5)	3802 (30.9)	3628 (29.5)
Moderate	4524 (36.8)	4460 (36.3)	4413 (35.9)	4290 (34.9)	4001 (32.5)
Low	2049 (16.7)	2316 (18.8)	2452 (19.9)	2600 (21.2)	2870 (23.3)
Processed meat intake, g/d	3.8 +/- 6.1	11.6 +/- 12.2	17.3 +/- 16.7	24.2 +/- 21.6	38.6 +/- 34.6
Red meat intake, g/d	3.6 +/- 6.7	22.3 +/- 13.8	38.5 +/- 17.9	56.2 +/- 23.0	93.9 +/- 42.0
Fruits and vegetables intake, g/d	496.9 +/- 257.7	458.7 +/- 211.6	442.4 +/- 203.0	431.6 +/- 198.8	411.4 +/- 203.5
Energy intake, kcal/d	1720 +/- 443.9	1769.1 +/- 425.2	1805 +/- 433.3	1844.5 +/- 436.0	1962.3 +/- 492.1
Total lipid intake, g/d	73 +/- 24.5	76.4 +/- 23.1	79.3 +/- 23.4	82.6 +/- 24.2	90.9 +/- 27.8
Alcohol intake, g/d	6.3 +/- 10.1	7.8 +/- 11.0	8.7 +/- 11.7	9.9 +/- 13.3	11.9 +/- 15.9
Oral contraception, yes	1021 (11.1)	1118 (12.2)	1112 (12.1)	1176 (12.8)	1299 (14.1)
Hormonal treatment for menopause, yes	750 (8.2)	830 (9.0)	882 (9.6)	827 (9.0)	699 (7.6)
Menopausal, yes	4557 (49.6)	4546 (49.5)	4698 (51.1)	4537 (49.4)	4056 (44.2)

^a Values are means ±SDs or n (%). Cut-offs for quintiles of red and processed meat intake were 32.00; 59.82; 86.81 and 122.14 g/d in men and 18.21 ; 39.73 ; 60.00 and 87.68 g/d in women.

^c Among first-degree relatives.

^d Missing for 7,842 (12.76%) subjects.

Table 2 Associations between quintiles of red and processed meat intake and overall, breast, and prostate cancer risk, from multivariable Cox proportional hazard models, NutriNet-Santé cohort, France, 2009 – 2016 (n=61476)^a

	Quintile 1	Quintile 2	Quintile 3	Quintile 4	Quintile 5	p-trend
Red meat						
All cancers						0.01
N for cases/non-cases	233/12101	359/11898	307/12001	358/11876	352/11991	
Multivariable HR (95%CI)	1	1.24 (1.05 , 1.47)	1.06 (0.89 , 1.26)	1.22 (1.03 , 1.45)	1.31 (1.10 , 1.55)	
Breast cancer						0.002
N for cases/non-cases	59/9160	124/9030	114/9076	123/9010	124/9110	
Multivariable HR (95%CI)	1	1.68 (1.23 , 2.31)	1.58 (1.14 , 2.17)	1.70 (1.24 , 2.34)	1.83 (1.33 , 2.51)	
Prostate cancer						0.9
N for cases/non-cases	28/3087	66/3037	33/3085	54/3047	41/3068	
Multivariable HR (95%CI)	1	1.70 (1.09 , 2.68)	0.87 (0.52 , 1.45)	1.38 (0.86 , 2.20)	1.28 (0.78 , 2.11)	
Processed meat						
All cancers						0.5
N for cases/non-cases	403/17148	221/6830	350/11929	351/11949	284/12011	
Multivariable HR (95%CI)	1	1.08 (0.91 , 1.28)	1.03 (0.88 , 1.19)	1.05 (0.90 , 1.22)	0.93 (0.79 , 1.10)	
Breast cancer						0.4
N for cases/non-cases	133/13809	63/4380	113/9055	134/9057	101/9085	
Multivariable HR (95%CI)	1	1.19 (0.88 , 1.62)	1.08 (0.83 , 1.39)	1.28 (1.00 , 1.64)	1.05 (0.80 , 1.38)	
Prostate cancer						0.3
N for cases/non-cases	37/3572	42/2566	57/3054	45/3064	41/3068	
Multivariable HR (95%CI)	1	1.21 (0.77 , 1.91)	1.39 (0.91 , 2.13)	1.17 (0.74 , 1.84)	1.35 (0.84 , 2.20)	
Red and processed meat						
All cancers						0.3
N for cases/non-cases	266/12026	339/11959	342/11961	344/11943	318/11978	
Multivariable HR (95%CI)	1	1.11 (0.94 , 1.30)	1.08 (0.91 , 1.27)	1.10 (0.93 , 1.30)	1.12 (0.94 , 1.33)	
Breast cancer						0.05
N for cases/non-cases	80/9105	101/9086	128/9065	126/9053	109/9077	
Multivariable HR (95%CI)	1	1.10 (0.82 , 1.49)	1.35 (1.02 , 1.81)	1.36 (1.02 , 1.81)	1.26 (0.93 , 1.71)	
Prostate cancer						0.8
N for cases/non-cases	37/3070	48/3063	54/3056	42/3066	41/3069	
Multivariable HR (95%CI)	1	1.07 (0.69 , 1.65)	1.21 (0.79 , 1.85)	0.93 (0.59 , 1.48)	1.17 (0.72 , 1.89)	

CI, confidence interval, HR, Hazard ratio

^a Multivariable models were adjusted for age (timescale), sex, energy intake without alcohol, number of 24h-dietary records, smoking status, educational level, physical activity, height, BMI, alcohol intake, family history of cancers, lipids intake, fruits, vegetables, menopausal status and number of children (breast cancer models), red meat intake (where processed meat was analyzed) and processed meat intake (where red meat was analyzed).

Sex-specific cut-offs for quintiles of red meat intake were 12.59; 37.14 ; 57.15 and 86.75 g/d in men and 0.14 ; 24.67 ; 42.15 and 65.71 g/d in women.

Sex-specific cut-offs for quintiles of processed meat intake were 0.20; 11.61 ; 25.45 and 45.86 g/d in men and 0.06 ; 5.36 ; 14.64 and 29.00 g/d in women.

Sex-specific cut-offs for quintiles of red and processed meat intake were 32.00; 59.82 ; 86.81 and 122.14 g/d in men and 18.21 ; 39.73 ; 60.00 and 87.68 g/d in women.

Table 3 Associations between quintiles of red and processed meat intake and breast cancer risk according to menopausal status from multivariable Cox proportional hazards models, NutriNet-Santé cohort, France, 2009 – 2016 (n=46474)^a

	Quintile 1	Quintile 2	Quintile3	Quintile 4	Quintile 5	p-trend
Red meat						
Pre-menopausal breast cancer						0.4
N for cases/non-cases	12/4732	50/4502	36/4618	43/4608	28/4622	
Multivariable HR (95%CI)	1	3.36 (1.77, 6.38)	2.37 (1.22, 4.60)	2.91 (1.52, 5.57)	2.04 (1.03, 4.06)	
Post-menopausal breast cancer						0.002
N for cases/non-cases	48/5347	73/5307	81/5308	78/5319	95/5281	
Multivariable HR (95%CI)	1	1.28 (0.88, 1.86)	1.46 (1.02, 2.09)	1.40 (0.97, 2.01)	1.79 (1.26, 2.55)	
Processed meat						
Pre-menopausal breast cancer						0.5
N for cases/non-cases	32/6591	28/2645	32/4619	40/4614	37/4613	
Multivariable HR (95%CI)	1	1.62 (0.96, 2.73)	1.09 (0.66, 1.80)	1.34 (0.83, 2.17)	1.30 (0.79, 2.15)	
Post-menopausal breast cancer						0.7
N for cases/non-cases	101/8309	36/2327	79/5299	93/5306	66/5321	
Multivariable HR (95%CI)	1	1.08 (0.73, 1.60)	1.07 (0.79, 1.44)	1.28 (0.95, 1.72)	0.95 (0.69, 1.32)	
Red and processed meat						
Pre-menopausal breast cancer						0.8
N for cases/non-cases	23/4609	36/4632	41/4612	40/4608	29/4621	
Multivariable HR (95%CI)	1	1.29 (0.76 , 2.19)	1.40 (0.83 , 2.36)	1.40 (0.83 , 2.37)	1.05 (0.59 , 1.86)	
Post-menopausal breast cancer						0.02
N for cases/non-cases	57/5331	66/5304	80/5325	88/5299	84/5303	
Multivariable HR (95%CI)	1	1.06 (0.74 , 1.52)	1.26 (0.90 , 1.77)	1.40 (0.99 , 1.96)	1.41 (0.99 , 2.01)	

^a Multivariable models were adjusted for age (timescale), energy intake without alcohol, number of 24h-dietary records, smoking status, educational level, physical activity, height, BMI, alcohol intake, family history of cancers, lipids intake, fruits, vegetables, hormone replacement therapy (for postmenopausal group), number of children, contraception (for premenopausal group), red meat intake (where processed meat was analyzed) and processed meat intake (where red meat was analyzed).

CI, confidence interval, HR, Hazard ratio

In premenopausal women: cut-offs for quintiles of red meat intake were 0.29; 24.00; 42.14; 67.7 g/d; cut-offs for quintiles of processed meat intake were 0.11; 6.79; 16.43; 31.89 g/d; cut-offs for quintiles of red and processed meat intake were 18.57; 40.40; 61.79; 91.16 g/d.

In postmenopausal women: cut-offs for quintiles of red meat intake were 2.68; 25.37; 42.68; 65.00 g/d; cut-offs for quintiles of processed meat intake were 0.06; 5.14; 14.29; 27.26 g/d; cut-offs for quintiles of red and processed meat intake were 18.21; 39.29; 58.79; 85.06 g/d.