



HAL
open science

The Aryl Hydrocarbon Receptor-Activating Effect of Uremic Toxins from Tryptophan Metabolism: A New Concept to Understand Cardiovascular Complications of Chronic Kidney Disease

Marion Sallée, Laetitia Dou, Claire Cerini, Stéphane Poitevin, Philippe Brunet, Stéphane Burtey

► To cite this version:

Marion Sallée, Laetitia Dou, Claire Cerini, Stéphane Poitevin, Philippe Brunet, et al.. The Aryl Hydrocarbon Receptor-Activating Effect of Uremic Toxins from Tryptophan Metabolism: A New Concept to Understand Cardiovascular Complications of Chronic Kidney Disease. *Toxins*, 2014, 6 (3), pp.934-949. 10.3390/toxins6030934 . hal-01610428

HAL Id: hal-01610428

<https://hal.science/hal-01610428v1>

Submitted on 31 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Review

The Aryl Hydrocarbon Receptor-Activating Effect of Uremic Toxins from Tryptophan Metabolism: A New Concept to Understand Cardiovascular Complications of Chronic Kidney Disease

Marion Sallée ^{1,2}, Laetitia Dou ^{1,*}, Claire Cerini ¹, Stéphane Poitevin ¹, Philippe Brunet ^{1,2} and Stéphane Burtey ^{1,2}

¹ Aix Marseille Université, Inserm, VRCM, UMR_S 1076, Marseille13005, France; E-Mails: marionsallee@yahoo.fr (M.S.); claire.cerini@univ-amu.fr (C.C.); stephane.poitevin@univ-amu.fr (S.P.); drphilippe.brunet@outlook.com (P.B.); stephane.burtey@ap-hm.fr (S.B.)

² Assistance Publique Hôpitaux de Marseille, CHU Conception, Centre de Néphrologie Dialyse Transplantation Rénale, Marseille 13005, France

* Author to whom correspondence should be addressed; E-Mail: laetitia.dou@univ-amu.fr; Tel.: +33-4-9183-5600; Fax: +33-4-9183-5602.

Received: 25 November 2013; in revised form: 19 February 2014 / Accepted: 24 February 2014 /

Published: 4 March 2014

Abstract: Patients with chronic kidney disease (CKD) have a higher risk of cardiovascular diseases and suffer from accelerated atherosclerosis. CKD patients are permanently exposed to uremic toxins, making them good candidates as pathogenic agents. We focus here on uremic toxins from tryptophan metabolism because of their potential involvement in cardiovascular toxicity: indolic uremic toxins (indoxyl sulfate, indole-3 acetic acid, and indoxyl- β -D-glucuronide) and uremic toxins from the kynurenine pathway (kynurenine, kynurenic acid, anthranilic acid, 3-hydroxykynurenine, 3-hydroxyanthranilic acid, and quinolinic acid). Uremic toxins derived from tryptophan are endogenous ligands of the transcription factor aryl hydrocarbon receptor (AhR). AhR, also known as the dioxin receptor, interacts with various regulatory and signaling proteins, including protein kinases and phosphatases, and Nuclear Factor-Kappa-B. AhR activation by 2,3,7,8-tetrachlorodibenzo-p-dioxin and some polychlorinated biphenyls is associated with an increase in cardiovascular disease in humans and in mice. In addition, this AhR activation mediates cardiotoxicity, vascular inflammation, and a procoagulant and

prooxidant phenotype of vascular cells. Uremic toxins derived from tryptophan have prooxidant, proinflammatory, procoagulant, and pro-apoptotic effects on cells involved in the cardiovascular system, and some of them are related with cardiovascular complications in CKD. We discuss here how the cardiovascular effects of these uremic toxins could be mediated by AhR activation, in a “dioxin-like” effect.

Keywords: tryptophan-derived uremic toxins; aryl hydrocarbon receptor; chronic kidney disease; cardiovascular diseases

1. Introduction

Patients with chronic kidney disease (CKD) are at high risk for cardiovascular diseases. CKD leads to accelerated atherosclerosis and consequently to a marked increase in cardiovascular morbi-mortality [1,2]. For patients with CKD, the risk of dying prematurely of cardiovascular disease is much higher than the risk of progressing to dialysis or transplantation [3]. This higher risk cannot be totally explained by classical cardiovascular risk factors such as diabetes mellitus, tobacco use, hypercholesterolemia, high blood pressure, and obesity. The uremic environment itself is harmful and uremic toxins have emerged as a key factor to explain cardiovascular disease [4]. These uremic toxins are classified in three groups by their behavior during dialysis [5]: the small water-soluble molecules, the middle molecules, and the protein-bound molecules, which are hardly removed by conventional hemodialysis treatments. Among uremic toxins, those derived from tryptophan are of particular interest because they have cardiovascular toxicity and they are Aryl Hydrocarbon Receptor (AhR) ligands [6,7]. These toxins include toxins from the kynurenine and indolic pathways. This paper reviews how toxins derived from tryptophan can play a role in cardiovascular diseases associated with CKD via their property of AhR activation.


2. Uremic Toxins from Tryptophan Metabolism

Tryptophan is an essential amino acid found in the diet. Ninety-five percent of tryptophan can be metabolized through the kynurenine metabolic pathway (Figure 1). This pathway is mediated by the rate-limiting enzymes tryptophan 2,3-dioxygenase (TDO) and indoleamine 2,3-dioxygenase (IDO) [8]. TDO is highly expressed in the liver and is also found in the brain. Two isoenzymes of IDO exist, IDO-1 and IDO-2. IDO-1 expression is found in most tissues [8]. IDO activity leading from tryptophan to kynurenine is reflected by the tryptophan/kynurenine (TRP/KYN) ratio. In CKD patients, tryptophan serum level is decreased whereas metabolites of the kynurenine pathway, *i.e.*, kynurenine, 3-hydroxykynurenine, kynurenic acid, anthranilic acid, and quinolinic acid, are increased [9,10].

Two other metabolic pathways of tryptophan metabolism are also described: the serotonin pathway that leads to melatonin, and the indolic pathway that leads to indolic components such as indoxyl sulfate (IS), indole acetic acid (IAA), and indoxyl- β -D glucuronide (IDG) (Figure 1) [7,11,12]. Indolic components are produced by intestinal bacteria via tryptophan degradation before absorption [12]. Indoles are metabolized to IS and IDG in the liver [13]. IAA is metabolized directly in the intestine [14]

or in tissue via tryptamine [11]. IS, IAA, and IDG are commonly excreted in urines and accumulate in case of CKD [5,15].

Figure 1. Uremic toxins from tryptophan metabolism.



3. AhR Activation by Toxins Derived from Tryptophan Metabolism

AhR is a ligand-activated transcription factor involved in biological detoxication of ligands. It is a member of the family of basic helix-loop-helix-Per-ARNT-Sim (PAS) containing transcription factors. AhR ligands include a wide variety of ubiquitous environmental pollutants like halogenated aromatic hydrocarbons (HAH) such as 2,3,7,8-tetrachlorodibenzo-*p*-dioxin or dioxin (TCDD), polycyclic aromatic hydrocarbons (PAH) and co-planar polychlorinated biphenyls (PCBs) [16,17]. However, there are many naturally occurring ligands such as tryptophan derivatives [17]. Historically, the action of AhR was determined in the setting of chemical carcinogenesis. AhR binds to the promoters of AhR target genes such as cytochromes P450 (CYP) 1A1, CYP1B1, and AhR repressor, which is the inhibitor protein of AhR. AhR induces the expression of xenobiotic metabolizing enzymes, such as CYP genes, needed for the detoxication of AhR toxic ligands [18]. Prior to AhR activation, AhR is found in the cytoplasm, coupled with its chaperone proteins such as Hsp90 (heat shock protein90).

When a ligand binds to AhR, the AhR/ligand/Hsp90 complex translocates into the nucleus and dimerizes with ARNT (Aryl hydrocarbon receptor nuclear translocator). The AhR/ARNT complex recognizes a DNA specific site, 5'-GCGTG-3', the XRE (Xenobiotic Responsive Element) sequence in promoters of target genes, and induces their transcription [19,20]. A non-genomic inflammatory pathway of AhR activation that leads to other transcription factor activation such as NF- κ B and AP-1 is also described [20]. The cytosolic AhR can activate a number of other cytosolic proteins, including proteins of the Mitogen-Activated Protein Kinase (MAPK) family Extracellular signal-regulated kinase (ERK), p38, and Jun-NH2-terminal kinase (JNK) [21–25]. AhR activation of MAPK depends on cell type. All three MAPKs phosphorylate downstream transcription factors and thereby alter gene expression to direct a specific cellular response. The AhR ligand TCDD induces ERK1/2 and p38 phosphorylation in rodent hepatoma cells [22] and in a mouse leukocyte cell line [23,24] and promotes JNK phosphorylation in a breast cancer cell line, MCF7 [25]. Furthermore, because the tyrosine kinase c-Src can be associated with cytosolic AhR complex, AhR activation by TCDD can also indirectly induce the MAPK cascade via c-Src [26,27]. C-Src phosphorylation results in EGFR phosphorylation, which triggers MAPK signal transduction and enhances the expression of target genes [27]. AhR also interacts with transcription factors such as AP-1 [24] and NF- κ B, notably the RelA subunit. The interaction between AhR and NF- κ B can lead to activation or mutual transrepression of their activities, depending on the duration, the cell type used, and the promoter being assayed [28,29]. These pathways were described in the setting of chemical carcinogenesis, their relevance in cardiovascular physiopathology has to be demonstrated.

Uremic toxins derived from tryptophan activate the AhR pathway. This was notably demonstrated for indolic toxins IS [6,30] and IAA [7,30], at uremic concentrations, in hepatic cells [6] and endothelial cells [30]. IS and IAA induce the nuclear translocation of AhR [30]. Using microarray analyses of endothelial cells, we found that IS and IAA induce an upregulation of eight genes regulated by AhR, including CYP1A1 and CYP1B1 [30]. The other uremic toxins derived from tryptophan metabolism described as AhR ligands are from the kynurenine pathway. In glioma cells, kynurenine induces AhR translocation into the nucleus and activates AhR target genes [31,32], but the concentrations of kynurenine used in those experiments are at least 10 times higher than those reported in uremic serum. For human NK cell lines, Shin *et al.* reported AhR activation by kynurenine [33] at uremic concentration [10,34]. Dinatale *et al.* found that kynurenic acid can stimulate the AhR pathway [35] at concentrations found in CKD patients [10,34].

4. Involvement of AhR-Activating Uremic Toxins in Cardiovascular Diseases

The cardiovascular toxicity of AhR ligands was first described with environmental pollutants like TCDD and co-planar PCBs. A recent meta-analysis highlighted a consistent association between TCDD exposure and increased risk of cardiovascular mortality, especially from ischemic diseases [36]. Moreover, people exposed to co-planar PCBs present an increased risk of acute myocardial infarction and ischemic stroke [37]. An association between plasma concentration of PCB serum level and hypertension was reported in the general population [38]. Furthermore, PCB levels in serum are strongly associated with the number of atherosclerotic plaques in carotid arteries [39]. In rodents, chronic TCDD or PCB exposure increases cardiomyopathy [40,41] and systemic blood pressure [41],

especially in female rats [42]. In an atherosclerosis model of apoE^{-/-} mice, PCB or TCDD exposure leads to a higher incidence of atherosclerotic lesions [43–45].

The role of AhR-activating uremic toxins in cardiovascular disease, and notably the role of IS, has been described. Serum IS is associated with overall and cardiovascular mortality in CKD patients (stage 2 to hemodialyzed stage 5) [46], and with all-cause mortality in incident hemodialyzed patients [47]. IS is associated with coronary lesions in non-hemodialyzed diabetic CKD patients even after eGFR adjustment [48]. IS is also associated with left ventricular diastolic dysfunction in CKD non-hemodialyzed patients [49]. In a model of salt-sensitive hypertensive rat, increased IS serum levels stimulate cardiac fibrosis [50]. The role of IS is supported by a decrease in myocardial fibrosis by AST-120, an oral charcoal adsorbent limiting indole absorption in the intestine, resulting in decreased IS serum levels [51]. In subtotaly nephrectomized apoE^{-/-} mice, AST-120 improves atherosclerotic lesions and reduces IS deposition in aorta [52].

Uremic toxins from the kynurenine pathway may also play a role in cardiovascular diseases. Higher levels of 3-hydroxykynurenine are most frequent in patients with cardiovascular diseases [53]. Interestingly, for patients with recent ischemic stroke, Darlington et al found an association between serum kynurenic acid levels and patient mortality at 24h [54]. In CKD patients, the TRP/KYN ratio, which inversely correlates with kidney function [34], is related to carotid intima-media thickness, a presymptomatic predictor of atherosclerosis [55,56]. In these patients, higher IDO activity is associated with larger carotid plaques [57].

5. AhR-Activating Uremic Toxins Induce Endothelial Dysfunction

Endothelial dysfunction is reflected by impaired endothelium-dependent vasodilatation. In vivo, flow-mediated dilatation (FMD) is a good clinical marker of endothelial dysfunction and is associated with cardiovascular events [58,59]. AhR activation by environmental ligands inhibits endothelium-dependent vasodilatory response via an induction of CYP1A1 in mice [41,60]. In CKD patients, the FMD decreases with kidney function [61], and one can suppose that AhR-activating uremic toxins impair the endothelium-dependent vasodilatation. Indeed, the decrease in IS plasma level after the administration of oral AST-120 in CKD patients is associated with improved FMD [62]. Furthermore, administration of AST-120 in rats with subtotal nephrectomy improves acetylcholine-dependent vasodilatation of the aorta, in a dose-dependent manner [63].

Environmental pollutants TCDD and co-planar PCBs that induce AhR activation have deleterious effects on endothelial cells *in vitro*. The AhR agonist 3-methylcholanthrene inhibits endothelial cell adhesion, migration, and proliferation [64] through p38 MAPK phosphorylation; PCB126 decreases endothelial nitric oxide (NO) production [65]; 3-methylcholanthrene decreases endothelial stress fiber formation [66]; and PAHs decrease endothelial progenitor cell number and colonies [67]. All these mechanisms are also observed for uremic toxins derived from tryptophan. In vitro, IS inhibits endothelial proliferation and wound repair [68] and increases cell senescence [62]. IS also inhibits endothelial cell migration by a tardive inhibition of ERK1/2 phosphorylation [69]. IS decreases endothelial NO production [62], and the inhibition of endothelial cell migration is reversed by NO [69]. IS induces the endothelial cell adherens junction disassembly and stress fiber reorganization mediated by ERK1/2 phosphorylation that is induced by oxidative stress overproduction [70]. In CKD patients,

IAA serum level is negatively correlated with the number of immature progenitor cells [71], and IAA induces progenitor cell apoptosis *in vitro* [71].

6. AhR-Activating Uremic Toxins Increase Oxidative Stress in Cardiovascular Cells

The role of AhR activation in oxidative stress is supported by studies demonstrating that AhR activation by TCDD and PCB induces reactive oxygen species (ROS) production [41,72–74] in the aorta, heart, and kidney [41,60] and in endothelial cells [73] via CYP1A1 expression [65,72,74].

In CKD patients, high IS [62], kynurenine [53], 3-hydroxykynurenine [53], and KYN/TRP ratio [75] are all associated with oxidative stress markers. Furthermore, in a model of salt-sensitive hypertensive rat, increased IS serum levels enhance cardiac oxidative stress [50]. In CKD patients, administration of AST-120, which reduces IS serum level, decreases plasma levels of oxidative stress markers [62]. AST-120 improves aortic [63] and cardiac [51] oxidative/nitrative stress in CKD rat models. *In vitro*, AhR-activating indolic uremic toxins such as IDG [76], IS [77], and IAA [personal data] induce endothelial ROS production. Induction of endothelial ROS production by IS occurs through activation of NADPH oxidase and inhibition of antioxidant defense mechanisms [76,77]. The link between a pro-oxidant effect of IS and AhR activation is supported by experiments demonstrating that the inductions of both Nox4 gene expression [78] and superoxide production by IS are inhibited by the AhR inhibitors ANF and CH-223191 [78].

In vascular smooth muscle cells (VSMC), IS stimulates NADPH oxidase-dependent ROS generation, promotes osteoblast differentiation [79], and induces VSMC senescence through oxidative stress pathway [80,81]. IS promotes the migration of VSMCs by a mechanism that involves ROS production [82,83] and phosphorylation of ERK1/2 [82,84] and p38 [82]. The involvement of AhR activation in the effects of IS on VSMC has not been studied. However, effects of IS mimic those of the AhR agonist Benzo [a]pyrene, which promotes VSMC migration [85] and ROS production [85,86] in an AhR-dependent manner.

7. AhR-Activating Uremic Toxins Induce Leukocyte Activation and Inflammation

The involvement of AhR activation in inflammation was demonstrated using environmental pollutants. In mice, PCB and TCDD exposure induces the expression of proinflammatory cytokines [45] and aortic expression of monocyte chemoattractant protein-1 (MCP-1) [87,88], but not in mice invalidated for AhR nuclear translocation [88]. In T cell immune response, TCDD induces lymphocyte activation via modifications in dendritic cell phenotype and functions [89]. Furthermore, AhR participates in T cell differentiation [90]. Exposure to AhR-activating environmental pollutants increases monocyte adhesion [91,92]. Macrophages treated with TCDD show an up-regulation of inflammatory genes IL-8 and MMP-12 and of VEGF in an AhR- and IL-8-dependent manner [45]. *In vitro*, AhR activation by PCB induces MCP-1 expression [73,87,91,93] through a ROS/NF- κ B pathway [73]. PCB induces the expression of endothelial adhesion molecules E-selectin [91,94], VCAM-1 [92,94], and ICAM-1 [89,91,95] via MAPK signaling pathways [92,95]. Finally, AhR agonists increase endothelial COX-2 expression and COX-2 metabolites in urine [43,65].

A growing body of data supports the hypothesis that AhR activation by tryptophan-derived uremic toxins is involved in inflammation associated with CKD. Kynurenine and 3-hydroxykynurenine levels

correlate with CRP in dialysis patients [53], and KYN/TRP ratio is positively associated with markers of inflammation in CKD patients [75]. Kynurenine and its metabolites induce T cell differentiation [96], and kynurenic acid activates IL6 in MCF-7 cells in an AhR-dependent manner [35]. IS stimulates inflammatory pathways, notably the p38/NF- κ B pathway that leads to IL-6 and IL-1 β expression [97,98]. IS enhances leukocyte adhesion in a subtotal nephrectomy mouse model after vascular lesion [99]. IS also enhances leukocyte adhesion to the endothelium *in vitro* by a ROS/p38 MAPK pathway [98] and a ROS/JNK/NF- κ B pathway [99]. In monocytes, IS induces ROS production in an NADPH oxidase-dependent manner [98]. Furthermore, IS induces endothelial E-selectin [99], VCAM-1 [100], and ICAM-1 [100,101] expression. IS also induces the expression of MCP-1 [78,100–102], which is abolished by an AhR antagonist [78]. MCP-1 induction by IS involves a ROS/MAPK/NF- κ B pathway [101,102], which evokes the involvement of the inflammatory non-genomic pathway of AhR. Finally, we have observed that IS and IAA induce COX-2 expression in endothelial cells [30] in an AhR-dependent manner [103].

8. Involvement of AhR-Activating Uremic Toxins in Thrombosis

AhR activation by tryptophan-derived uremic toxins could be involved in thrombosis. In hemodialyzed patients, the KYN/TRP ratio is positively associated with thrombosis markers [75], and IS level correlates with vascular access intervention [104]. In CKD patients, the levels of the indolic uremic toxins IS and IAA correlate with the levels and activity of tissue factor [30], the main initiator of the blood coagulation cascade. In endothelial cells and in peripheral blood mononuclear cells, IS and IAA increase tissue factor expression and tissue factor-dependent procoagulant activity [30]. Using siRNA and a pharmacological AhR inhibitor, we demonstrated that AhR activation is involved in tissue factor production induced by indolic uremic toxins [30]; we also showed that the canonical AhR agonist TCDD induces endothelial tissue factor expression [30]. The increase in tissue factor activity induced by IS and IAA is associated with enhanced clot formation by a pathway encompassing inhibition of tissue factor ubiquitination, which increases tissue factor half-life [105]. Interestingly, AHR was reported to interact with ubiquitination machinery [106].

9. Is AhR Activation Involved in CardioVascular Toxicity of CKD?

AhR can be activated by structurally diverse chemicals. The dramatic variety of effects produced by TCDD and other AhR ligands suggests significant mechanistic diversity in the action of these chemicals [20]. There are species- and tissue-specific toxic effects [20]. AhR-dependent toxic and biological effects are modulated by the structure and metabolism of AhR ligands leading to differences in AhR-dependent gene expression profiles [107]. TCDD results in a persistent activation of AhR due to its accumulation in adipose tissue [44,108] whereas other AhR ligands such as benzo [a]pyrene are rapidly metabolized and fail to induce sustained AhR activation. Little is known about the metabolism and the tissue distribution of tryptophan-derived uremic toxins in CKD. However the accumulation of these uremic toxins in patients suggests the possibility of a persistent activation of AhR. The hypothesis that AhR activation is involved in the cardiovascular toxicity of tryptophan-derived uremic toxins has to be confirmed in animal and human studies.

10. Conclusions

Tryptophan-derived uremic toxins share, with environmental pollutants like dioxins, the ability to activate the AhR pathway. These toxins, like environmental pollutants, induce endothelial dysfunction and leukocyte activation, favor inflammation and thrombosis, and increase vascular oxidative stress (Table 1). The ability of tryptophan derived of uremic toxins to activate AhR may explain how these toxins contribute to cardiovascular diseases in CKD patients. These recent insights into the mechanisms of toxicity of tryptophan-derived uremic toxins may lead to new therapeutic strategies targeting AhR activation.

Table 1. Similarities between the effects of aryl hydrocarbon receptor (AhR)-activating pollutants and the effects of tryptophan-derived uremic toxins.

Effects of AhR-activating-pollutants	Effects of tryptophan-derived uremic toxins
Association with cardiovascular events	
Association with cardiovascular mortality [36]	Association with cardiovascular mortality [46]
Association with higher hypertension [37]	Association with overall mortality [46,47]
Association with hospitalization for ischemic stroke [37]	Association with the mortality of ischemic stroke [54]
Associated with number of atherosclerotic carotid plaques [39]	Association with greater size of atherosclerotic carotid plaques [57] Association with coronary lesions [48]
Decrease in endothelial-dependent vasodilation [41,60]	Decrease in endothelial-dependent vasodilation [62,63]
Induction of cardiomyopathy in rodents [40,41]	Induction of cardiac fibrosis in rodents [50,51]
Higher incidence of atherosclerotic lesions in mice [43–45]	Induction of atherosclerotic lesions in mice [52]
Endothelial dysfunction	
Inhibition of endothelial cell proliferation [64]	Inhibition of endothelial cell proliferation [62,68]
Inhibition of endothelial NO production [65]	Inhibition of endothelial NO production [62]
Inhibition of endothelial cell migration [64]	Inhibition of endothelial cell migration [69]
Decrease in endothelial progenitor cells [67]	Decrease in progenitor cells [71]
Induction of stress fiber formation [66]	Induction of stress fiber reorganization [70]
Oxidative stress	
Association with oxidative stress markers [41,60,72,73]	Association with oxidative stress markers [53,62,75]
Induction of endothelial ROS [65,72–74]	Induction of endothelial ROS [76,77]
Induction of VSMC migration by ROS [85,86]	Induction of VSMC migration by ROS [82,83]
Thrombosis	
Induction of TF production and activity [30]	Induction of TF production and activity [30]
Inflammation	
Association with inflammatory markers [45,88]	Association with inflammatory markers [53,75]
Increase in monocyte expression of inflammatory cytokines [45,89]	Increase in monocyte expression of inflammatory cytokines [97,98]
Increase in monocyte adhesion [92,94]	Increase in monocyte adhesion [98,99]
Induction of VCAM-1 [92,100], MCP-1 [73,87,91,93], E-selectin [91,94] ICAM-1 [89,91,95] and COX-2 [65]	Induction of VCAM-1 [100], MCP-1 [78,100–102], E-selectin [99], ICAM-1 [100,101] and COX-2 [30]
Induction of urinary COX-2 metabolites in mice [43]	
Induction of T cell differentiation [90]	Induction of T cell differentiation [96]

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Vanholder, R.; Massy, Z.; Argiles, A.; Spasovski, G.; Verbeke, F.; Lameire, N. Chronic kidney disease as cause of cardiovascular morbidity and mortality. *Nephrol. Dial. Transplant.* **2005**, *20*, 1048–1056.
2. Go, A.S.; Chertow, G.M.; Fan, D.; McCulloch, C.E.; Hsu, C.Y. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N. Engl. J. Med.* **2004**, *351*, 1296–1305.
3. Keith, D.S.; Nichols, G.A.; Gullion, C.M.; Brown, J.B.; Smith, D.H. Longitudinal follow-up and outcomes among a population with chronic kidney disease in a large managed care organization. *Arch. Intern. Med.* **2004**, *164*, 659–663.
4. Jourde-Chiche, N.; Dou, L.; Cerini, C.; Dignat-George, F.; Brunet, P. Vascular incompetence in dialysis patients—Protein-bound uremic toxins and endothelial dysfunction. *Semin. Dial.* **2011**, *24*, 327–337.
5. Vanholder, R.; De Smet, R.; Glorieux, G.; Argiles, A.; Baurmeister, U.; Brunet, P.; Clark, W.; Cohen, G.; De Deyn, P.P.; Deppisch, R.; *et al.* Review on uremic toxins: Classification, concentration, and interindividual variability. *Kidney Int.* **2003**, *63*, 1934–1943.
6. Schroeder, J.C.; Dinatale, B.C.; Murray, I.A.; Flaveny, C.A.; Liu, Q.; Laurenzana, E.M.; Lin, J.M.; Strom, S.C.; Omiecinski, C.J.; Amin, S.; *et al.* The uremic toxin 3-indoxyl sulfate is a potent endogenous agonist for the human aryl hydrocarbon receptor. *Biochemistry* **2010**, *49*, 393–400.
7. Heath-Pagliuso, S.; Rogers, W.J.; Tullis, K.; Seidel, S.D.; Cenijn, P.H.; Brouwer, A.; Denison, M.S. Activation of the Ah receptor by tryptophan and tryptophan metabolites. *Biochemistry* **1998**, *37*, 11508–11515.
8. Fatokun, A.A.; Hunt, N.H.; Ball, H.J. Indoleamine 2,3-dioxygenase 2 (IDO2) and the kynurenine pathway: Characteristics and potential roles in health and disease. *Amino Acids* **2013**, *45*, 1319–1329.
9. Saito, K.; Fujigaki, S.; Heyes, M.P.; Shibata, K.; Takemura, M.; Fujii, H.; Wada, H.; Noma, A.; Seishima, M. Mechanism of increases in L-kynurenine and quinolinic acid in renal insufficiency. *Am. J. Physiol. Renal Physiol.* **2000**, *279*, F565–F572.
10. Pawlak, D.; Pawlak, K.; Malyszko, J.; Mysliwiec, M.; Buczko, W. Accumulation of toxic products degradation of kynurenine in hemodialyzed patients. *Int. Urol. Nephrol.* **2001**, *33*, 399–404.
11. Zhu, W.; Stevens, A.P.; Dettmer, K.; Gottfried, E.; Hoves, S.; Kreutz, M.; Holler, E.; Canelas, A.B.; Kema, I.; Oefner, P.J. Quantitative profiling of tryptophan metabolites in serum, urine, and cell culture supernatants by liquid chromatography-tandem mass spectrometry. *Anal. Bioanal. Chem.* **2011**, *401*, 3249–3261.
12. Aronov, P.A.; Luo, F.J.; Plummer, N.S.; Quan, Z.; Holmes, S.; Hostetter, T.H.; Meyer, T.W. Colonic contribution to uremic solutes. *J. Am. Soc. Nephrol.* **2011**, *22*, 1769–1776.
13. Meyer, T.W.; Hostetter, T.H. Uremic solutes from colon microbes. *Kidney Int.* **2012**, *81*, 949–954

14. Koga, J.; Syono, K.; Ichikawa, T.; Adachi, T. Involvement of L-tryptophan aminotransferase in indole-3-acetic acid biosynthesis in *Enterobacter cloacae*. *Biochim. Biophys. Acta* **1994**, *1209*, 241–247.
15. Duranton, F.; Cohen, G.; De Smet, R.; Rodriguez, M.; Jankowski, J.; Vanholder, R.; Argiles, A. Normal and pathologic concentrations of uremic toxins. *J. Am. Soc. Nephrol.* **2012**, *23*, 1258–1270.
16. Bock, K.W. The human Ah receptor: Hints from dioxin toxicities to deregulated target genes and physiological functions. *Biol. Chem.* **2013**, *394*, 729–739.
17. Denison, M.S.; Nagy, S.R. Activation of the aryl hydrocarbon receptor by structurally diverse exogenous and endogenous chemicals. *Annu. Rev. Pharmacol. Toxicol.* **2003**, *43*, 309–334.
18. Schmidt, J.V.; Bradfield, C.A. Ah receptor signaling pathways. *Annu. Rev. Cell. Dev. Biol.* **1996**, *12*, 55–89.
19. Fisher, J.M.; Jones, K.W.; Whitlock, J.P., Jr. Activation of transcription as a general mechanism of 2,3,7,8-tetrachlorodibenzo-p-dioxin action. *Mol. Carcinog.* **1989**, *1*, 216–221.
20. Denison, M.S.; Soshilov, A.A.; He, G.; DeGroot, D.E.; Zhao, B. Exactly the same but different: Promiscuity and diversity in the molecular mechanisms of action of the aryl hydrocarbon (dioxin) receptor. *Toxicol. Sci.* **2011**, *124*, 1–22.
21. Matsumura, F. The significance of the nongenomic pathway in mediating inflammatory signaling of the dioxin-activated Ah receptor to cause toxic effects. *Biochem. Pharmacol.* **2009**, *77*, 608–626.
22. Tan, Z.; Chang, X.; Puga, A.; Xia, Y. Activation of mitogen-activated protein kinases (MAPKs) by aromatic hydrocarbons: Role in the regulation of aryl hydrocarbon receptor (AHR) function. *Biochem. Pharmacol.* **2002**, *64*, 771–780.
23. Park, S.J.; Yoon, W.K.; Kim, H.J.; Son, H.Y.; Cho, S.W.; Jeong, K.S.; Kim, T.H.; Kim, S.H.; Kim, S.R.; Ryu, S.Y. 2,3,7,8-Tetrachlorodibenzo-p-dioxin activates ERK and p38 mitogen-activated protein kinases in RAW 264.7 cells. *Anticancer Res.* **2005**, *25*, 2831–2836.
24. Weiss, C.; Faust, D.; Durk, H.; Kolluri, S.K.; Pelzer, A.; Schneider, S.; Dietrich, C.; Oesch, F.; Gottlicher, M. TCDD induces c-jun expression via a novel Ah (dioxin) receptor-mediated p38-MAPK-dependent pathway. *Oncogene* **2005**, *24*, 4975–4983.
25. Diry, M.; Tomkiewicz, C.; Koehle, C.; Coumoul, X.; Bock, K.W.; Barouki, R.; Transy, C. Activation of the dioxin/aryl hydrocarbon receptor (AhR) modulates cell plasticity through a JNK-dependent mechanism. *Oncogene* **2006**, *25*, 5570–5574.
26. Park, S.; Mazina, O.; Kitagawa, A.; Wong, P.; Matsumura, F. TCDD causes suppression of growth and differentiation of MCF10A, human mammary epithelial cells by interfering with their insulin receptor signaling through c-Src kinase and ERK activation. *J. Biochem. Mol. Toxicol.* **2004**, *18*, 322–331.
27. Xie, G.; Peng, Z.; Raufman, J.P. Src-mediated aryl hydrocarbon and epidermal growth factor receptor cross talk stimulates colon cancer cell proliferation. *Am. J. Physiol. Gastrointest. Liver Physiol.* **2012**, *302*, G1006–G1015.
28. Tian, Y.; Rabson, A.B.; Gallo, M.A. Ah receptor and NF-kappaB interactions: mechanisms and physiological implications. *Chem. Biol. Interact.* **2002**, *141*, 97–115.
29. Vogel, C.F.; Sciallo, E.; Li, W.; Wong, P.; Lazennec, G.; Matsumura, F. RelB, a new partner of aryl hydrocarbon receptor-mediated transcription. *Mol. Endocrinol.* **2007**, *21*, 2941–2955.

30. Gondouin, B.; Cerini, C.; Dou, L.; Sallee, M.; Duval-Sabatier, A.; Pletinck, A.; Calaf, R.; Lacroix, R.; Jourde-Chiche, N.; Poitevin, S.; *et al.* Indolic uremic solutes increase tissue factor production in endothelial cells by the aryl hydrocarbon receptor pathway. *Kidney Int.* **2013**, *84*, 733–744.
31. Frumento, G.; Rotondo, R.; Tonetti, M.; Damonte, G.; Benatti, U.; Ferrara, G.B. Tryptophan-derived catabolites are responsible for inhibition of T and natural killer cell proliferation induced by indoleamine 2,3-dioxygenase. *J. Exp. Med.* **2002**, *196*, 459–468.
32. Opitz, C.A.; Litzenburger, U.M.; Sahm, F.; Ott, M.; Tritschler, I.; Trump, S.; Schumacher, T.; Jestaedt, L.; Schrenk, D.; Weller, M.; *et al.* An endogenous tumour-promoting ligand of the human aryl hydrocarbon receptor. *Nature* **2011**, *478*, 197–203.
33. Shin, J.H.; Zhang, L.; Murillo-Sauca, O.; Kim, J.; Kohrt, H.E.; Bui, J.D.; Sunwoo, J.B. Modulation of natural killer cell antitumor activity by the aryl hydrocarbon receptor. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 12391–12396.
34. Schefold, J.C.; Zeden, J.P.; Fotopoulou, C.; von Haehling, S.; Pschowski, R.; Hasper, D.; Volk, H.D.; Schuett, C.; Reinke, P. Increased indoleamine 2,3-dioxygenase (IDO) activity and elevated serum levels of tryptophan catabolites in patients with chronic kidney disease: A possible link between chronic inflammation and uraemic symptoms. *Nephrol. Dial. Transplant.* **2009**, *24*, 1901–1908.
35. DiNatale, B.C.; Murray, I.A.; Schroeder, J.C.; Flaveny, C.A.; Lahoti, T.S.; Laurenzana, E.M.; Omiecinski, C.J.; Perdew, G.H. Kynurenic acid is a potent endogenous aryl hydrocarbon receptor ligand that synergistically induces interleukin-6 in the presence of inflammatory signaling. *Toxicol. Sci.* **2010**, *115*, 89–97.
36. Humblet, O.; Birnbaum, L.; Rimm, E.; Mittleman, M.A.; Hauser, R. Dioxins and cardiovascular disease mortality. *Environ. Health Perspect* **2008**, *116*, 1443–1448.
37. Institute of Medicine of the National Academies. *Veterans and Agent Orange: Update 2012*; The National Academies Press: Washington, DC, USA, 2013.
38. Everett, C.J.; Mainous, A.G., 3rd; Frithsen, I.L.; Player, M.S.; Matheson, E.M. Association of polychlorinated biphenyls with hypertension in the 1999–2002 national health and nutrition examination survey. *Environ. Res.* **2008**, *108*, 94–97.
39. Lind, P.M.; van Bavel, B.; Salihovic, S.; Lind, L. Circulating levels of persistent organic pollutants (POPs) and carotid atherosclerosis in the elderly. *Environ. Health Perspect* **2012**, *120*, 38–43.
40. Jokinen, M.P.; Walker, N.J.; Brix, A.E.; Sells, D.M.; Haseman, J.K.; Nyska, A. Increase in cardiovascular pathology in female Sprague-Dawley rats following chronic treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin and 3,3',4,4',5-pentachlorobiphenyl. *Cardiovasc. Toxicol.* **2003**, *3*, 299–310.
41. Kopf, P.G.; Huwe, J.K.; Walker, M.K. Hypertension, cardiac hypertrophy, and impaired vascular relaxation induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin are associated with increased superoxide. *Cardiovasc. Toxicol.* **2008**, *8*, 181–193.
42. Lind, P.M.; Orberg, J.; Edlund, U.B.; Sjoblom, L.; Lind, L. The dioxin-like pollutant PCB 126 (3,3',4,4',5-pentachlorobiphenyl) affects risk factors for cardiovascular disease in female rats. *Toxicol. Lett.* **2004**, *150*, 293–299.

43. Dalton, T.P.; Kerzee, J.K.; Wang, B.; Miller, M.; Dieter, M.Z.; Lorenz, J.N.; Shertzer, H.G.; Nerbert, D.W.; Puga, A. Dioxin exposure is an environmental risk factor for ischemic heart disease. *Cardiovasc. Toxicol.* **2001**, *1*, 285–298.
44. Arsenescu, V.; Arsenescu, R.; Parulkar, M.; Karounos, M.; Zhang, X.; Baker, N.; Cassis, L.A. Polychlorinated biphenyl 77 augments angiotensin II-induced atherosclerosis and abdominal aortic aneurysms in male apolipoprotein E deficient mice. *Toxicol. Appl. Pharmacol.* **2011**, *257*, 148–154.
45. Wu, D.; Nishimura, N.; Kuo, V.; Fiehn, O.; Shahbaz, S.; Van Winkle, L.; Matsumura, F.; Vogel, C.F. Activation of aryl hydrocarbon receptor induces vascular inflammation and promotes atherosclerosis in apolipoprotein E^{-/-} mice. *Arterioscler. Thromb. Vasc. Biol.* **2011**, *31*, 1260–1267.
46. Barreto, F.C.; Barreto, D.V.; Liabeuf, S.; Meert, N.; Glorieux, G.; Temmar, M.; Choukroun, G.; Vanholder, R.; Massy, Z.A. Serum indoxyl sulfate is associated with vascular disease and mortality in chronic kidney disease patients. *Clin. J. Am. Soc. Nephrol.* **2009**, *4*, 1551–1558.
47. Melamed, M.L.; Plantinga, L.; Shafi, T.; Parekh, R.; Meyer, T.W.; Hostetter, T.H.; Coresh, J.; Powe, N.R. Retained organic solutes, patient characteristics and all-cause and cardiovascular mortality in hemodialysis: Results from the retained organic solutes and clinical outcomes (ROSCO) investigators. *BMC Nephrol.* **2013**, *14*, 134.
48. Chiu, C.A.; Lu, L.F.; Yu, T.H.; Hung, W.C.; Chung, F.M.; Tsai, I.T.; Yang, C.Y.; Hsu, C.C.; Lu, Y.C.; Wang, C.P.; *et al.* Increased levels of total P-Cresylsulphate and indoxyl sulphate are associated with coronary artery disease in patients with diabetic nephropathy. *Rev. Diabet. Stud.* **2010**, *7*, 275–284.
49. Sato, B.; Yoshikawa, D.; Ishii, H.; Suzuki, S.; Inoue, Y.; Takeshita, K.; Tanaka, M.; Kumagai, S.; Matsumoto, M.; Okumura, S.; *et al.* Relation of plasma indoxyl sulfate levels and estimated glomerular filtration rate to left ventricular diastolic dysfunction. *Am. J. Cardiol.* **2013**, *111*, 712–716.
50. Yisireyili, M.; Shimizu, H.; Saito, S.; Enomoto, A.; Nishijima, F.; Niwa, T. Indoxyl sulfate promotes cardiac fibrosis with enhanced oxidative stress in hypertensive rats. *Life Sci.* **2013**, *92*, 1180–1185.
51. Fujii, H.; Nishijima, F.; Goto, S.; Sugano, M.; Yamato, H.; Kitazawa, R.; Kitazawa, S.; Fukagawa, M. Oral charcoal adsorbent (AST-120) prevents progression of cardiac damage in chronic kidney disease through suppression of oxidative stress. *Nephrol. Dial. Transplant.* **2009**, *24*, 2089–2095.
52. Yamamoto, S.; Zuo, Y.; Ma, J.; Yancey, P.G.; Hunley, T.E.; Motojima, M.; Fogo, A.B.; Linton, M.F.; Fazio, S.; Ichikawa, I.; *et al.* Oral activated charcoal adsorbent (AST-120) ameliorates extent and instability of atherosclerosis accelerated by kidney disease in apolipoprotein E-deficient mice. *Nephrol. Dial. Transplant.* **2011**, *26*, 2491–2497.
53. Pawlak, K.; Domaniewski, T.; Mysliwiec, M.; Pawlak, D. The kynurenines are associated with oxidative stress, inflammation and the prevalence of cardiovascular disease in patients with end-stage renal disease. *Atherosclerosis* **2009**, *204*, 309–314.
54. Darlington, L.G.; Mackay, G.M.; Forrest, C.M.; Stoy, N.; George, C.; Stone, T.W. Altered kynurenine metabolism correlates with infarct volume in stroke. *Eur. J. Neurosci.* **2007**, *26*, 2211–2221.

55. Pertovaara, M.; Raitala, A.; Juonala, M.; Lehtimäki, T.; Huhtala, H.; Oja, S.S.; Jokinen, E.; Viikari, J.S.; Raitakari, O.T.; Hurme, M. Indoleamine 2,3-dioxygenase enzyme activity correlates with risk factors for atherosclerosis: The cardiovascular risk in young finns study. *Clin. Exp. Immunol.* **2007**, *148*, 106–111.
56. Niinisalo, P.; Raitala, A.; Pertovaara, M.; Oja, S.S.; Lehtimäki, T.; Kahonen, M.; Reunanen, A.; Jula, A.; Moilanen, L.; Kesaniemi, Y.A.; *et al.* Indoleamine 2,3-dioxygenase activity associates with cardiovascular risk factors: The health 2000 study. *Scand. J. Clin. Lab. Invest.* **2008**, *68*, 767–770.
57. Kato, A.; Suzuki, Y.; Suda, T.; Suzuki, M.; Fujie, M.; Takita, T.; Furuhashi, M.; Maruyama, Y.; Chida, K.; Hishida, A. Relationship between an increased serum kynurenine/tryptophan ratio and atherosclerotic parameters in hemodialysis patients. *Hemodial. Int.* **2010**, *14*, 418–424.
58. Gokce, N.; Keaney, J.F., Jr.; Hunter, L.M.; Watkins, M.T.; Menzoian, J.O.; Vita, J.A. Risk stratification for postoperative cardiovascular events via noninvasive assessment of endothelial function: A prospective study. *Circulation* **2002**, *105*, 1567–1572.
59. Gokce, N.; Keaney, J.F., Jr.; Hunter, L.M.; Watkins, M.T.; Nedeljkovic, Z.S.; Menzoian, J.O.; Vita, J.A. Predictive value of noninvasively determined endothelial dysfunction for long-term cardiovascular events in patients with peripheral vascular disease. *J. Am. Coll. Cardiol.* **2003**, *41*, 1769–1775.
60. Kopf, P.G.; Scott, J.A.; Agbor, L.N.; Boberg, J.R.; Elased, K.M.; Huwe, J.K.; Walker, M.K. Cytochrome P4501A1 is required for vascular dysfunction and hypertension induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol. Sci.* **2010**, *117*, 537–546.
61. Perticone, F.; Maio, R.; Tripepi, G.; Zoccali, C. Endothelial dysfunction and mild renal insufficiency in essential hypertension. *Circulation* **2004**, *110*, 821–825.
62. Yu, M.; Kim, Y.J.; Kang, D.H. Indoxyl sulfate-induced endothelial dysfunction in patients with chronic kidney disease via an induction of oxidative stress. *Clin. J. Am. Soc. Nephrol.* **2011**, *6*, 30–39.
63. Namikoshi, T.; Tomita, N.; Satoh, M.; Sakuta, T.; Kuwabara, A.; Kobayashi, S.; Higuchi, Y.; Nishijima, F.; Kashihara, N. Oral adsorbent AST-120 ameliorates endothelial dysfunction independent of renal function in rats with subtotal nephrectomy. *Hypertens. Res.* **2009**, *32*, 194–200.
64. Juan, S.H.; Lee, J.L.; Ho, P.Y.; Lee, Y.H.; Lee, W.S. Antiproliferative and antiangiogenic effects of 3-methylcholanthrene, an aryl-hydrocarbon receptor agonist, in human umbilical vascular endothelial cells. *Eur. J. Pharmacol.* **2006**, *530*, 1–8.
65. Andersson, H.; Garscha, U.; Brittebo, E. Effects of PCB126 and 17beta-oestradiol on endothelium-derived vasoactive factors in human endothelial cells. *Toxicology* **2011**, *285*, 46–56.
66. Chang, C.C.; Tsai, S.Y.; Lin, H.; Li, H.F.; Lee, Y.H.; Chou, Y.; Jen, C.Y.; Juan, S.H. Aryl-hydrocarbon receptor-dependent alteration of FAK/RhoA in the inhibition of HUVEC motility by 3-methylcholanthrene. *Cell. Mol. Life Sci.* **2009**, *66*, 3193–3205.
67. Van Grevenynghe, J.; Monteiro, P.; Gilot, D.; Fest, T.; Fardel, O. Human endothelial progenitors constitute targets for environmental atherogenic polycyclic aromatic hydrocarbons. *Biochem. Biophys. Res. Commun.* **2006**, *341*, 763–769.
68. Dou, L.; Bertrand, E.; Cerini, C.; Faure, V.; Sampol, J.; Vanholder, R.; Berland, Y.; Brunet, P. The uremic solutes p-cresol and indoxyl sulfate inhibit endothelial proliferation and wound repair. *Kidney Int.* **2004**, *65*, 442–451.

69. Kharait, S.; Haddad, D.J.; Springer, M.L. Nitric oxide counters the inhibitory effects of uremic toxin indoxyl sulfate on endothelial cells by governing ERK MAP kinase and myosin light chain activation. *Biochem. Biophys. Res. Commun.* **2011**, *409*, 758–763.
70. Peng, Y.S.; Lin, Y.T.; Chen, Y.; Hung, K.Y.; Wang, S.M. Effects of indoxyl sulfate on adherens junctions of endothelial cells and the underlying signaling mechanism. *J. Cell. Biochem.* **2012**, *113*, 1034–1043.
71. Jourde-Chiche, N.; Dou, L.; Sabatier, F.; Calaf, R.; Cerini, C.; Robert, S.; Camoin-Jau, L.; Charpiot, P.; Argiles, A.; Dignat-George, F.; *et al.* Levels of circulating endothelial progenitor cells are related to uremic toxins and vascular injury in hemodialysis patients. *J. Thromb. Haemost.* **2009**, *7*, 1576–1584.
72. Slim, R.; Toborek, M.; Robertson, L.W.; Hennig, B. Antioxidant protection against PCB-mediated endothelial cell activation. *Toxicol. Sci.* **1999**, *52*, 232–239.
73. Majkova, Z.; Layne, J.; Sunkara, M.; Morris, A.J.; Toborek, M.; Hennig, B. Omega-3 fatty acid oxidation products prevent vascular endothelial cell activation by coplanar polychlorinated biphenyls. *Toxicol. Appl. Pharmacol.* **2011**, *251*, 41–49.
74. Kopf, P.G.; Walker, M.K. 2,3,7,8-tetrachlorodibenzo-p-dioxin increases reactive oxygen species production in human endothelial cells via induction of cytochrome P4501A1. *Toxicol. Appl. Pharmacol.* **2010**, *245*, 91–99.
75. Pawlak, K.; Domaniewski, T.; Mysliwiec, M.; Pawlak, D. Kynurenines and oxidative status are independently associated with thrombomodulin and von Willebrand factor levels in patients with end-stage renal disease. *Thromb. Res.* **2009**, *124*, 452–457.
76. Itoh, Y.; Ezawa, A.; Kikuchi, K.; Tsuruta, Y.; Niwa, T. Protein-bound uremic toxins in hemodialysis patients measured by liquid chromatography/tandem mass spectrometry and their effects on endothelial ROS production. *Anal. Bioanal. Chem.* **2012**, *403*, 1841–1850.
77. Dou, L.; Jourde-Chiche, N.; Faure, V.; Cerini, C.; Berland, Y.; Dignat-George, F.; Brunet, P. The uremic solute indoxyl sulfate induces oxidative stress in endothelial cells. *J. Thromb. Haemost.* **2007**, *5*, 1302–1308.
78. Watanabe, I.; Tatebe, J.; Namba, S.; Koizumi, M.; Yamazaki, J.; Morita, T. Activation of aryl hydrocarbon receptor mediates indoxyl sulfate-induced monocyte chemoattractant protein-1 expression in human umbilical vein endothelial cells. *Circ. J.* **2013**, *77*, 224–230.
79. Muteliefu, G.; Enomoto, A.; Niwa, T. Indoxyl sulfate promotes proliferation of human aortic smooth muscle cells by inducing oxidative stress. *J. Ren. Nutr.* **2009**, *19*, 29–32.
80. Shimizu, H.; Yisireyili, M.; Nishijima, F.; Niwa, T. Stat3 contributes to indoxyl sulfate-induced inflammatory and fibrotic gene expression and cellular senescence. *Am. J. Nephrol.* **2012**, *36*, 184–189.
81. Muteliefu, G.; Shimizu, H.; Enomoto, A.; Nishijima, F.; Takahashi, M.; Niwa, T. Indoxyl sulfate promotes vascular smooth muscle cell senescence with upregulation of p53, p21, and prelamin A through oxidative stress. *Am. J. Physiol. Cell. Physiol.* **2012**, *303*, C126–C134.
82. Shimizu, H.; Hirose, Y.; Nishijima, F.; Tsubakihara, Y.; Miyazaki, H. ROS and PDGF-beta [corrected] receptors are critically involved in indoxyl sulfate actions that promote vascular smooth muscle cell proliferation and migration. *Am. J. Physiol. Cell. Physiol.* **2009**, *297*, C389–C396.

83. Shimizu, H.; Hirose, Y.; Goto, S.; Nishijima, F.; Zrelli, H.; Zghonda, N.; Niwa, T.; Miyazaki, H. Indoxyl sulfate enhances angiotensin II signaling through upregulation of epidermal growth factor receptor expression in vascular smooth muscle cells. *Life Sci.* **2012**, *91*, 172–177.
84. Yamamoto, H.; Tsuruoka, S.; Ioka, T.; Ando, H.; Ito, C.; Akimoto, T.; Fujimura, A.; Asano, Y.; Kusano, E. Indoxyl sulfate stimulates proliferation of rat vascular smooth muscle cells. *Kidney Int.* **2006**, *69*, 1780–1785.
85. Meng, D.; Lv, D.D.; Zhuang, X.; Sun, H.; Fan, L.; Shi, X.L.; Fang, J. Benzo [a]pyrene induces expression of matrix metalloproteinases and cell migration and invasion of vascular smooth muscle cells. *Toxicol. Lett.* **2009**, *184*, 44–49.
86. Kerzee, J.K.; Ramos, K.S. Activation of c-Ha-ras by benzo(a)pyrene in vascular smooth muscle cells involves redox stress and aryl hydrocarbon receptor. *Mol. Pharmacol.* **2000**, *58*, 152–158.
87. Knaapen, A.M.; Curfs, D.M.; Pachen, D.M.; Gottschalk, R.W.; de Winther, M.P.; Daemen, M.J.; Van Schooten, F.J. The environmental carcinogen benzo [a]pyrene induces expression of monocyte-chemoattractant protein-1 in vascular tissue: a possible role in atherogenesis. *Mutat. Res.* **2007**, *621*, 31–41.
88. Vogel, C.F.; Nishimura, N.; Sciallo, E.; Wong, P.; Li, W.; Matsumura, F. Modulation of the chemokines KC and MCP-1 by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in mice. *Arch. Biochem. Biophys.* **2007**, *461*, 169–175.
89. Vorderstrasse, B.A.; Kerkvliet, N.I. 2,3,7,8-Tetrachlorodibenzo-p-dioxin affects the number and function of murine splenic dendritic cells and their expression of accessory molecules. *Toxicol. Appl. Pharmacol.* **2001**, *171*, 117–125.
90. Kimura, A.; Naka, T.; Nohara, K.; Fujii-Kuriyama, Y.; Kishimoto, T. Aryl hydrocarbon receptor regulates Stat1 activation and participates in the development of Th17 cells. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 9721–9726.
91. Choi, W.; Eum, S.Y.; Lee, Y.W.; Hennig, B.; Robertson, L.W.; Toborek, M. PCB 104-induced proinflammatory reactions in human vascular endothelial cells: Relationship to cancer metastasis and atherogenesis. *Toxicol. Sci.* **2003**, *75*, 47–56.
92. Han, S.G.; Eum, S.Y.; Toborek, M.; Smart, E.; Hennig, B. Polychlorinated biphenyl-induced VCAM-1 expression is attenuated in aortic endothelial cells isolated from caveolin-1 deficient mice. *Toxicol. Appl. Pharmacol.* **2010**, *246*, 74–82.
93. Majkova, Z.; Smart, E.; Toborek, M.; Hennig, B. Up-regulation of endothelial monocyte chemoattractant protein-1 by coplanar PCB77 is caveolin-1-dependent. *Toxicol. Appl. Pharmacol.* **2009**, *237*, 1–7.
94. Choi, Y.J.; Arzuaga, X.; Kluemper, C.T.; Caraballo, A.; Toborek, M.; Hennig, B. Quercetin blocks caveolae-dependent pro-inflammatory responses induced by co-planar PCBs. *Environ. Int.* **2010**, *36*, 931–934.
95. Oesterling, E.; Toborek, M.; Hennig, B. Benzo [a]pyrene induces intercellular adhesion molecule-1 through a caveolae and aryl hydrocarbon receptor mediated pathway. *Toxicol. Appl. Pharmacol.* **2008**, *232*, 309–316.
96. Mezrich, J.D.; Fechner, J.H.; Zhang, X.; Johnson, B.P.; Burlingham, W.J.; Bradfield, C.A. An interaction between kynurenine and the aryl hydrocarbon receptor can generate regulatory T cells. *J. Immunol.* **2010**, *185*, 3190–3198.

97. Lekawanvijit, S.; Adrahtas, A.; Kelly, D.J.; Kompa, A.R.; Wang, B.H.; Krum, H. Does indoxyl sulfate, a uraemic toxin, have direct effects on cardiac fibroblasts and myocytes? *Eur. Heart J.* **2010**, *31*, 1771–1779.
98. Ito, S.; Higuchi, Y.; Yagi, Y.; Nishijima, F.; Yamato, H.; Ishii, H.; Osaka, M.; Yoshida, M. Reduction of indoxyl sulfate by AST-120 attenuates monocyte inflammation related to chronic kidney disease. *J. Leukoc Biol.* **2013**, *93*, 837–845.
99. Ito, S.; Osaka, M.; Higuchi, Y.; Nishijima, F.; Ishii, H.; Yoshida, M. Indoxyl sulfate induces leukocyte-endothelial interactions through up-regulation of E-selectin. *J. Biol. Chem.* **2010**, *285*, 38869–38875.
100. Lee, C.T.; Lee, Y.T.; Ng, H.Y.; Chiou, T.T.; Cheng, C.I.; Kuo, C.C.; Wu, C.H.; Chi, P.J.; Lee, W.C. Lack of modulatory effect of simvastatin on indoxyl sulfate-induced activation of cultured endothelial cells. *Life Sci.* **2012**, *90*, 47–53.
101. Tumor, Z.; Shimizu, H.; Enomoto, A.; Miyazaki, H.; Niwa, T. Indoxyl sulfate upregulates expression of ICAM-1 and MCP-1 by oxidative stress-induced NF-kappaB activation. *Am. J. Nephrol.* **2010**, *31*, 435–441.
102. Masai, N.; Tatebe, J.; Yoshino, G.; Morita, T. Indoxyl sulfate stimulates monocyte chemoattractant protein-1 expression in human umbilical vein endothelial cells by inducing oxidative stress through activation of the NADPH oxidase-nuclear factor-kappaB pathway. *Circ. J.* **2010**, *74*, 2216–2224.
103. Dou, L.; Sallee, M. Aix Marseille Université, France, Unpublished work, 2014.
104. Lin, C.J.; Pan, C.F.; Liu, H.L.; Chuang, C.K.; Jayakumar, T.; Wang, T.J.; Chen, H.H.; Wu, C.J. The role of protein-bound uremic toxins on peripheral artery disease and vascular access failure in patients on hemodialysis. *Atherosclerosis* **2012**, *225*, 173–179.
105. Chitalia, V.C.; Shivanna, S.; Martorell, J.; Balcells, M.; Bosch, I.; Kolandaivelu, K.; Edelman, E.R. Uremic serum and solutes increase post-vascular interventional thrombotic risk through altered stability of smooth muscle cell tissue factor. *Circulation* **2013**, *127*, 365–376.
106. Ohtake, F.; Baba, A.; Takada, I.; Okada, M.; Iwasaki, K.; Miki, H.; Takahashi, S.; Kouzmenko, A.; Nohara, K.; Chiba, T.; *et al.* Dioxin receptor is a ligand-dependent E3 ubiquitin ligase. *Nature* **2007**, *446*, 562–566.
107. Kopec, A.K.; Burgoon, L.D.; Ibrahim-Aibo, D.; Burg, A.R.; Lee, A.W.; Tashiro, C.; Potter, D.; Sharratt, B.; Harkema, J.R.; Rowlands, J.C.; *et al.* Automated dose-response analysis and comparative toxicogenomic evaluation of the hepatic effects elicited by TCDD, TCDF, and PCB126 in C57BL/6 mice. *Toxicol. Sci.* **2010**, *118*, 286–297.
108. Li, W.; Matsumura, F. Significance of the nongenomic, inflammatory pathway in mediating the toxic action of TCDD to induce rapid and long-term cellular responses in 3T3-L1 adipocytes. *Biochemistry* **2008**, *47*, 13997–14008.