



HAL
open science

Diffusing-Wave Spectroscopy of Cold Atoms in Ballistic Motion

Aurélien Eloy, Zhibin Yao, Romain Bachelard, William Guerin, Mathilde Fouché, Robin Kaiser

► **To cite this version:**

Aurélien Eloy, Zhibin Yao, Romain Bachelard, William Guerin, Mathilde Fouché, et al.. Diffusing-Wave Spectroscopy of Cold Atoms in Ballistic Motion. 2017. hal-01611345v1

HAL Id: hal-01611345

<https://hal.science/hal-01611345v1>

Preprint submitted on 5 Oct 2017 (v1), last revised 3 Jan 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Diffusing-Wave Spectroscopy of Cold Atoms in Ballistic Motion

Aurélien Eloy¹, Zhibin Yao¹, Romain Bachelard², William Guerin¹, Mathilde Fouché^{1,*} and Robin Kaiser¹

¹*Université Côte d'Azur, CNRS, INPHYNI, France and*

²*Departamento de Física, Universidade Federal de São Carlos,*

Rod. Washington Luís, km 235 - SP-310, 13565-905 São Carlos, SP, Brazil

(Dated: October 5, 2017)

Diffusing-wave spectroscopy is a powerful technique which consists in measuring the temporal correlation function of the intensity of light multiply scattered by a medium. In this paper, we apply this technique to cold atoms under purely ballistic motion and we investigate the transition between the single and the multiple scattering regime. The intensity correlation function changes from a simple Gaussian function, whose width reveals the sample temperature, to a more complex decay shortened by the frequency redistribution in the thick cloud. These features are quantitatively compared to simulations with a phase-coherent and an incoherent model. Both show a very good agreement with the experiments.

I. INTRODUCTION

The study of fluctuations and correlations often gives access to information not contained in averaged values. Among the many statistical properties of a fluctuating field, the intensity correlation function is largely used in a number of areas, from astronomy [1, 2], to quantum optics [3–6], particle physics [7], and to mesoscopic optics. In the latter, it has been applied to the fluctuations of light scattered by a disordered medium. First used in the single scattering regime with a technique known as dynamic light scattering or quasielastic light scattering [8, 9], it was then extended to strong multiple scattering regime. This powerful technique, called diffusing-wave spectroscopy (DWS) [10–12], allows probing extremely small displacement, much smaller than the optical wavelength. It has found a wide range of applications, going from particle sizing [12–14], Brownian motion [15, 16], hydrodynamic interactions [17, 18], microscopic dynamics [19, 20] and medical applications [21, 22].

DWS consists in measuring the temporal intensity correlation $g^{(2)}(\tau)$ of the light scattered by the medium under investigation. In the case of a stationary process, this quantity is expressed as:

$$g^{(2)}(\tau) = \frac{\langle I(t)I(t+\tau) \rangle}{\langle I(t) \rangle^2}, \quad (1)$$

where $\langle \rangle$ denotes time averaging. If light is scattered by a large number of particles it can be considered as chaotic light for which $g^{(2)}(\tau = 0)$ is equal to 2, representing photon bunching, and it goes to 1 for large τ when events become uncorrelated. The temporal decay of $g^{(2)}(\tau)$ gives access to the mean square displacement of the scatterers. On the other hand, the shape of $g^{(2)}(\tau)$ on different time scales provides information on the scatterers dynamics [15, 20, 23]. The standard theoretical approach to describe DWS is to use the diffusion theory

for light. The first order correlation function $g^{(1)}(\tau)$ can be derived [12], and the second order is obtained using the Siegert relation [5, 12]:

$$g^{(2)}(\tau) = 1 + \beta |g^{(1)}(\tau)|^2. \quad (2)$$

The factor β is related to the number of detected spatial and polarization modes. For a detector whose radius is much smaller than the spatial coherence length of the scattered light and for polarized light β is equal to unity [5].

Here, we apply the DWS technique on a cold atomic cloud. The light beam is set close to resonance and its frequency is changed to probe the transition from the single to the multiple scattering regime. This is a first difference from other DWS measurements where light is always considered to be far from any resonance. Cold atoms also allow for a pure ballistic regime, which is usually hardly reached in standard DWS measurements due to the surrounding fluid in which scatterers are immersed [10, 11, 15, 16]. The intensity correlation function of light scattered by cold atoms has already been measured in several experiments and signatures of the Doppler broadening, Lamb-Dicke narrowing, radiation trapping and inelastic scattering (Mollow triplet) could be obtained [24–29]. However, these measurements were done with fluorescence from atoms trapped in a magneto-optical trap (MOT) or during a molasse phase, with thus large beams (probing different parts of the cloud and thus different optical thicknesses) and quite strong intensities, leading to a complicated interplay between all these effects.

In this paper, we perform intensity correlation measurements while the atoms are ballistically expanding with a small and weak probe beam, which we can tune independently from the laser-cooling beams. At the cost of a long integration time, this allows us to avoid inelastic effects and precisely investigate the transition from the single-scattering to the multiple scattering regime with a well-defined optical thickness. The behavior of the $g^{(2)}$ function is distinctively different from that obtained in other media (scatterers in a fluid) and it is only related to the velocity distribution of the atoms. We observe in

* mathilde.fouche@inphyni.cnrs.fr

particular a change of shape of the $g^{(2)}$ function, from a Gaussian in the single-scattering regime to a more complex shape in the multiple scattering regime, which in good agreement with simulations.

II. EXPERIMENTAL SETUP

In our experiment, the scattering sample is a cold atomic vapor produced by a MOT that can contain up to a few 10^9 atoms of ^{85}Rb with a Gaussian density distribution of rms size $R \simeq 1$ mm giving a typical peak-density $\rho_0 \simeq 10^{11} \text{ cm}^{-3}$. The temperature is of the order of $200 \mu\text{K}$, measured by a standard time-of-flight (TOF) technique. Our sample is also characterized by its on-resonance optical thickness b_0 . This quantity is inferred from the measurement of the transmission of a small probe beam going through the center of the atomic cloud as a function of the frequency detuning on the $|3\rangle \rightarrow |4'\rangle$ D_2 hyperfine transition as shown on Fig. 1b (frequency ω_0 , wavelength $\lambda = 780.24$ nm, linewidth $\Gamma/2\pi = 6.07$ MHz) [30].

To measure the intensity correlations of the scattered light, we use the following time sequence. First, the MOT is loaded from a background vapor for 30 ms, followed by a compression stage of 35 ms. Then, the MOT trapping beams and the magnetic field gradient are switched off allowing for a 2-ms free expansion. Next, we apply two pulses of a weak probe beam with a waist of $250 \mu\text{m}$, linearly polarized and detuned by $\delta = \omega_L - \omega_0$ with $\delta > 0$, ω_L being the laser frequency. This probe beam is delivered by a distributed-feedback (DFB) laser, amplified by a tapered amplifier. Although it is known that DFB lasers have strong frequency noise [30, 31], it does not affect our measurements of intensity correlations [32]. The duration of each probe pulse is $100 \mu\text{s}$, corresponding to a time-window where the intensity correlations are recorded, separated by a $50\text{-}\mu\text{s}$ pulse of repumper and $50 \mu\text{s}$ of free expansion. From one cycle to the other, atoms are recycled and we reach a steady state with $N \simeq 3 \times 10^9$ atoms and $b_0 = 105 \pm 4$, measured between each acquisition of the $g^{(2)}$ function.

The setup for measuring $g^{(2)}(\tau)$ is represented in Fig. 1. The scattered light is collected by a single mode fiber (SMF) at a distance $L = 40$ cm with an angle of $\theta = 41^\circ$ from propagation direction of the probe beam. Only one polarization is coupled into the SMF thanks to a polarizing beam splitter (PBS). The combination of the SMF and the PBS ensures the condition of maximum spatial coherence, $\beta = 1$. The coupled light is then split using a 50:50 fiber beam splitter and finally detected on two single-photon avalanche photodiodes (APDs, SPCM-AQRH from Excelitas Technologies), gated on the probe pulses. Time tags, with a resolution of 162 ps, are finally obtained from a multichannel time-to-digital converter (TDC, ID800 from IDQuantique) and sent to a computer to record the histogram of the coincidences. Number of counts during the $200 \mu\text{s}$ total probe pulse range from 7




FIG. 1. Experimental set-up. Light scattered in the MOT is collected using a single-mode fiber (SMF), after passing through a polarizing beam splitter (PBS) to select only one polarization. The light is then split in a 50:50 fiber beam splitter (FBS). The outputs of the FBS are connected to two APDs. The single counts in each APDs are time-tagged by a time-to-digital converter (TDC) and analyzed with a computer. Left bottom: Relevant atomic levels of the D_2 hyperfine transition of ^{85}Rb .

to 40, well above the dark count rate (< 0.02 during the total probe pulse). Typical integration times range from a few hours to a few days.

To reconstruct the $g^{(2)}$ function, we need to normalize the recorded coincidences. For delays much larger than the coherence time, $g^{(2)}(\tau)$ is expected to be flat and equal to 1. However, in this range, because of the finite time-window of the measurement, the number of recorded coincidences decreases linearly with the delay and reaches 0 at the pulse duration. The $g^{(2)}$ function is thus obtained by dividing the data by the fitted slope at large delays. The detection scheme and the normalisation were tested by measuring the light scattered by a piece of paper at rest illuminated by the probe laser, giving $g^{(2)}(\tau) = 1$ within the statistical uncertainty.

Exploiting the narrow atomic resonance, we can easily tune the optical thickness of the sample

$$b(\delta) = \frac{b_0}{1 + 4\delta^2/\Gamma^2}, \quad (3)$$

by varying the detuning δ of the probe. We can thus explore the transition from the single scattering regime, where $b(\delta) \ll 1$, to the multiple scattering regime, $b(\delta) \gg 1$, in principle up to $b \sim 100$. However, one limitation is the amount of light scattered in the detection direction, which decreases with b for $b > 1$ [33]. To keep a reasonable integration time, we have thus limited our investigation to $b(\delta) \leq 2$. Note that the intensity of the probe is adapted to each detuning in order to keep the saturation parameter constant at $s \simeq 5 \times 10^{-2}$, with $s = (I/I_{\text{sat}})/(1 + 4\delta^2/\Gamma^2)$ and $I_{\text{sat}} \simeq 3.7 \text{ mW/cm}^2$. This low value of s ensures that inelastic scattering is negligible.




FIG. 2. Intensity correlation function $g^{(2)}(\tau)$ with $b_0 \simeq 105$ and different detunings $\delta/\Gamma = \{20, 5, 3\}$ from left to right, corresponding to different optical thicknesses, (a) $b(\delta) = 0.07$, (b) $b(\delta) = 1$ and (c) $b(\delta) = 2$. Grey continuous curve: experimental data; red dotted dashed line: random walk simulations; black dashed line: coupled-dipole simulations. The curves obtained by the simulations have been renormalized to take into account the decrease of the contrast (measured on the experimental data on the first 30 ns) and due to spontaneous Raman-scattering. Insets : $g^{(2)}(\tau) - 1$ obtained experimentally in semi-log scale. A quadratic behavior is added (dashed line) to guide the eyes.

III. RESULTS AND DISCUSSION

We present in Fig. 2 several measurements of $g^{(2)}(\tau)$ for different $b(\delta)$, from the single scattering to the multiple scattering regime. From these experimental curves, three observations can be made, as detailed in the following.

First, a systematic analysis of the contrast, defined as $\mathcal{C} = g^{(2)}(0) - 1$, with respect to the detuning shows that it is almost one near resonance, but slightly decreases at large detuning. We have checked that this loss of contrast could not be explained by any correlation artefacts (due to the light scattered by the vacuum cell or by the hot vapor for example) or by the dark count of the detectors. We attribute this decrease to spontaneous Raman scattering between the two hyperfine ground states, $|3\rangle \rightarrow |2\rangle$, via off-resonant excitation of the $|3'\rangle$ and $|2'\rangle$ levels (see Fig. 1b). As the detuning from the $|4'\rangle$ level increases, the Rayleigh scattering decreases, whereas the Raman scattering is almost constant. Thus, the relative weight of Raman scattering increases. Since the two ground states are separated by 3 GHz, the interference between Rayleigh- and Raman-scattered light induces a beating that is too fast to be resolved with the ~ 350 ps time resolution of our APDs. This leads to a time averaging and thus a reduction of the contrast, as already seen in [34] with hot vapors.

Second, increasing the optical thickness leads to a narrowing of the intensity correlation function, i.e., to a decrease of the temporal coherence. This is a direct signature of the frequency redistribution induced by multiple scattering. Each scattering event is indeed associated to a frequency shift $\delta\omega = \vec{\Delta k} \cdot \vec{v}$ due to Doppler effect, where $\vec{\Delta k}$ is the difference between the wavevectors of the scattered photon and of the incident one, and \vec{v} is the velocity of the scattering atom. Since all velocity directions are

equiprobable, $\delta\omega$ is a centered random variable [35]. As a consequence this frequency redistribution induces a diffusion in frequency space, leading in average to a broadening of the spectrum, and correspondingly a decrease of the temporal correlation with the number of scattering events, and thus with the optical thickness. Note that previous signatures of this effect have been indirectly observed in radiation-trapping experiments [36–38]. This reduction of the coherence time with multiple scattering is the key idea of DWS, since it corresponds to an increased sensitivity to the motion of scatterers. It should also be noted that contrary to previous reports [24, 26], multiple scattering does not induce here any reduction of the contrast of $g^{(2)}(\tau)$, because we only have here coherent (Rayleigh) scattering.

A more detailed study of the evolution of the coherence time τ_{coh} , defined here as the half width at half maximum (HWHM) of the $g^{(2)}$ function, as a function of the optical thickness $b(\delta)$ is reported in Fig. 3. The HWHM extracted from the experimental $g^{(2)}$ function correspond to the black triangles. It shows a plateau at very small b and a decrease as soon as b is not much smaller than unity. The plateau is due to Doppler broadening in the single scattering regime. It is directly related to the temperature of the atomic sample but also strongly depends on the direction of observation θ from the incident wave vector. The Doppler shift for each scattering event is indeed given by $\delta\omega(\theta) = \vec{\Delta k} \cdot \vec{v} = k[v_{\parallel}(\cos\theta - 1) + v_{\perp} \sin\theta]$, where v_{\parallel}, v_{\perp} are the longitudinal and transverse components of the atomic velocity along the direction defined by the incident wave vector, and $k = 2\pi/\lambda$. For cold atoms, $\langle v_{\parallel}^2 \rangle = \langle v_{\perp}^2 \rangle = k_{\text{B}}T/M$, M being the atomic mass of ^{85}Rb and k_{B} the Boltzmann constant, which gives a Doppler width $\Delta\omega(\theta) = k\sqrt{2(1 - \cos\theta)}k_{\text{B}}T/M$ [35]. Since the atomic velocity is given by the Maxwell-Boltzmann dis-




FIG. 3. Coherence time, defined as the HWHM of $g^{(2)}(\tau)$, as a function of the detuning-dependent optical thickness. The black triangles with error bars are extracted from experimental data, the red diamonds from the random walk simulations and the orange circles from the coupled-dipole simulations.

tribution, the Doppler-broadened spectrum is Gaussian:

$$S(\omega_L, \theta) \propto e^{-(\omega_L - \omega_0)^2 / 2\Delta\omega^2(\theta)}. \quad (4)$$

Therefore, in the single scattering limit, the first order coherence function $g^{(1)}(\tau)$, which is the Fourier transform of the optical spectrum, is also Gaussian:

$$g^{(1)}(\tau, \theta) = e^{-\tau^2(1 - \cos\theta)/4\tau_c^2}, \quad (5)$$

with $\tau_c^{-1} = 2k\sqrt{k_B T/M}$ and, according to the Siegert relation (2), so is $g^{(2)}(\tau)$:

$$g^{(2)}(\tau) = 1 + e^{-\tau^2(1 - \cos\theta)/2\tau_c^2}. \quad (6)$$

Taking the data at the lowest optical thickness $b = 0.02$, we extract a HWHM $\tau_{\text{coh}} = (961 \pm 30)$ ns, corresponding to a temperature of (245 ± 15) μ K.

Finally, the third observation featured on Fig. 2 is the progressive change of shape of the $g^{(2)}$ function. At low optical thicknesses, $g^{(2)}$ is well fitted by a Gaussian. As expected and explained before, this is obtained due to the Maxwell-Boltzmann distribution of the atomic velocity but also because we can assume that the atoms undergo purely ballistic motions during the whole scattering process. It is important to note that, for standard DWS performed on suspended particles in a liquid, this regime is hardly reached. The interactions with the fluid induce in particular a long-time diffusive motion for the particles and thus an exponential decay for $g^{(2)}$ [10, 11, 15].

At short time scale, $g^{(2)}$ remains Gaussian with a width that decreases as b is increased, as shown on the insets of Fig. 2. On the other hand, in the multiple scattering regime and for larger time-scale, the shape clearly deviates from a Gaussian, with a decrease that gets closer to linear in semi-log scale instead of quadratic, and becomes more complex to infer analytically. In order to understand this change, we have performed numerical simulations to probe the role of multiple scattering, based on the two following models.




FIG. 4. Verification of the Siegert relation with a cloud of $N = 5000$ atoms, $b_0 = 2.8$, $b(\delta) = 2$ and $\theta = 41^\circ$. Black dashed line: $g^{(2)}(\tau)$; red dashed line: $1 + |g^{(1)}(\tau)|^2$. The Siegert relation is well satisfied.

The first model, known as the coupled-dipole model (CD) [39–42], describes the sample as an ensemble of identical atoms driven by a laser field. Atoms are modeled as point-like dipoles interacting via the light in the scalar approximation. Their position changes in time following the velocity distribution corresponding to the temperature, and they are spatially distributed with a Gaussian shape that mimics the geometry of the cloud. In the recent years, this model has been widely used in the context of single-photon superradiance and subradiance [43–47]. It is valid at low intensity, when the saturation parameter $s \ll 1$, but it includes coherent and cooperative effects. Since this coherent model allows computing the time-dependent radiated field of each dipole, it provides access to both the $g^{(1)}(\tau)$ and $g^{(2)}(\tau)$, so it can be used to check the validity of the Siegert relation, as demonstrated in Fig. 4 for $b = 2$. The use of the CD model is however limited to a few thousands atoms and thus to a maximum resonant optical thickness of $b_0 \sim 10$ for dilute samples. Still, by adapting the detuning, the same $b(\delta)$ as in the experiment can be simulated. Simulations of the CD model reveal that the intensity correlation function indeed depends only on $b(\delta)$.

The second model is a Monte-Carlo simulation modeling an incoherent random walk (RW) of photons inside the atomic cloud. The step of a photon sent into the medium is randomly sampled taking into account a step-length distribution due to the Gaussian density profile of the cloud and the dependent-detuning isotropic scattering cross-section. Doppler effect is included in the model by adding a frequency shift after each scattering event, randomly sampled from the velocity distribution. It is then possible to select all photons exiting at a particular angle and to get the output spectrum of the scattered light $S(\omega_L, \theta)$, which corresponds to the Fourier transform of the $g^{(1)}$ function. We can finally access the $g^{(2)}$ function by assuming the Siegert relation (2), validated by the CD model.

The RW simulations are useful to understand the shape of the intensity correlation function. It gives access

to the distribution $P(n)$ of photons undergoing n scattering events before escaping, to the corresponding optical spectrum $S_n(\omega_L, \theta)$ and its Fourier transform $g_n^{(1)}(\tau, \theta)$. While an analytical expression of $P(n)$ is hardly accessible, we have checked that, for small b ($b \lesssim 2$), $g_n^{(1)}(\tau, \theta)$ can be obtained analytically. For one scattering event, one gets the spectrum and the first order correlation function given by Eq. (4) and Eq. (5). For two scattering events, let us first consider the case of a first photon scattered in the θ_1 direction. Since the measurement is done in the θ direction, the second photon must be scattered at the angle $\theta - \theta_1$ compared to the first one. The spectrum is then given by the convolution of the two individual Doppler spectra $S(\omega_L, \theta_1) * S(\omega_L, \theta - \theta_1)$. Taking the Fourier transform, the corresponding $g^{(1)}$ function is $g_1^{(1)}(\tau, \theta_1) \times g_1^{(1)}(\tau, \theta - \theta_1)$. Finally, the total $g_2^{(1)}$ is the sum over all the possible angles:

$$g_2^{(1)}(\tau, \theta) = \int_0^{2\pi} g_1^{(1)}(\tau, \theta_1) g_1^{(1)}(\tau, \theta - \theta_1) d\theta_1 \quad (7)$$

$$= e^{-\frac{2\tau^2}{4\tau_c^2}} I_0 \left(\frac{\tau^2}{4\tau_c^2} \sqrt{2(1 + \cos \theta)} \right), \quad (8)$$

with I_0 the modified Bessel function of the first kind. While in the single scattering regime the $g_2^{(1)}$ is a Gaussian function, its shape is more complex in the double scattering regime. The principle of the calculation is the same when the number of scattering events is increased and one gets:

$$g_n^{(1)}(\tau, \theta) = \int_0^{2\pi} \left[\prod_{i=1}^{n-1} g_1^{(1)}(\tau, \theta_i) \right] g_1^{(1)} \left(\tau, \theta - \sum_{j=1}^{n-1} \theta_j \right) d\theta_1 \dots d\theta_{n-1}, \quad (9)$$

When n gets larger, the contribution of θ is blurred. Moreover the Gaussian part decreases more rapidly than the Bessel function and one recovers the Gaussian shape:

$$g_n^{(1)}(\tau, \theta) \xrightarrow{n \rightarrow \infty} e^{-\frac{n\tau^2}{4\tau_c^2}}, \quad (10)$$

with a HWHM that depends on the number of scattering events as $1/\sqrt{n}$. As said before, the $g_n^{(2)}$ functions are calculated using the Siegert relation. The analytical expressions are compared to the ones obtained by the RW simulations in Fig. 5, showing a good agreement between them. The total $g^{(2)}$ function is finally given by:

$$g^{(2)}(\tau) = 1 + \left| \sum_{n>0} P(n) g_n^{(1)}(\tau, \theta) \right|^2. \quad (11)$$

Its shape depends on the individual shape of $g_n^{(1)}$ as well as $P(n)$ which is only only accessible exploiting the RW simulations.

To compare the experimental results and the simulations, we first extract the coherence time. The results




FIG. 5. Intensity correlation function $g_n^{(2)}(\tau, \theta = 41^\circ)$ for photons that have been scattered n times before escaping. Solid line: obtained using the analytical expressions from (5) to (10). Dashed line: extracted from the RW simulations with $b(\delta) = 2$. In diminishing order of width: $n = 1$ (black), $n = 2$ (blue) and $n = 7$ (green).

are plotted in Fig. 3 with the two models. The RW simulations are performed with the actual parameters of the experiment and the CD simulations use a lower b_0 and correspondingly lower detunings to match the same $b(\delta)$. The only free parameter is the temperature, which is chosen in the simulations in order to match the one extracted from the τ_{coh} measured with the lowest b . Despite the differences between the two models, there is a remarkable agreement between them and with the experimental data.

Finally, the shape of $g^{(2)}(\tau)$ obtained experimentally is compared with the simulations in Fig. 2. The simulated curves are renormalized to take into account the decrease of the contrast due to spontaneous Raman-scattering and measured on the experimental data on the first 30 ns. We also see a good agreement between all curves, showing that the effect of the multiple scattering is well described by both the CD and the RW models.

IV. CONCLUSION

To summarize, we have precisely measured the temporal intensity correlation function $g^{(2)}(\tau)$ of light scattered by cold atoms in ballistic motion, a regime hardly reached in standard DWS measurements. Different regimes of light transport have been investigated, from the single scattering to the multiple scattering regime. We have observed a decrease of the width of $g^{(2)}(\tau)$ as we enter the multiple scattering regime and we have evidenced the corresponding change of shape in $g^{(2)}(\tau)$. Numerical simulations, based on the coupled-dipole model or on random walk simulations, have been performed to further understand the role of multiple scattering of light in intensity correlations. A remarkable agreement is obtained between the two simulations and with experimental data.

DWS is a powerful technique that has been widely used in soft condensed matter but it can find many applications in other field of research, ranging from atomic physics to astrophysics. In hot atomic vapors, DWS might allow to study time dependance in Levy flights [48, 49] and the connection between Lévy flights and Lévy walks. In cold atomic samples, it might be used to quantify the friction induced by Doppler or sub-Doppler cooling. Anomalous diffusion, as expected close to the decrochage could be investigated with an alternative approach to that used in [50, 51]. DWS might also prove to be a powerful tool to investigate phase transitions, as for instance Anderson localization of light by cold atoms [52, 53], and to study the impact of Dicke type super- and subradiance [45–47] on fluctuations of scattered or emitted light. Finally, intensity correlations are also a well known tool to distinguish classical from quantum optics [3]. This will allow us to probe the onset

of gain and random lasing in our cloud of cold atoms [54]. Gain and lasing being also expected in astrophysical systems [55], the recent success of intensity correlations measured on three bright stars [2] opens this quantum eye to lasing in astrophysics.

ACKNOWLEDGEMENTS

We thank Antoine Dussaux for the TDC programs and S. Vartabi Kashanian for the early stage of the DWS experiment. A.E. is supported by a grant from the DGA. R. B. benefited from Grants from São Paulo Research Foundation (FAPESP) (Grant Nos. 2015/50422-4 and 2014/01491-0). R. B., M. F and R. K. received support from project CAPES-COFECUB (Ph879-17/CAPES 88887.130197/2017-01). The Titan X Pascal used for this research was donated by the NVIDIA Corporation.

-
- [1] R. Hanbury Brown and R. Q. Twiss, “A test of a new type of stellar interferometer on Sirius,” *Nature* **178**, 1046–1048 (1956).
- [2] W. Guerin, A. Dussaux, M. Fouché, G. Labeyrie, J.-P. Rivet, D. Vernet, F. Vakili, and R. Kaiser, “Temporal intensity interferometry: photon bunching on three bright stars,” *MNRAS* **472**, 4126 (2017).
- [3] R. J. Glauber, “The quantum theory of optical coherence,” *Phys. Rev. A* **130** (1963).
- [4] F. T. Arecchi, E. Gatti, and A. Sona, “Time distribution of photons from coherent and Gaussian sources,” *Phys. Lett.* **20**, 27–29 (1966).
- [5] R. Loudon, *The Quantum Theory of Light*, 3rd ed. (Oxford University Press, 2000).
- [6] H.-A. Bachor and T. C. Ralph, *A Guide to Experiments in Quantum Optics*, 2nd ed. (Wiley, 2004).
- [7] G. Alexander, “Bose-Einstein and Fermi-Dirac interferometry in particle physics,” *Rep. Prog. Phys.* **66**, 481–522 (2003).
- [8] N. A. Clark, J. H. Lunacek, and G. B. Benedek, “A study of brownian motion using light scattering,” *American Journal of Physics* **38**, 575–585 (1970).
- [9] B. J. Berne and R. Pecora, *Dynamic light scattering* (Wiley, New York, 1976).
- [10] G. Maret and P. E. Wolf, “Multiple light scattering from disordered media. The effect of brownian motion of scatterers,” *Zeitschrift für Physik B Condensed Matter* **65**, 409–413 (1987).
- [11] D. J. Pine, D. A. Weitz, P. M. Chaikin, and E. Herbolzheimer, “Diffusing wave spectroscopy,” *Phys. Rev. Lett.* **60**, 1134–1137 (1988).
- [12] D. J. Pine, D. A. Weitz, J. X. Zhu, and E. Herbolzheimer, “Diffusing-wave spectroscopy: dynamic light scattering in the multiple scattering limit,” *J. Phys. France* **51**, 2101–2127 (1990).
- [13] C. Rega, C. J. Lloyd, D. Attwood, D. Clarke, and P. Gerahty, “Temporal autocorrelation function for a diffusing-wave spectroscopy experiment with a point source and backscattering detection,” *Appl. Opt.* **40**, 4204–4209 (2001).
- [14] F. Scheffold, “Particle sizing with diffusing wave spectroscopy,” *Journal of Dispersion Science and Technology* **23**, 591–599 (2002).
- [15] D. A. Weitz, D. J. Pine, P. N. Pusey, and R. J. A. Tough, “Nondiffusive brownian motion studied by diffusing-wave spectroscopy,” *Phys. Rev. Lett.* **63**, 1747–1750 (1989).
- [16] M. H. Kao, A. G. Yodh, and D. J. Pine, “Observation of brownian motion on the time scale of hydrodynamic interactions,” *Phys. Rev. Lett.* **70**, 242–245 (1993).
- [17] S. Fraden and G. Maret, “Multiple light scattering from concentrated, interacting suspensions,” *Phys. Rev. Lett.* **65**, 512–515 (1990).
- [18] D. A. Weitz, J. X. Zhu, D. J. Durain, Hu Gang, and D. J. Pine, “Diffusing-wave spectroscopy: The technique and some applications,” *Physica Scripta* **T49**, 610 (1993).
- [19] P. Hébraud, F. Lequeux, J. P. Munch, and D. J. Pine, “Yielding and rearrangements in disordered emulsions,” *Phys. Rev. Lett.* **78**, 4657–4660 (1997).
- [20] N. Menon and D. J. Durian, “Diffusing-wave spectroscopy of dynamics in a three-dimensional granular flow,” *Science* **275**, 1920–1922 (1997).
- [21] T. Durduran, G. Yu, M. G. Burnett, J. A. Detre, J. H. Greenberg, J. Wang, C. Zhou, and A. G. Yodh, “Diffuse optical measurement of blood flow, blood oxygenation, and metabolism in a human brain during sensorimotor cortex activation,” *Opt. Lett.* **29**, 1766–1768 (2004).
- [22] M. Ninck, M. Untenberger, and T. Gisler, “Diffusing-wave spectroscopy with dynamic contrast variation: disentangling the effects of blood flow and extravascular tissue shearing on signals from deep tissue,” *Biome. Opt. Express* **1**, 1502 (2001).
- [23] M. J. Stephen, “Temporal fluctuations in wave propagation in random media,” *Phys. Rev. B* **37**, 1–5 (1988).
- [24] S. Bali, D. Hoffmann, J. Simán, and T. Walker, “Measurements of intensity correlations of scattered light from laser-cooled atoms,” *Phys. Rev. A* **53**, 3469–3472 (1996).
- [25] C. Jurczak, B. Desruelle, K. Sengstock, J.-Y. Courtois, C. I. Westbrook, and A. Aspect, “Atomic transport in

- an optical lattice: An investigation through polarization-selective intensity correlations,” *Phys. Rev. Lett.* **77**, 1727–1730 (1996).
- [26] R. Stites, M. Beeler, L. Feeney, S. Kim, and S. Bali, “Sensitive measurement of radiation trapping in cold-atom clouds by intensity correlation detection,” *Opt. Lett.* **29**, 2713–5 (2004).
- [27] K. Nakayama, Y. Yoshikawa, H. Matsumoto, Y. Torii, and T. Kuga, “Precise intensity correlation measurement for atomic resonance fluorescence from optical molasses,” *Opt. Express* **18**, 6604 (2010).
- [28] J. A. Grover, P. Solano, L. A. Orozco, and S. L. Rolston, “Photon-correlation measurements of atomic-cloud temperature using an optical nanofiber,” *Phys. Rev. A* **92**, 013850 (2015).
- [29] K. Muhammed Shafi, D. Pandey, B. Suryabrahmam, B.S. Girish, and H. Ramachandran, “Time-delayed intensity-interferometry of the emission from ultracold atoms in a steady-state magneto-optical trap,” *J. Phys. B: At. Mol. Opt. Phys.* **49**, 025301 (2016).
- [30] S. Vartabi Kashanian, A. Eloy, W. Guerin, M. Lintz, M. Fouché, and R. Kaiser, “Noise spectroscopy with large clouds of cold atoms,” *Phys. Rev. A* **94**, 043622 (2016).
- [31] S. Kraft, A. Deninger, C. Trüch, J. Fortágh, F. Lison, and C. Zimmermann, “Rubidium spectroscopy at 778–780 nm with a distributed feedback laser diode,” *Laser Phys. Lett.* **2**, 71–76 (2005).
- [32] The laser frequency noise does not affect the $g^{(2)}$ function as soon as the typical path length of the photon inside the atomic sample is much lower than the laser coherence length.
- [33] G. Labeyrie, D. Delande, C. A. Müller, C. Miniatura, and R. Kaiser, “Multiple scattering of light in a resonant medium,” *Opt. Commun.* **243**, 157–164 (2004).
- [34] A. Dussaux, T. Passerat de Silans, W. Guerin, O. Alibert, S. Tanzilli, F. Vakili, and R. Kaiser, “Temporal intensity correlation of light scattered by a hot atomic vapor,” *Phys. Rev. A* **93**, 043826 (2016).
- [35] Note that this is only a good approximation for cold atoms. For hot vapors, see, e.g., ref. [56].
- [36] G. Labeyrie, E. Vaujour, C. A. Müller, D. Delande, C. Miniatura, D. Wilkowski, and R. Kaiser, “Slow diffusion of light in a cold atomic cloud,” *Phys. Rev. Lett.* **91**, 223904 (2003).
- [37] G. Labeyrie, R. Kaiser, and D. Delande, “Radiation trapping in a cold atomic gas,” *Appl. Phys. B* **81**, 1001–1008 (2005).
- [38] R. Pierrat, B. Grémaud, and D. Delande, “Enhancement of radiation trapping for quasis resonant scatterers at low temperature,” *Phys. Rev. A* **80**, 013831 (2009).
- [39] A. A. Svidzinsky, J.-T. Chang, and M. O. Scully, “Cooperative spontaneous emission of N atoms: Many-body eigenstates, the effect of virtual Lamb shift processes, and analogy with radiation of N classical oscillators,” *Phys. Rev. A* **81**, 053821 (2010).
- [40] Ph. W. Courteille, S. Bux, E. Lucioni, K. Lauber, T. Bienaimé, R. Kaiser, and N. Piovella, “Modification of radiation pressure due to cooperative scattering of light,” *Eur. Phys. J. D.* **58**, 69–73 (2010).
- [41] T. Bienaimé, M. Petruzzo, D. Bigerni, N. Piovella, and R. Kaiser, “Atom and photon measurement in cooperative scattering by cold atoms,” *J. Mod. Opt.* **58**, 1942–1950 (2011).
- [42] T. Bienaimé, R. Bachelard, P. W. Courteille, N. Piovella, and R. Kaiser, “Cooperativity in light scattering by cold atoms,” *Fortschr. Phys.* **61**, 377–392 (2013).
- [43] M. O. Scully, E. S. Fry, C. H. Raymond Ooi, and K. Wódkiewicz, “Directed spontaneous emission from an extended ensemble of N atoms: Timing is everything,” *Phys. Rev. Lett.* **96**, 010501 (2006).
- [44] M. O. Scully and A. A. Svidzinsky, “The super of superradiance,” *Science* **325**, 1510–1511 (2009).
- [45] W. Guerin, M. O. Araújo, and R. Kaiser, “Subradiance in a Large Cloud of Cold Atoms,” *Phys. Rev. Lett.* **116**, 083601 (2016).
- [46] M. O. Araújo, I. Krešić, R. Kaiser, and W. Guerin, “Superradiance in a Large and Dilute Cloud of Cold Atoms in the Linear-Optics Regime,” *Phys. Rev. Lett.* **117**, 073002 (2016).
- [47] S. J. Roof, K. J. Kemp, M. D. Havey, and I. M. Sokolov, “Observation of Single-Photon Superradiance and the Cooperative Lamb Shift in an Extended Sample of Cold Atoms,” *Phys. Rev. Lett.* **117**, 073003 (2016).
- [48] N. Mercadier, W. Guerin, M. Chevrollier, and R. Kaiser, “Lévy flights of photons in hot atomic vapours,” *Nat. Phys.* **5**, 602–605 (2009).
- [49] Q. Baudouin, R. Pierrat, A. Eloy, E. J. Nunes-Pereira, P.-A. Cuniasse, N. Mercadier, and R. Kaiser, “Signatures of Lévy flights with annealed disorder,” *Phys. Rev. E* **90**, 052114 (2014).
- [50] J. Jersblad, H. Ellmann, K. Støchkel, A. Kastberg, L. Sanchez-Palencia, and R. Kaiser, “Non-Gaussian velocity distributions in optical lattices,” *Phys. Rev. A* **69**, 013410 (2004).
- [51] P. Douglas, S. Bergamini, and F. Renzoni, “Tunable Tsallis distributions in dissipative optical lattices,” *Phys. Rev. Lett.* **96**, 110601 (2006).
- [52] S. E. Skipetrov and I. M. Sokolov, “Magnetic-Field-Driven Localization of Light in a Cold-Atom Gas,” *Phys. Rev. Lett.* **114**, 053902 (2015).
- [53] G. L. Celardo, M. Angeli, and R. Kaiser, “Localization of light in subradiant Dicke states: a mobility edge in the imaginary axis,” arXiv:1702.04506 (2017).
- [54] Q. Baudouin, N. Mercadier, V. Guarrera, W. Guerin, and R. Kaiser, “A cold-atom random laser,” *Nat. Phys.* **9**, 357–360 (2013).
- [55] V. S. Letokhov and S. Johansson, *Astrophysical lasers*, edited by Oxford University Press (2009).
- [56] J. C. de A. Carvalho, M. Oriá, M. Chevrollier, H. L. D. de Souza Cavalcante, and T. Passerat de Silans, “Redistribution of light frequency by multiple scattering in a resonant atomic vapor,” *Phys. Rev. A* **91**, 053846 (2015).