

Risk assessment using suprema data Christophette Blanchet-Scalliet, Diana Dorobantu, Laura Gay, Véronique Maume-Deschamps, Pierre Ribereau

▶ To cite this version:

Christophette Blanchet-Scalliet, Diana Dorobantu, Laura Gay, Véronique Maume-Deschamps, Pierre Ribereau. Risk assessment using suprema data. 2017. hal-01615196v1

HAL Id: hal-01615196 https://hal.science/hal-01615196v1

Preprint submitted on 12 Oct 2017 (v1), last revised 15 Oct 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Risk assessment using suprema data^{*}

Christophette Blanchet-ScallietDiana DorobantuLaura GayVéronique Maume-DeschampsPierre Ribereau \parallel

Abstract

The dynamic of temperatures can be modelled by a mean-reverting process such as an Ornstein-Uhlenbeck one. In this study, we estimate the parameters of this process thanks to daily observed suprema of temperatures, which are the only data gathered by some weather stations. The expression of the cumulative distribution function of the process supremum is obtained. The parameters are estimated by a least square method quantiles based on this function. Theoretical results, including mixing property and consistency of model parameter estimation, are provided. The parameters estimation will allow us to estimate risk measures, such as the probability of heat wave. Numerical illustrations are given on simulated data and real ones.

Keywords. Ornstein-Uhlenbeck process, supremum law, parameters estimation, heat wave risk assessment.

1 Introduction

Forecasting and assessing the risk of heat waves is a crucial public policy stake. It requires measure tools in order to evaluate the probability of heat waves and their severity. The available information depends on meteorological stations. Daily extremes (maximum and / or minimum) might be the only available data. Since temperature does not deviate from its mean level, a mean-reverting process such as an Ornstein-Uhlenbeck (OU) process is commonly used to model temperature process (see [8], [9] for example). The authors of [10] and [2] propose to use an ARMA version of the OU process while [6] propose a fractional Brownian motion (to take into account the long range dependence) instead of the classical Brownian motion in the OU process.

The main purpose of this paper is to estimate the parameters of this OU process. Estimation of OU parameters has been done using observations of the process (see [13]) or more recently using hitting time data in [22] for the neuronal activity. However, weather stations do not record either of these data. That's why we propose an estimation based on daily observed suprema of temperatures. Once the parameter estimation is done, risk measures related to heat waves may be obtained from Monte Carlo simulations of the dynamic of temperatures with the estimated parameters. For example, we would like to estimate the probability of heat waves, namely the probability for outdoor air temperature to exceed a threshold (26.67°C during 3 days, see [15]) or two thresholds (one during night and one during day, see [18]). Other interesting measures would be the corresponding expected area over the threshold or the mean time over the threshold.

Recently, lots of results on the first passage time of the process have been obtained. In [3], different expressions for the density function of the first hitting time to a fixed level by an OU process are given. Since hitting time and suprema are related, the cumulative distribution function (cdf) of the

^{*}This work was supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007) opera operated by the French National Research Agency (ANR).

[†]University of Lyon, CNRS UMR 5208, Ecole Centrale de Lyon, Institut Camille Jordan, France, christophette.blanchet@ec-lyon.fr

[‡]University of Lyon, University Lyon 1, ISFA, LSAF (EA 2429) France, diana.dorobantu@univ-lyon1.fr

[§]University of Lyon, CNRS UMR 5208, Ecole Centrale de Lyon, Institut Camille Jordan, France, laura.gay@ec-lyon.fr

 $[\]label{eq:constraint} \ensuremath{^{\P}}\xspace{University} university of Lyon, CNRS UMR 5208, Institut Camille Jordan, France, veronique.maume@univ-lyon1.fr$

^{II}University of Lyon, CNRS UMR 5208, Institut Camille Jordan, France, pierre.ribereau@univ-lyon1.fr

supremum is obtained.

Unlike classical quantile estimation (such as done in [7] or [23]), we do not use the cdf inverse and propose though a new approach to estimate the parameters. Thanks to the cdf, we perform a least square method to estimate the OU parameters.

The paper is organized as follows. In the next section, the estimation problem is presented. Section 3 is concerned with the theoretical tools. Finally, Section 4 is devoted to the numerical illustrations of the estimation and the related risk measures.

2 Estimation Problem

We suppose that the temperature variations process $X = (X_t)_{t \ge 0}$ is modelled by a stationary OU process given by :

$$dX_t = (\phi_0 - \lambda_0 X_t) dt + \sqrt{\beta_0} dB_t, \quad X_0 \sim \mathcal{N}\left(\frac{\phi_0}{\lambda_0}, \frac{\beta_0}{2\lambda_0}\right)$$

where $\phi_0 \in \mathbb{R}, \lambda_0, \beta_0 \in \mathbb{R}^*_+$, and $(B_t)_{t\geq 0}$ is a standard Brownian motion¹. Suppose that X_0 and $(B_t)_{t\geq 0}$ are independent. We recall that the measure $\mathcal{N}\left(\frac{\phi_0}{\lambda_0}, \frac{\beta_0}{2\lambda_0}\right)$ is the stationary measure. This modelling is reasonable, as, in the applications, we consider observations only from a sub-period of annual observations (e.g. from summer).

Let us note for $s, r \in \mathbb{R}^+$, $S_{[s,r[} = \sup_{s \leqslant t < r} X_t$, $I_{[s,r[} = \inf_{s \leqslant t < r} X_t$ and $\theta_0 = (\phi_0, \lambda_0, \beta_0)$.

Assume that we observe the suprema on a period [0, T] with a partition $(t_i)_{i \ge 0}$ of constant step $h \ge 0$. We then have n suprema $S_{[t_{i-1},t_i[}$ for $i \in [\![1,n]\!]$ on disjoint intervals. Let us remark here that in our problem of daily observations we will take h = 1.

Classical estimation methods are not well suited for the parameter estimation from the supremum observations. Indeed, the likelihood maximization requires the probability density function of the supremum and in order to use quantile methods, one needs to know the supremum's cdf inverse. These two functions can only be obtained by numerical approximations that are more time consuming than numerical methods to get the cdf itself.

This is why we propose to use the cdf of the supremum, denoted F^* , whose expression is given in Proposition 3.3.

Let $N_q \in \mathbb{N}^*$ and s_j , $j = 1, \ldots, N_q$ be real numbers. Let us denote F_n^* the empirical distribution function on the sample $S_{[t_{i-1}, t_i]}$, $i = 1, \ldots, n$.

A natural way to estimate θ_0 is to use a least square method by minimizing the sum of squares of the differences between theoretical and empirical cdf. Then, we want to minimize the following function Q_n :

$$Q_{n}(\theta) = \sum_{j=1}^{N_{q}} \left[F^{*}(s_{j}, \theta, h) - F_{n}^{*}(s_{j}) \right]^{2}$$

where $F^*(a, \theta, h) = \mathbb{P}\left(S_{[0,h]} \leq a\right)$ and s_1, \ldots, s_{N_q} are real numbers (to be chosen later). Thus, θ_0 is estimated by

$$\widehat{\theta_n} = (\widehat{\phi_n}, \widehat{\lambda_n}, \widehat{\beta_n}) = \operatorname*{argmin}_{\theta \in \mathbb{R}_+^* \times \mathbb{R}_+^*} Q_n(\theta) \tag{1}$$

Remark 2.1. The problem is stated here with suprema but the same reasoning may be applied to the infima (or both infima and suprema) to deduce the estimation.

3 Theoretical Tools

In this section, we present some useful results to estimate the parameters.

¹If $\beta_0 = 0$, the process is deterministic. If $\lambda_0 = 0$, then X is a drifted Brownian motion.

3.1 Cdf of the supremum

To minimize the function Q_n , we need to compute the cdf F^* of the supremum. Using the hitting time density for an OU process (see [3]), we can deduce the following result on the conditional cdf.

Proposition 3.1. For $x \in \mathbb{R}$, $t \in \mathbb{R}^+$ and a > x, the cdf F^c of the conditional supremum of the OU process X with parameter $\theta = (\phi, \lambda, \beta) \in \mathbb{R} \times \mathbb{R}^*_+ \times \mathbb{R}^*_+$ is given by

$$F^{c}(a,\theta,t,x) = \mathbb{P}\left(S_{[0,t[} \leqslant a \mid X_{0} = x\right)$$
$$= 1 - \int_{0}^{t\beta} e^{-\frac{\lambda}{2\beta} \left[\left(a - \frac{\phi}{\lambda}\right)^{2} - \left(x - \frac{\phi}{\lambda}\right)^{2} - u\right]} \frac{a - x}{\sqrt{2\pi u^{3}}} e^{-\frac{(a - x)^{2}}{2u}} \mathbb{E}\left[e^{-\frac{\lambda^{2}}{2\beta^{2}} \int_{0}^{u} (r_{s} - a + \frac{\phi}{\lambda})^{2} ds}\right] du$$

where r is a 3-dimensional Bessel bridge over the interval [0, u] between 0 and a - x. For $a \leq x$, $F^{c}(a, \theta, t, x) = 0$.

Proof. Let a > x be given and fixed.

Let set $U_t = X_{\frac{t}{\beta}} - \frac{\phi}{\lambda}$, $\mu = \frac{\lambda}{\beta}$ and $W_t = \sqrt{\beta}B_{\frac{t}{\beta}}$ (which is thus a standard Brownian motion). Then the dynamic of $(U_t)_{t\geq 0}$ is

$$dU_t = -\mu U_t dt + dW_t, \quad U_0 = u_0 = x - \frac{\phi}{\lambda} \in \mathbb{R}$$

For $b > u_0$, we introduce the first passage time $H_b = \inf\{s \ge 0; U_s = b\}$. Since $\mathbb{P}\left(\sup_{0 \le u < t} U_u \le b \mid X_0 = x\right) = \mathbb{P}(H_b > t \mid X_0 = x)$, we have :

$$\mathbb{P}\left(S_{[0,t[} \leqslant a \mid X_0 = x\right) = \mathbb{P}\left(\sup_{0 \leqslant u < t\beta} U_u \leqslant a - \frac{\phi}{\lambda} \mid X_0 = x\right) = \mathbb{P}\left(H_{a - \frac{\phi}{\lambda}} > t\beta \mid X_0 = x\right).$$

We conclude using the density f_{H_b} of H_b given (in [3]) by :

$$f_{H_b}(u) = e^{-\frac{\mu}{2}(b^2 - u_0^2 - u)} \frac{b - u_0}{\sqrt{2\pi u^3}} e^{-\frac{(b - u_0)^2}{2u}} \mathbb{E}\left[e^{\frac{-\mu^2}{2}\int_0^u (r_s - b)^2 \,\mathrm{d}s}\right]$$

where r is a 3-dimensional Bessel bridge over the interval [0, u] between 0 and $b - u_0$. Then, we have

$$\mathbb{P}\left(S_{[0,t]} \leqslant a \mid X_0 = x\right) = 1 - \int_0^{t\beta} \mathrm{e}^{-\frac{\lambda}{2\beta} \left[\left(a - \frac{\phi}{\lambda}\right)^2 - \left(x - \frac{\phi}{\lambda}\right)^2 - u\right]} \frac{a - x}{\sqrt{2\pi u^3}} \,\mathrm{e}^{-\frac{(a - x)^2}{2u}} \,\mathbb{E}\left[\mathrm{e}^{\frac{-\lambda^2}{2\beta^2} \int_0^u \left(r_s - a + \frac{\phi}{\lambda}\right)^2 \,\mathrm{d}s}\right] \mathrm{d}u$$

 \square

with r a 3-dimensional Bessel bridge over the interval [0, u] between 0 and a - x.

Remark 3.2. Similarly, we can obtain the cdf F_c of the conditional infimum of X. For $x \in \mathbb{R}$, $t \in \mathbb{R}^+$, a < x and $\theta \in \mathbb{R} \times \mathbb{R}^*_+ \times \mathbb{R}^*_+$, we have

$$F_c(a,\theta,t,x) = \mathbb{P}\left(I_{[0,t]} \leqslant a \mid X_0 = x\right)$$
$$= \int_0^{t\beta} e^{-\frac{\lambda}{2\beta} \left[\left(a - \frac{\phi}{\lambda}\right)^2 - \left(x - \frac{\phi}{\lambda}\right)^2 - u\right]} \frac{x-a}{\sqrt{2\pi u^3}} e^{-\frac{(a-x)^2}{2u}} \mathbb{E}\left[e^{-\frac{\lambda^2}{2\beta^2} \int_0^u (r_s + a - \frac{\phi}{\lambda})^2 \, \mathrm{d}s}\right] \mathrm{d}u$$

with r a 3-dimensional Bessel bridge over the interval [0, u] between 0 and x - a. For $x \leq a$, $F_c(a, \theta, t, x) = 0$.

Integrating with respect to the law of X_0 , we can express the cdf F^* of the supremum of the OU process X with parameter $\theta = (\phi, \lambda, \beta) \in \mathbb{R} \times \mathbb{R}^*_+ \times \mathbb{R}^*_+$.

Proposition 3.3. For $t \in \mathbb{R}^+$ and $a \in \mathbb{R}$, the cdf F^* is given by

$$\begin{split} F^*(a,\theta,t) &= \mathbb{P}\left(S_{[0,t]} \leqslant a\right) \\ &= \int_{-\infty}^a F^c(a,\theta,t,x) \,\mathrm{d}x \\ &= \Phi\left(\frac{a - \frac{\phi}{\lambda}}{\sqrt{\frac{\beta}{2\lambda}}}\right) - \int_{-\infty}^a \int_0^{t\beta} \mathrm{e}^{-\frac{\lambda}{2\beta} \left[\left(a - \frac{\phi}{\lambda}\right)^2 + \left(x - \frac{\phi}{\lambda}\right)^2 - u\right]} \frac{a - x}{\pi\sqrt{2u^3\frac{\beta}{\lambda}}} \,\mathrm{e}^{-\frac{(a - x)^2}{2u}} \,\mathbb{E}\left[\mathrm{e}^{-\frac{\lambda^2}{2\beta^2} \int_0^u (r_s - a + \frac{\phi}{\lambda})^2 \,\mathrm{d}s}\right] \mathrm{d}u \,\mathrm{d}x \end{split}$$

where r is a 3-dimensional Bessel bridge over the interval [0, u] between 0 and a - x and Φ is the cdf of the standard normal distribution.

3.2 Mixing property

In order to get statistical properties of estimators, some mixing properties are usually required. Indeed, statistics beyond independence have received a deep attention from the 90's. Mixing is used instead of independence and results such as Laws of Large Numbers or Central Limit Theorems may still hold. There is a very large literature on that subject and we refer to [5, 11, 25] and the references therein for definitions and main results.

Roughly speaking, mixing properties of a process $(Y_t)_{t\in\mathbb{R}}$ quantify the convergence to 0 as r goes to infinity of

$$\operatorname{Cov}\left(f(Y_{z_1},\ldots,Y_{z_j}),g(Y_{z_{j+1}},\ldots,Y_{z_\ell})\right)$$

for f and g in an appropriate class of measurable functions and $0 < z_1 < \cdots < z_j \leq z_j + r \leq z_{j+1} < \cdots < z_\ell$. The following proposition means that $(S_{[s,t]})_{0 \leq s < t}$ is exponentially ρ -mixing.

Proposition 3.4 (Mixing property). Let us consider an OU with parameter $\theta = (\phi, \lambda, \beta)$. For any $s, r \ge 0$, for any functions $f : C^0([0,s], \mathbb{R}) \to \mathbb{R}$, $g : C^0([s+r,+\infty], \mathbb{R}) \to \mathbb{R}$ such that f, g are square-integrable with respect to the law of S, and for any $0 \le u \le s \le s + r \le v$, we have

$$\left|\operatorname{Cov}\left[f\left(\left(S_{[0,u[}\right)_{u\leqslant s}\right),g\left(\left(S_{[s+r,v[}\right)_{v\geqslant s+r}\right)\right]\right|\leqslant e^{-\lambda r}\sqrt{\operatorname{Var}\left[f\left(\left(S_{[0,u[}\right)_{u\leqslant s}\right)\right]\operatorname{Var}\left[g\left(\left(S_{[s+r,v[}\right)_{v\geqslant s+r}\right)\right]\right)}\right]\right|$$

Proof. We can easily adapt the proof of Theorem 2.1 in [14] in the one-dimensional case of the OU process satisfying the equation

$$dX_t = (\phi - \lambda X_t) dt + \sqrt{\beta} dB_t, \quad X_0 \sim \mathcal{N}\left(\frac{\phi}{\lambda}, \frac{\beta}{2\lambda}\right)$$

Then, keeping the notation from [14], one may take $\varphi = f \circ \sup$ and $\phi = g \circ \sup$ which are squareintegrable with respect to the law of X by hypothesis.

3.3 Consistency of the estimation

Following the idea of the proof of Theorem II.5.1 in [4], we may prove the consistency of our estimation of the parameter θ_0 provided that the s_j , $j = 1, \ldots, N_q$ are chosen such that the function

$$\Psi: \begin{array}{ccc} \mathbb{R} \times \mathbb{R}^*_+ \times \mathbb{R}^*_+ & \longrightarrow & [0,1]^{N_q} \\ \theta & \longmapsto & (F^*(s_j,\theta,h))_{j=1,\dots,N_q} \end{array}$$

is injective and that the parameter θ belong to a compact subset of $\mathbb{R} \times \mathbb{R}^*_+ \times \mathbb{R}^*_+$.

Proposition 3.5. Consider an OU process with parameters $\theta_0 = (\phi_0, \lambda_0, \beta_0)$. Assume that the parameters θ_0 belong to Θ a compact subset of $\mathbb{R} \times \mathbb{R}^*_+ \times \mathbb{R}^*_+ 3$. For any $n \in \mathbb{N}^*$, let $\hat{\theta}_n = (\hat{\phi}_n, \hat{\lambda}_n, \hat{\beta}_n)$ be given by (1). Then, any a.s. limit point θ^* of $(\hat{\theta}_n)_{n \in \mathbb{N}^*}$ satisfies $\Psi(\theta^*) = \Psi(\theta_0)$.

Proof. We adapt the lines of the proof of Theorem II.5.1 in [4] and use the ergodic theorem for mixing sequences (see [5] e.g.).

We denote

$$\widehat{\theta}_n = \operatorname{argmin}_{\theta \in \Theta} Q_n(\theta).$$

Since Θ is a compact set, the sequence $\hat{\theta}_n$ has limit points. Let θ^* be any limit point of $\hat{\theta}_n$. Since no confusion can be made, and in order to simplify notations, we use $F^*(\cdot, \cdot)$ instead of $F^*(\cdot, \cdot, h)$ for this proof. For $j = 1, \ldots N_q$, let $\varepsilon_n(s_j) = F^*(s_j, \theta_0) - F_n(s_j)$ we write

$$Q_n(\theta) = \sum_{j=1}^{N_q} \varepsilon_n(s_j)^2 + \sum_{j=1}^{N_q} (F^*(s_j, \theta_0) - F^*(s_j, \theta))^2 - 2\sum_{j=1}^{N_q} \varepsilon_n(s_j) (F^*(s_j, \theta_0) - F^*(s_j, \theta)).$$
(2)

The ergodic theorem for mixing sequences implies that $\varepsilon_n(s_j)$ goes to 0 a.e. as n goes to infinity for $j = 1, \ldots, N_q$. Now,

$$Q_n(\widehat{\theta}_n) \leqslant Q_n(\theta_0).$$

Let n_k be a subsequence such that $\hat{\theta}_{n_k}$ goes to θ^* , using (2), we have

$$Q_{n_k}(\widehat{\theta}_{n_k}) \longrightarrow \sum_{j=1}^{N_q} (F^*(s_j, \theta_0) - F^*(s_j, \theta^*))^2$$
 a.e.

and $Q_n(\theta_0) \longrightarrow 0$ a.e. We deduce that

$$F^*(s_j, \theta_0) - F^*(s_j, \theta^*) = 0, \ j = 1, \dots, N_q$$

which gives the announced result.

Remark 3.6. Of course, if the application Ψ is injective, then Proposition 3.5 implies that $\theta^* = \theta_0$ and thus $\hat{\theta}_n$ goes to θ_0 a.s. as n goes to infinity. From some numerical tests, it seems that the injectivity is satisfied but we were unable to prove it.

4 Numerical Applications

In this section, we want to estimate the parameters, first on some simulated data, and then on real ones. First of all, we need to implement the cdf of the supremum of the OU process X with 3 parameters.

From now on, to simplify the study and regarding the form of F^* , we make a new parametrisation for the optimisation problem (1). Instead of working with $\theta = (\phi, \lambda, \beta)$, we use in an equivalent way² $\theta' = \left(\beta, \frac{\phi}{\lambda}, \frac{\lambda}{\beta}\right)$.

Now, functions taking θ for argument are going to be used with θ' as argument.

We want to make an estimation on real data with daily suprema observations. This is why without any precisions, h will be equal to 1 for numerical applications in the rest of the paper.

4.1 Cdf Numerical Computation

We describe here the used method for the numerical computation. Contrary to what is written in [3], the process r is the unique solution of the following SDE ([12])

$$\mathrm{d}r_s = \left(-\frac{r_s}{u-s} + \frac{a-x}{(u-s)\tanh\left(\frac{r_s(a-x)}{u-s}\right)}\right)\mathrm{d}s + \mathrm{d}B_s, \quad 0 < s < u, \quad r_0 = 0.$$

Since the process starts from 0 here, the Euler scheme cannot be applied for this SDE. Recall that the process $(r_s)_{s \leq u}$ with $r_0 = 0$ and $r_u = a - x$ and the process $(\tilde{r}_s)_{s \leq u}$ defined by $\tilde{r}_s = r_{u-s}$ with

²Since $\beta \neq 0$ and $\lambda \neq 0$.

 $\tilde{r}_0 = a - x$ and $\tilde{r}_u = 0$ have same distributions (Exercise XI.3.7 of [24]). Therehence, we can use the Euler scheme on the switched Bessel bridge $(\tilde{r}_s)_{s \leq u}$ which verifies the SDE ([24])

$$\mathrm{d}\tilde{r}_s = \left(-\frac{\tilde{r}_s}{u-s} + \frac{1}{\tilde{r}_s}\right)\mathrm{d}s + \mathrm{d}B_s, \quad 0 < s < u, \quad \tilde{r}_0 = a - x.$$

Finally, the integrals are computed by considering the corresponding Riemann sum and the expectation by a Monte-Carlo method with M = 10000 simulations.

The code is written in C++ and the evaluation of the function is very long. Consequently, we had to make a parallel code. Yet, the function "rand" in C++ is not thread safe. Thus, we propose to use the Mersenne Twister generator for the simulation of the random numbers. With the parallelisation, the time for one evaluation of the function Q_n to be minimized has been divided approximately by 5 but is still long (around 52 secs for $N_q = 4$, $\theta' = (47.5, 22, 0.02)$, with a 40 cores machine). The duration is not a problem since the optimisation needs to be done once and for all.

4.2 Bounding parameters

The problem (1) is solved by an algorithm which performs a Nelder-Mead method. More precisely, we use optim procedure on the software R. Initial values for the parameters to be optimized over are required. To set those initial values, we propose to bound each parameter. For each of them, we give here a lower and an upper bound.

As well as we observe the maxima, suppose we also have the minima : $I_{[t_{i-1},t_i]}$ (still for $(t_i)_{i\geq 0}$ a partition of [0,T] of constant step $h \geq 0$). Then, the available quantities for bounding the parameters are the minima mean, denoted m_{\min} ; the maxima mean, denoted m_{\max} ; the smallest observed temperature, denoted rec_{min} and the largest one, denoted rec_{max}.

Let us recall the OU process is assumed to be stationary, then we have, for all $t \ge 0$,

$$\mathbb{E}[X_t] = \frac{\phi_0}{\lambda_0} \qquad \qquad \text{Var}[X_t] = \frac{\beta_0}{2\lambda_0}$$

The expectation gives us natural bounds for the parameter $\frac{\phi_0}{\lambda_0}$:

$$m_{\min} \leqslant \frac{\phi_0}{\lambda_0} \leqslant m_{\max}$$

Moreover, we have $\frac{\lambda_0}{\beta_0} = \frac{1}{2 \operatorname{Var}[X_t]}$. As for all $i, |X_{t_i} - \bar{X}| \leq \max(|\operatorname{rec}_{\min} - m_{\max}|; |\operatorname{rec}_{\max} - m_{\min}|),$ one may say that it is natural to upper bound the variance by

$$\operatorname{Var}[X_t] \leq (\max(|\operatorname{rec}_{\min} - m_{\max}|; |\operatorname{rec}_{\max} - m_{\min}|))^2$$

It then gives us a lower-bound $\left(\frac{\lambda}{\beta}\right)_{\min}$ for $\frac{\lambda_0}{\beta_0}$. For the upper-bound, we use Theorem 2.7 of [20]. For all x > 0, we have

$$p_x := \mathbb{P}\left(S_{[0,1[} - \mathbb{E}\left[S_{[0,1[}\right] \ge x\right) \le \exp\left(-\frac{\lambda_0 x^2}{\beta_0}\right)\right)$$

Hence

Finally, $\hat{\theta}_n$

$$\frac{\lambda_0}{\beta_0} \leqslant \left(\frac{\lambda}{\beta}\right)_{\max} = \inf_{x>0} \left[\frac{-\ln(p_x)}{x^2}\right].$$

It remains to find the domain of β_0 . First of all, $\beta_0 > 0$. Since $\beta_0 = \frac{\langle X \rangle_T}{T}$, a classical estimator of β_0 (see [19]) is $\hat{\beta}_0 = \frac{1}{T} \sum_{i=0}^{n-1} (X_{t_{i+1}} - X_{t_i})^2$. Then,

$$\beta_0 \leqslant \beta_{\max} = \frac{1}{T} \sum_{i=1}^{n-1} \max \left[\left(S_{[t_i, t_{i+1}[} - I_{[t_{i-1}, t_i[}] \right)^2, \left(I_{[t_i, t_{i+1}[} - S_{[t_{i-1}, t_i[}] \right)^2 \right] \right]$$

= $\operatorname{argmin}_{\theta \in C} Q_n(\theta)$ where $C = [0, \beta_{\max}] \times [m_{\min}, m_{\max}] \times \left[\left(\frac{\lambda}{\beta} \right)_{\min}, \left(\frac{\lambda}{\beta} \right)_{\max} \right] \subset \mathbb{R} \times \mathbb{R}^*_+ \times \mathbb{R}^*_+$

4.3Parameters estimation on simulated data

We are going to test our method on simulated data. To choose realistic parameters, we use some temperatures data. The mean temperature leads us to take $\frac{\phi_0}{\lambda_0} = 22$. Using the difference between the maximal (respectively minimal) temperature and the mean temperature, we take $\sqrt{\frac{\beta_0}{2\lambda_0}} = 5$. The

hourly correlation allows us to set $\lambda_0 = 0.95$ (see e.g. [15] and [6]). Then, $\theta'_0 = (47.5, 22, 0.02)$.

We made several tests to make a compromise between the algorithm complexity and the precision of the estimation (with RMSE) which lead us to take $N_q = 4$ here.

The algorithm is launched on R on 50 samples simulated over T = n = 1000 days for each, with $X_0 \sim \mathcal{N}(22, 25)$. The parameters found minimizing Q_{1000} are presented in the following boxplots:



Figure 1: Boxplots of the estimated parameters

The median parameters are satisfying. Yet, we observe a big variation in the estimators of β_0 . It is confirmed by the relative RMSE which are for β_0 , $\frac{\phi_0}{\lambda_0}$ and $\frac{\lambda_0}{\beta_0}$ respectively equal to 0.462, 0.0418 and 0.0932. Better results are obtained if β_0 is fixed and performing a 2D-estimation, as in [22]. Indeed, the relatives RMSE for $\frac{\phi_0}{\lambda_0}$ and $\frac{\lambda_0}{\beta_0}$ are then respectively equal to 0.0107 and 0.0929. It is consistent with the results in [21] where β_0 is assumed to be known.

Real data **4.4**

4.4.1**Parameters** estimation

In [17], daily temperature dataset in Paris through the ECA&D project is provided (Data and metadata available at http://www.ecad.eu). This dataset is one of the longest in temperature measurement

since it begins in 1900 but it records only maximum, minimum and mean daily temperature. In our application, we study daily summer temperature. In that way, we select maximal and minimal temperatures from 15th of june to the 14th of august (61 days) each year between 1950 and 1984 included, representing 35 years of records and 2135 days. These years are selected in order to avoid climate change influence so we can consider the dataset as maximum observations of a stationary process (see [1]).

When we apply the estimation procedure presented in Section 2, we find $\hat{\theta}'_0 = (34.35, 19.04, 0.02633)$. In order to assess the quality of this estimation, we propose to compare some theoretical quantities with empirical ones. That is done in the next section.

4.4.2 Estimation validation

To verify the estimation, we propose two models validation indicators: : comparison of quantiles and prediction.



Figure 2: Quantile-Quantile plot

The first thing to check is the match of quantiles. To this aim, we draw a quantile-quantile plot (see Figure 2). The plotted points fall near the line y = x which indicates that the quantiles of the theoretical and data distributions agree.

We also want to assess the estimation quality by a prediction method. The estimation ends on the 14/08/1984. We take the mean temperature of the 14/06/1985 as an initial point to simulate processes on 10 days with the estimated parameters. Then, we make a confidence interval for the maxima over each of those days and compare it with the real values (between 15/06/1985 and 24/06/1985).



Figure 3: Confidence limits for the maxima between 15/06/1985 and 24/06/1985

We observe that the real values are all in the confidence interval which confirms the pertinence of our model.

4.4.3 Risk measures

Let us recall the goal of our study. We want to estimate some risk measures related to heat waves. Let us note that one may estimate any risk measures of his choice. Indeed, once the minimisation is performed, we can use a Monte-Carlo method simulating independent processes with estimated parameters.

A heat wave is classically defined by as a sequence of consecutive days (Δ days) for which the maximum daily temperature is larger than a high-level threshold (a_{\max}) and the minimum daily temperature is greater than a low level one (a_{\min}) . Those temperatures thresholds depend on the geographical zone. As we have daily observations, to simplify the expression, we take $t_i = i \in [0, n]$ and $S_{[0,1[}$ is then the supremum on the first day for example.

We define the two random variables

$$m_{[i,i+\delta[}^S = \min\left(S_{[i,i+1[},\ldots,S_{[i+\delta-1,i+\delta[})\right)\right)$$

and

$$m_{[i,i+\delta[}^{I} = \min\left(I_{[i,i+1[},\ldots,I_{[i+\delta-1,i+\delta[})\right)$$

for $\delta \in [\![1, n-i]\!]$.

Then, we can express the probability of heat wave

$$\mathbb{P}\left(\exists i \in \llbracket 0, n-\delta \rrbracket, \ m_{[i,i+\delta[}^S \ge a_{\max} \ , \ m_{[i,i+\delta[}^I \ge a_{\min}\right).\right.$$

Another interesting measure is the duration of an heat wave. Let us note, when there exists

$$\tau_{\rm in} = \min\{i \in [\![0, n-\delta]\!], \ m_{[i,i+\delta[}^S \ge a_{\rm max} \ , \ m_{[i,i+\delta[}^I \ge a_{\rm min}\}\}$$

and

$$\tau_{\text{out}} = \tau_{\text{in}} + \max\{\delta \ge \Delta, \ m_{[i,i+\delta[}^S \ge a_{\max} \ , \ m_{[i,i+\delta[}^I \ge a_{\min}\}.$$

Then, the mean duration of an heat wave is

$$\mathbb{E}\left[\tau_{\text{out}} - \tau_{\text{in}} \mid m^{S}_{[\tau_{\text{in}}, \tau_{\text{out}}[} \geqslant a_{\text{max}}, m^{I}_{[\tau_{\text{in}}, \tau_{\text{out}}[} \geqslant a_{\text{min}}]\right]$$

We use the markers of Météo France for Paris (see [18]), we take $\Delta = 3$, $a_{\min} = 21^{\circ}$ C and $a_{\max} = 31^{\circ}$ C. As we want to estimate the measures for a summer, we take n = 61 days. Those measures are calculated with the estimated parameters on the real data, namely $\hat{\theta}'_0 = (34.35, 19.04, 0.02633)$

For the probability of heat waves, the Monte-Carlo is performed with the simulation of 10^8 years of 61 days and we obtain a probability of 2.57×10^{-2} for a summer. There were 2 heat waves between 1985 to 2011 then a proportion of 7.41×10^{-2} . This highlights the deviation of the temperatures in the last decades, due to climate change ([1]).

With 10^6 simulations for the Monte-Carlo, we obtain a mean duration for an heat wave of 3.2 days. The 2 heat waves had lasted respectively 3 and 10 days.

5 Conclusion and and future research directions

In this paper, a new method to estimate the parameters of an OU process is proposed. Indeed, the proposed method includes a least square estimation based on the suprema observations. To this aim, the cdf of the suprema of an OU is given and theoretical results, including consistency of model parameter estimation, are established.

The numerical applications on real and simulated data prove the goodness of the estimation and its relevance. Risk measures such as the probability of heat wave or the duration of one have been studied and compared with the reality. The proposed model is also able to predict temperatures for a few days.

Some directions for further investigations are summarized as follows. For example, in continuity with this work, obtaining explicit expressions of risk measures may be interesting in the model. To this aim, one may know the joint law of the supremum and the process. Moreover, another interesting estimation for the parameters of the process might be done using Maximum Simulated Likelihood Estimation (see [16]).

References

- IPCC, 2014: Climate Change 2014: Synthesis report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.
- [2] P. Alaton, B. Djehiche, and D. Stillberger. Applied Mathematics in Finance, (9):1–20, 2002.
- [3] L. Alili, P. Patie, and J. L. Pedersen. Representations of first hitting time density of an Ornstein-Uhlenbeck process. *Stochastic Models*, 21(4):967–980, 2005.
- [4] A. Antoniadis, J. Berruyer, and R. Carmona. Régression non linéaire et applications. Collection "Economie et statistiques avancées.": Série Ecole nationale de la statistique et de l'administration économique et Centre d'études des programmes économiques. Economica, 1992.
- [5] P. Billingsley. *Ergodic Theory and Information*. Wiley series in probability and mathematical statistics. Wiley, 1965.
- [6] D. Brody, J. Syroka, and M. Zervos. Dynamical pricing of weather derivatives. *Quantitative Finance*, 2(3):189–198, 2002.
- [7] E. Castillo, A. Hadi, N. Balakrishnan, and J. Sarabia. Extreme Value and Related Models With Applications in Engineering and Science. 01 2005.
- [8] B. Dischel. At least: a model for weather risk, weather risk special report. Energy and Power Risk Management, (March):30–32, 1998.
- [9] B. Dischel. Black-Scholes won't do it, weather risk special report. Energy and Power Risk Management, (Ocober):8–9, 1998.
- [10] F. Dornier and M. Queruel. Caution to the wind, Weather risk special report. Energy Power Risk Management, pages 30–32, 2000.
- [11] P. Doukhan. Mixing. In *Mixing*, pages 15–23. Springer, 1994.
- [12] A. N. Downes and K. Borovkov. First Passage Densities and Boundary Crossing Probabilities for Diffusion Processes. *Methodology and Computing in Applied Probability*, 10(4):621–644, 2008.
- [13] J. C. G. Franco. Maximum likelihood estimation of mean reverting processes. *Real Options Practice*, 2003.
- [14] E. Gobet and G. Matulewicz. Parameter estimation of Ornstein-Uhlenbeck process generating a stochastic graph. *Statistical Inference for Stochastic Processes*, pages 1–25, 2016.
- [15] I. Gringorten. Estimating finite-time maxima and minima of a stationary Gaussian Ornstein-Uhlenbeck process by Monte Carlo simulation. Journal of the American Statistical Association, 63(324):1517–1521, 1968.
- [16] I. Jeliazkov and A. Lloro. Maximum Simulated Likelihood Estimation: Techniques and Applications in Economics, pages 85–100. Springer Berlin Heidelberg, Berlin, Heidelberg, 2011.

- [17] A. M. G. Klein Tank and al. Daily dataset of 20th-century surface air temperature and precipitation series for the European Climate Assessment. Int. J. of Climatol., (22):1441–1453, 2002.
- [18] K. Laaidi, M. Pascal, M. Ledrans, A. Le Tertre, S. Medina, C. Casério, J.-C. Cohen, J. Manach, P. Beaudeau, and P. Empereur Bissonnet. Le système français d'alerte canicule et santé 2004 (sacs 2004) un dispositif intégré au plan national canicule. *Bulletin épidémiologique hebdomadaire*, pages 134–136, 2004.
- [19] A. Le Breton and M. Musiela. A study of a one-dimensional bilinear differential model for stochastic processes (STMA V26 2481). Probability and Mathematical Statistics, 4:91–107, 1984.
- [20] W. V. Li and Q.-M. Shao. Gaussian processes: Inequalities, small ball probabilities and applications. Stochastic Processes : Theory and Methods . Handbook of Statistics, (19):533–597, 2001.
- [21] B. Martin Bibby and M. Sørensen. Martingale estimation functions for discretely observed diffusion processes. *Bernoulli*, 1(1-2):17–39, 03 1995.
- [22] P. Mullowney and S. Iyengar. Parameter estimation for a leaky integrate-and-fire neuronal model from ISI data. *Journal of Computational Neuroscience*, 24(2):179–194, 2008.
- [23] D. Pekasiewicz. Application of quantile methods to estimation of Cauchy distribution parameters. Statistics in Transition, 15(1):133–144, 2014.
- [24] D. Revuz and M. Yor. Continuous martingales and Brownian motion, volume 293. Springer Science & Business Media, 2013.
- [25] E. Rio. Covariance inequalities for strongly mixing processes. Annales de l'IHP, 29(4):587–597, 1993.