



HAL
open science

Finite type invariants of knots in homology 3-spheres with respect to null LP-surgeries

Delphine Moussard

► **To cite this version:**

Delphine Moussard. Finite type invariants of knots in homology 3-spheres with respect to null LP-surgeries. 2017. hal-01620930v1

HAL Id: hal-01620930

<https://hal.science/hal-01620930v1>

Preprint submitted on 22 Oct 2017 (v1), last revised 27 Oct 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Finite type invariants of knots in homology 3-spheres with respect to null LP-surgeries

Delphine Moussard

Abstract

We study a theory of finite type invariants for null-homologous knots in rational homology 3-spheres with respect to null Lagrangian-preserving surgeries. It is an analogue in the setting of the rational homology of the Goussarov-Rozansky theory for knots in integral homology 3-spheres. We give a partial combinatorial description of the graded space associated with our theory and determine some cases when this description is complete. For null-homologous knots in rational homology 3-spheres with a trivial Alexander polynomial, we show that the Kriker lift of the Kontsevich integral and the Lescop equivariant invariant built from integrals in configuration spaces are universal finite type invariants for this theory; in particular it implies that they are equivalent for such knots.

Contents

1	Introduction	2
2	Statement of the results	6
2.1	Filtration defined by null LP-surgeries	6
2.2	Borromean surgeries	8
2.3	Spaces of diagrams	9
2.4	The Lescop invariant and the Kriker invariant	11
2.5	About the map $\psi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\delta)$ and perspectives	13
2.6	The case of knots in \mathbb{Z} -spheres	15
3	Equivariant clasper calculus	16
4	Colored diagrams and Y-links	20
5	The surjective map $\varphi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$	25
6	Extension of the Lescop/Kriker invariant	30

1 Introduction

The notion of finite type invariants was first introduced independently by Goussarov and Vassiliev for the study of invariants of knots in the 3-dimensional sphere S^3 ; in this case, finite type invariants are also called Vassiliev invariants. The discovery of the Kontsevich integral, which is a universal invariant among all finite type invariants of knots in S^3 , revealed that this class of invariants is very prolific. It is known, for instance, that it dominates all Witten-Reshetikhin-Turaev's quantum invariants. The notion of finite type invariants was adapted to the setting of 3-dimensional manifolds by Ohtsuki [Oht96], who introduced the first examples for integral homology 3-spheres, and it has been widely developed and generalized since then. In particular, Goussarov and Habiro independently developed a theory which involves any 3-dimensional manifolds – and their knots – and which contains the Ohtsuki theory for \mathbb{Z} -spheres [GGP01, Hab00b].

In general, the finite type invariants of a set of objects are defined by their polynomial behavior with respect to some elementary move. For Vassiliev invariants of knots in S^3 , this move is the crossing change on a diagram of the knot. For 3-dimensional manifolds, the elementary move is a certain kind of surgery, for instance the Borromean surgery – a Lagrangian-preserving replacement of a genus 3 handlebody – in the Goussarov-Habiro theory.

In [GR04], Garoufalidis and Rozansky studied the theory of finite type invariants for \mathbb{Z} SK-pairs, *i.e.* knots in integral homology 3-spheres, with respect to the so-called null-move, which is a Borromean surgery defined on a handlebody that is null-homologous in the complement of the knot. In this paper, we study a theory of finite type invariants for \mathbb{Q} SK-pairs, *i.e.* null-homologous knots in rational homology 3-spheres (\mathbb{Q} -spheres). Our elementary move is the null Lagrangian-preserving surgery introduced by Lescop [Les13], which is the Lagrangian-preserving replacement of a rational homology handlebody that is null-homologous in the complement of the knot. This latter theory can be understood as an adaptation of the Garoufalidis-Rozansky theory to the setting of the rational homology; a great part of the results in this paper are stated in both settings.

In [Kri00], Kriker constructed a rational lift of the Kontsevich integral of \mathbb{Z} SK-pairs. In [GK04], he proved with Garoufalidis that his construction provides an invariant of \mathbb{Z} SK-pairs. This invariant takes values in a diagram space with a stronger structure than the target diagram space of the Kontsevich integral, hence it is much more structured than the Kontsevich integral, which it lifts. Garoufalidis and Kriker proved in [GK04] that the Kriker invariant satisfies some splitting formulas with respect to the null-move. These formulas imply in particular that the Kriker invariant is a series of finite type invariants of all degrees with respect to the null-move.

It appears that the null-move preserves the Blanchfield module – the Alexander mod-

ule equipped with the Blanchfield form – of the \mathbb{Z} SK-pair. Hence the study of the Garoufalidis-Rozansky theory of finite type invariants can be restricted to a class of \mathbb{Z} SK-pairs with a fixed Blanchfield module. In the case of a trivial Blanchfield module, Garoufalidis and Rozansky gave a combinatorial description of the associated graded space. Together with the splitting formulas of Garoufalidis and Kricker, this proves that the Kricker invariant is a universal finite type invariant of \mathbb{Z} SK-pairs with trivial Blanchfield module with respect to the null-move.

Another universal invariant in this context was constructed by Lescop in [Les11]. Lescop proved in [Les13] that her invariant satisfies the same splitting formulas as the Kricker invariant. Hence the Lescop invariant is also a universal finite type invariant of \mathbb{Z} SK-pairs with trivial Blanchfield module with respect to the null-move. It implies in particular that the Lescop invariant and the Kricker invariant are equivalent for \mathbb{Z} SK-pairs with trivial Blanchfield module. Lescop conjectured in [Les13] that this equivalence holds for knots with any Blanchfield module.

The Lescop invariant is indeed defined for \mathbb{Q} SK-pairs and Lescop’s splitting formulas are stated with respect to general null Lagrangian-preserving surgeries. In [Mou17], the Kricker invariant is extended to \mathbb{Q} SK-pairs and splitting formulas for this invariant with respect to null Lagrangian-preserving surgeries are given. Hence a combinatorial description of the graded space associated with finite type invariants of \mathbb{Q} SK-pairs with respect to null Lagrangian-preserving surgeries would allow to explicit the universality properties of these two invariants and provide a comparison between them, answering the above conjecture of Lescop for general \mathbb{Q} SK-pairs.

In analogy with the integral homology setting, null Lagrangian-preserving surgeries preserve the Blanchfield module defined over \mathbb{Q} and we study finite type invariants of \mathbb{Q} SK-pairs with a fixed Blanchfield module. In the case of a trivial Blanchfield module, we give a complete description of the associated graded space. This description and the above-mentioned splitting formulas imply that the Lescop invariant and the Kricker invariant are both universal finite type invariants of \mathbb{Q} SK-pairs with trivial Blanchfield module, up to degree one invariants given by the cardinality of the first homology group of the \mathbb{Q} -sphere. In particular, the Lescop invariant and the Kricker invariant are equivalent for \mathbb{Q} SK-pairs with trivial Blanchfield module when the cardinality of the first homology group of the \mathbb{Q} -sphere is fixed, as explicated in the next result, which follows from Theorem 2.10.

Theorem 1.1. *Let $Z_{Les} = (Z_{n,Les})_{n \in \mathbb{N}}$ and $Z_{Kri} = (Z_{n,Kri})_{n \in \mathbb{N}}$ denote the Lescop invariant and the Kricker invariant respectively. Let (M, K) and (N, J) be \mathbb{Q} SK-pairs with trivial Blanchfield module such that $H_1(M; \mathbb{Z})$ and $H_1(N; \mathbb{Z})$ have the same cardinality. Then, for any $n \in \mathbb{N}$, $Z_{k,Les}(M, K) = Z_{k,Les}(N, J)$ for all $k \leq n$ if and only if $Z_{k,Kri}(M, K) = Z_{k,Kri}(N, J)$ for all $k \leq n$.*

Let $(\mathfrak{A}, \mathfrak{b})$ be any Blanchfield module with annihilator $\delta \in \mathbb{Q}[t^{\pm 1}]$. The main goal of this paper is to give a combinatorial description of the graded space $\mathcal{G}(\mathfrak{A}, \mathfrak{b}) = \bigoplus_{n \in \mathbb{Z}} \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ associated with finite type invariants of \mathbb{Q} SK-pairs with Blanchfield module $(\mathfrak{A}, \mathfrak{b})$ – precise

definitions are given in the next section. The Lescop or Kricker invariant $Z = (Z_n)_{n \in \mathbb{N}}$ is a family of finite type invariants Z_n of degree n for n even (Z_n is trivial for n odd). For \mathbb{Q} SK-pairs with Blanchfield module $(\mathfrak{A}, \mathfrak{b})$, Z_n takes values in a space $\mathcal{A}_n(\delta)$ of trivalent graphs with edges labelled in $\frac{1}{\delta}\mathbb{Q}[t^{\pm 1}]$. The finiteness properties imply that Z_n induces a map on $\mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$. In order to take into account the degree 1 invariants, we construct from Z an invariant $Z^{aug} = (Z_n^{aug})_{n \in \mathbb{N}}$ of \mathbb{Q} SK-pairs with Z_n^{aug} of degree n . The invariant Z_n^{aug} takes values in a space $\mathcal{A}_n^{aug}(\delta)$ of trivalent graphs as before, which may in addition contain isolated vertices labelled by prime integers. Again by finiteness, Z_n^{aug} induces a map on $\mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$. This leads us to our main question.

Question 1. *Is the map $Z_n^{aug} : \mathcal{G}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n^{aug}(\delta)$ injective ?*

Injectivity of this map for any Blanchfield module $(\mathfrak{A}, \mathfrak{b})$ is equivalent to universality of the invariant Z as a finite type invariant of \mathbb{Q} SK-pairs, up to degree 0 and 1 invariants. This would imply the equivalence of the Lescop invariant and the Kricker invariant when the Blanchfield module and the cardinality of the first homology group of the \mathbb{Q} -sphere are fixed.

To deal with Question 1, we first construct another diagram space $\mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b})$ together with a surjective map $\varphi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$. Then we compose this map with Z_n^{aug} to get a map $\psi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n^{aug}(\delta)$. It appears that this composed map has a simple




Figure 1: Commutative diagram

diagrammatic description. Nevertheless, it is not easy to decide whether it is injective or not in general.

Question 2. *Is the map $\psi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n^{aug}(\delta)$ injective ?*

If Question 2 has a positive answer, then Question 1 also has, and $\mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ is completely described combinatorially by $\varphi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \xrightarrow{\cong} \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$.

Question 2 has a positive answer in the following cases – the last two cases are treated in [AM17]:

- for a trivial Blanchfield module and any value of n ,
- for a Blanchfield module which is a direct sum of N isomorphic Blanchfield modules and $n \leq \frac{2}{3}N$,
- for a Blanchfield module of \mathbb{Q} -dimension 2 and $n = 2$,
- for a Blanchfield module which is a direct sum of two isomorphic Blanchfield modules of \mathbb{Q} -dimension 2 and of order different from $t + 1 + t^{-1}$, and $n = 2$.

In the third case, the map ψ_n is not surjective, whereas in the other cases, it is an isomorphism. In particular, Z_n^{aug} is not surjective in general. Moreover, for a Blanchfield module which is a direct sum of two isomorphic Blanchfield modules of \mathbb{Q} -dimension 2 and of order $t + 1 + t^{-1}$, and $n = 2$, Question 2 has a negative answer [AM17], but Question 1 is open.

The fact that Question 1 remains open while Question 2 does not have a positive answer in general leads us to the following alternative:

- either Question 1 has a positive answer in general, in which case $\mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ is isomorphic to $\psi_n(\mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}))$,
- or we miss some invariant to add to the augmented Lescop/Kricker invariant and the Blanchfield module to get a universal finite type invariant of QSK-pairs.

We also treat the Garoufalidis-Rozansky theory of finite type invariants of ZSK-pairs in the case of a non-trivial Blanchfield module.

Notations For $\mathbb{K} = \mathbb{Z}, \mathbb{Q}$, a \mathbb{K} -sphere (resp. \mathbb{K} -ball, \mathbb{K} -torus, genus g \mathbb{K} -handlebody) is a compact connected oriented 3-manifold with the same homology with coefficients in \mathbb{K} as the standard 3-sphere (resp. 3-ball, solid torus, genus g standard handlebody). A \mathbb{K} SK-pair (M, K) is a pair made of a \mathbb{K} -sphere M and a knot K in M whose homology class in $H_1(M; \mathbb{Z})$ is trivial.

Plan of the paper In Section 2, we introduce the necessary notions and state the main results of the paper. Section 3 is devoted to clasper calculus in the equivariant setting. We apply this calculus in Section 4 to our diagrams. This provides a surjective map from a graded diagram space to the graded space associated with ZSK-pairs with respect to integral null Lagrangian-preserving surgeries. To get a similar map in the case of QSK-pairs, we need further arguments developed in Section 5. In Section 6, we show the universality property of the invariant Z^{aug} which combines the Lescop/Kricker invariant and the cardinality of the first homology group. In Section 7, we answer Question 2 for a Blanchfield module which is a direct sum of N isomorphic Blanchfield modules in degree at most $\frac{2}{3}N$.

2 Statement of the results

2.1 Filtration defined by null LP-surgeries

We first recall the definition of the Alexander module and the Blanchfield form. Let (M, K) be a QSK-pair. Let $T(K)$ be a tubular neighborhood of K . The *exterior* of K is $X = M \setminus \text{Int}(T(K))$. Consider the projection $\pi : \pi_1(X) \rightarrow \frac{H_1(X; \mathbb{Z})}{\text{torsion}} \cong \mathbb{Z}$ and the covering map $p : \tilde{X} \rightarrow X$ associated with its kernel. The covering \tilde{X} is the *infinite cyclic covering* of X . The automorphism group of the covering, $\text{Aut}(\tilde{X})$, is isomorphic to \mathbb{Z} . It acts on $H_1(\tilde{X}; \mathbb{Q})$. Denoting the action of a generator τ of $\text{Aut}(\tilde{X})$ as the multiplication by t , we get a structure of $\mathbb{Q}[t^{\pm 1}]$ -module on $\mathfrak{A}(M, K) = H_1(\tilde{X}; \mathbb{Q})$. This $\mathbb{Q}[t^{\pm 1}]$ -module is called the *Alexander module* of (M, K) . It is a torsion $\mathbb{Q}[t^{\pm 1}]$ -module.

On the Alexander module, the *Blanchfield form*, or *equivariant linking pairing*, $\mathfrak{b} : \mathfrak{A} \times \mathfrak{A} \rightarrow \frac{\mathbb{Q}(t)}{\mathbb{Q}[t^{\pm 1}]}$, is defined as follows. First define the equivariant linking number of two knots. Let J_1 and J_2 be two knots in \tilde{X} such that $p(J_1) \cap p(J_2) = \emptyset$. Let $\delta \in \mathbb{Q}(t)$ be the annihilator of \mathfrak{A} . There is a rational 2-chain S such that $\partial S = \delta(\tau)J_1$. The *equivariant linking number* of J_1 and J_2 is

$$lk_e(J_1, J_2) = \frac{1}{\delta(t)} \sum_{k \in \mathbb{Z}} \langle S, \tau^k(J_2) \rangle t^k,$$

where $\langle \cdot, \cdot \rangle$ stands for the algebraic intersection number. It is well-defined and we have $lk_e(J_1, J_2) \in \frac{1}{\delta(t)} \mathbb{Q}[t^{\pm 1}]$, $lk_e(J_2, J_1)(t) = lk_e(J_1, J_2)(t^{-1})$ and $lk_e(P(\tau)J_1, Q(\tau)J_2)(t) = P(t)Q(t^{-1})lk_e(J_1, J_2)(t)$. Now, if γ (resp. η) is the homology class of J_1 (resp. J_2) in \mathfrak{A} , define $\mathfrak{b}(\gamma, \eta)$ by:

$$\mathfrak{b}(\gamma, \eta) = lk_e(J_1, J_2) \text{ mod } \mathbb{Q}[t^{\pm 1}].$$

The Blanchfield form is *hermitian*: $\mathfrak{b}(\gamma, \eta)(t) = \mathfrak{b}(\eta, \gamma)(t^{-1})$ and $\mathfrak{b}(P(t)\gamma, Q(t)\eta)(t) = P(t)Q(t^{-1})\mathfrak{b}(\gamma, \eta)(t)$ for all $\gamma, \eta \in \mathfrak{A}$ and all $P, Q \in \mathbb{Q}[t^{\pm 1}]$. Moreover, it is *non degenerate* (see Blanchfield in [Bla57]) : $\mathfrak{b}(\gamma, \eta) = 0$ for all $\eta \in \mathfrak{A}$ implies $\gamma = 0$.

The Alexander module of a QSK-pair (M, K) endowed with its Blanchfield form is its *Blanchfield module* denoted by $(\mathfrak{A}, \mathfrak{b})(M, K)$. In the sequel, by a *Blanchfield module* $(\mathfrak{A}, \mathfrak{b})$, we mean a pair $(\mathfrak{A}, \mathfrak{b})$ which can be realized as the Blanchfield module of a QSK-pair. An isomorphism between Blanchfield modules is an isomorphism between the underlying Alexander modules which preserves the Blanchfield form.

We now define LP-surgeries. Note that the boundary of a genus g \mathbb{Q} -handlebody is homeomorphic to the standard genus g surface. The *Lagrangian* \mathcal{L}_A of a \mathbb{Q} -handlebody A is the kernel of the map $i_* : H_1(\partial A; \mathbb{Q}) \rightarrow H_1(A; \mathbb{Q})$ induced by the inclusion. Two \mathbb{Q} -handlebodies A and B have *LP-identified* boundaries if (A, B) is equipped with a homeomorphism $h : \partial A \rightarrow \partial B$ such that $h_*(\mathcal{L}_A) = \mathcal{L}_B$. The Lagrangian of a \mathbb{Q} -handlebody A is indeed a Lagrangian subspace of $H_1(\partial A; \mathbb{Q})$ with respect to the intersection form.

Let M be a \mathbb{Q} -sphere, let $A \subset M$ be a \mathbb{Q} -handlebody and let B be a \mathbb{Q} -handlebody whose boundary is LP-identified with ∂A . Set $M \left(\frac{B}{A} \right) = (M \setminus \text{Int}(A)) \cup_{\partial A = \partial B} B$. We say that the \mathbb{Q} -sphere $M \left(\frac{B}{A} \right)$ is obtained from M by *Lagrangian-preserving surgery*, or *LP-surgery*.

Given a QSK-pair (M, K) , a *\mathbb{Q} -handlebody null in $M \setminus K$* is a \mathbb{Q} -handlebody $A \subset M \setminus K$ such that the map $i_* : H_1(A; \mathbb{Q}) \rightarrow H_1(M \setminus K; \mathbb{Q})$ induced by the inclusion has a trivial image. A *null LP-surgery* on (M, K) is an LP-surgery $\left(\frac{B}{A} \right)$ such that A is null in $M \setminus K$. The QSK-pair obtained by surgery is denoted by $(M, K) \left(\frac{B}{A} \right)$.

Let \mathcal{F}_0 be the rational vector space generated by all QSK-pairs up to orientation-preserving homeomorphism. Let \mathcal{F}_n be the subspace of \mathcal{F}_0 generated by the

$$\left[(M, K); \left(\frac{B_i}{A_i} \right)_{1 \leq i \leq n} \right] = \sum_{I \subset \{1, \dots, n\}} (-1)^{|I|} (M, K) \left(\left(\frac{B_i}{A_i} \right)_{i \in I} \right)$$

for all QSK-pairs (M, K) and all families of \mathbb{Q} -handlebodies $(A_i, B_i)_{1 \leq i \leq n}$, where the A_i are null in $M \setminus K$ and disjoint, and each ∂B_i is LP-identified with the corresponding ∂A_i . Here and in all the article, $|\cdot|$ stands for the cardinality. Since $\mathcal{F}_{n+1} \subset \mathcal{F}_n$, this defines a filtration.

Definition 2.1. A \mathbb{Q} -linear map $\lambda : \mathcal{F}_0 \rightarrow \mathbb{Q}$ is a *finite type invariant of degree at most n of QSK-pairs with respect to null LP-surgeries* if $\lambda(\mathcal{F}_{n+1}) = 0$.

The following result provides a splitting of the filtration $(\mathcal{F}_n)_{n \in \mathbb{N}}$:

Theorem 2.2 ([Mou15] Theorem 1.14). *A null LP-surgery induces a canonical isomorphism between the Blanchfield modules of the involved QSK-pairs. Conversely, for any isomorphism ζ from the Blanchfield module of a QSK-pair (M, K) to the Blanchfield module of a QSK-pair (M', K') , there is a finite sequence of null LP-surgeries from (M, K) to (M', K') which induces the composition of ζ by the multiplication by a power of t .*

For an isomorphism class $(\mathfrak{A}, \mathfrak{b})$ of Blanchfield modules, let $\mathcal{P}(\mathfrak{A}, \mathfrak{b})$ be the set of all QSK-pairs, up to orientation-preserving homeomorphism, whose Blanchfield modules are isomorphic to $(\mathfrak{A}, \mathfrak{b})$. Let $\mathcal{F}_0(\mathfrak{A}, \mathfrak{b})$ be the subspace of \mathcal{F}_0 generated by the QSK-pairs $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $(\mathcal{F}_n(\mathfrak{A}, \mathfrak{b}))_{n \in \mathbb{N}}$ be the filtration defined on $\mathcal{F}_0(\mathfrak{A}, \mathfrak{b})$ by null LP-surgeries. Then, for $n \in \mathbb{N}$, \mathcal{F}_n is the direct sum over all isomorphism classes $(\mathfrak{A}, \mathfrak{b})$ of Blanchfield modules of the $\mathcal{F}_n(\mathfrak{A}, \mathfrak{b})$. Set $\mathcal{G}_n(\mathfrak{A}, \mathfrak{b}) = \mathcal{F}_n(\mathfrak{A}, \mathfrak{b}) / \mathcal{F}_{n+1}(\mathfrak{A}, \mathfrak{b})$ and $\mathcal{G}(\mathfrak{A}, \mathfrak{b}) = \bigoplus_{n \in \mathbb{N}} \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$. We wish to describe the graded space $\mathcal{G}(\mathfrak{A}, \mathfrak{b})$. By Theorem 2.2, $\mathcal{G}_0(\mathfrak{A}, \mathfrak{b}) \cong \mathbb{Q}$. In Section 5, as a consequence of Theorem 2.7, we prove:

Theorem 2.3. *Let $(\mathfrak{A}, \mathfrak{b})$ be a Blanchfield module. Let $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. For any prime integer p , let B_p be a \mathbb{Q} -ball such that $H_1(B_p; \mathbb{Z}) \cong \mathbb{Z}/p\mathbb{Z}$. Then:*

$$\mathcal{G}_1(\mathfrak{A}, \mathfrak{b}) = \bigoplus_{p \text{ prime}} \mathbb{Q} \left[(M, K); \frac{B_p}{B^3} \right].$$

2.2 Borromean surgeries

Let us define a specific type of LP-surgeries.

The *standard Y-graph* is the graph $\Gamma_0 \subset \mathbb{R}^2$ represented in Figure 2. The looped edges of Γ_0 are the *leaves*. The vertex incident to three different edges is the *internal vertex*. With Γ_0 is associated a regular neighborhood $\Sigma(\Gamma_0)$ of Γ_0 in the plane. The surface $\Sigma(\Gamma_0)$ is oriented with the usual convention. This induces an orientation of the leaves and an orientation of the internal vertex, *i.e.* a cyclic order of the three edges. Consider a 3-




Figure 2: The standard Y-graph

manifold M and an embedding $h : \Sigma(\Gamma_0) \rightarrow M$. The image Γ of Γ_0 is a *Y-graph*, endowed with its *associated surface* $\Sigma(\Gamma) = h(\Sigma(\Gamma_0))$. The Y-graph Γ is equipped with the framing induced by $\Sigma(\Gamma)$. A *Y-link* in a 3-manifold is a collection of disjoint Y-graphs.




Figure 3: Y-graph and associated surgery link

Let Γ be a Y-graph in a 3-manifold M . Let $\Sigma(\Gamma)$ be its associated surface. In $\Sigma \times [-1, 1]$, associate with Γ the six-component link L represented in Figure 3. The *Borromean surgery on Γ* is the surgery along the framed link L . The surgered manifold is denoted $M(\Gamma)$. As proved by Matveev in [Mat87], a Borromean surgery can be realised by cutting a genus 3 handlebody (a regular neighborhood of the Y-graph) and regluing it another

way, which preserve the Lagrangian. If (M, K) is a \mathbb{Q} SK-pair and if Γ is an n -component Y-link, null in $M \setminus K$, then $[(M, K); \Gamma] \in \mathcal{F}_0$ denotes the bracket defined by the n disjoint null LP-surgeries on the components of Γ .

For $n \geq 0$, let $\mathcal{G}_n^b(\mathfrak{A}, \mathfrak{b})$ be the subspace of $\mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ generated by the classes of the brackets defined by null Borromean surgeries. The following result is a consequence of Proposition 2.6 and Lemma 2.5.

Proposition 2.4. *For any Blanchfield module $(\mathfrak{A}, \mathfrak{b})$ and any $n \geq 0$, $\mathcal{G}_{2n+1}^b(\mathfrak{A}, \mathfrak{b}) = 0$.*

2.3 Spaces of diagrams

Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. Let $\delta \in \mathbb{Q}(t)$ be the annihilator of \mathfrak{A} . An $(\mathfrak{A}, \mathfrak{b})$ -colored diagram D is a uni-trivalent graph without strut (\downarrow), with the following data:

- trivalent vertices are oriented, where an *orientation of a trivalent vertex* is a cyclic order of the three half-edges that meet at this vertex;
- edges are oriented and colored by $\mathbb{Q}[t^{\pm 1}]$;
- univalent vertices are colored by \mathfrak{A} ;
- for all $v \neq v'$ in the set V of univalent vertices of D , a rational fraction $f_{vv'}^D(t) \in \frac{1}{\delta(t)}\mathbb{Q}[t^{\pm 1}]$ is fixed such that $f_{vv'}^D(t) \bmod \mathbb{Q}[t^{\pm 1}] = \mathfrak{b}(\gamma, \gamma')$, where γ (resp. γ') is the coloring of v (resp. v'), with $f_{v'v}^D(t) = f_{vv'}^D(t^{-1})$.

In the pictures, the orientation of trivalent vertices is given by $\begin{array}{c} \curvearrowright \\ \downarrow \end{array}$. When it does not seem to cause confusion, we write $f_{vv'}$ for $f_{vv'}^D$. The *degree* of a colored diagram is the number of trivalent vertices of its underlying graph. The unique degree 0 diagram is the empty diagram. For $n \geq 0$, set:

$$\tilde{\mathcal{A}}_n(\mathfrak{A}, \mathfrak{b}) = \frac{\mathbb{Q}\langle (\mathfrak{A}, \mathfrak{b})\text{-colored diagrams of degree } n \rangle}{\mathbb{Q}\langle \text{AS, IHX, LE, OR, Hol, LV, EV, LD} \rangle},$$

where the relations AS (anti-symmetry), IHX, LE (linearity for edges), OR (orientation reversal), Hol (holonomy), LV (linearity for vertices), EV (edge-vertex) and LD (linking difference) are described in Figure 4.

The automorphism group $Aut(\mathfrak{A}, \mathfrak{b})$ of the Blanchfield module $(\mathfrak{A}, \mathfrak{b})$ acts on $\tilde{\mathcal{A}}_n(\mathfrak{A}, \mathfrak{b})$ by acting on the colorings of all the univalent vertices of a diagram simultaneously. Denote by Aut the relation which identifies two diagrams obtained from one another by the action of an element of $Aut(\mathfrak{A}, \mathfrak{b})$. Set:

$$\mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) = \tilde{\mathcal{A}}_n(\mathfrak{A}, \mathfrak{b}) / \langle \text{Aut} \rangle \quad \text{and} \quad \mathcal{A}(\mathfrak{A}, \mathfrak{b}) = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}).$$

Since the opposite of the identity is an automorphism of $(\mathfrak{A}, \mathfrak{b})$, we have:

$$\begin{array}{c}
\begin{array}{ccc}
\begin{array}{c} \diagup \\ | \\ \diagdown \end{array} + \begin{array}{c} \diagup \\ | \\ \diagdown \\ | \\ \diagup \end{array} = 0 \\
\text{AS}
\end{array} &
\begin{array}{ccc}
\begin{array}{c} \diagup \\ | \\ \diagdown \\ | \\ \diagup \end{array} - \begin{array}{c} \diagup \\ | \\ \diagdown \\ | \\ \diagup \\ | \\ \diagdown \end{array} + \begin{array}{c} \diagup \\ | \\ \diagdown \\ | \\ \diagup \\ | \\ \diagdown \end{array} = 0 \\
\text{IHX}
\end{array} \\
\\
\begin{array}{ccc}
x \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} P + y \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} Q = \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} xP + yQ &
\begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} P(t) = \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} P(t^{-1}) &
\begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} P \\
\text{LE} & \text{OR} & \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} Q \\
\text{Hol} & & \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} tP \\
\\
x \begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} \gamma_1 + y \begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} \gamma_2 = \begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} x\gamma_1 + y\gamma_2 &
\begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} \gamma \\
\text{LV} & \begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} PQ = \begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} P \\
\text{EV} & & \begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} Q(t)\gamma \\
x f_{vv'}^{D_1}(t) + y f_{vv'}^{D_2}(t) = f_{vv'}^D(t) \quad \forall v' \neq v & & f_{vv'}^{D'}(t) = Q(t) f_{vv'}^D(t) \\
& & \forall v' \neq v \\
\\
\begin{array}{ccc}
\begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} \gamma_1 & \begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} \gamma_2 & \begin{array}{c} \bullet \\ \uparrow \\ | \\ \downarrow \end{array} \gamma \\
v_1 & v_2 & v \\
\begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} 1 & \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} 1 & \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} 1 \\
D & D' & D'' \\
f_{v_1 v_2}^D = f_{v_1 v_2}^{D'} + P & & \\
\text{LD} & &
\end{array}
\end{array}$$

Figure 4: Relations, where $x, y \in \mathbb{Q}$, $P, Q, R \in \mathbb{Q}[t^{\pm 1}]$, $\gamma, \gamma_1, \gamma_2 \in \mathfrak{A}$.

Lemma 2.5. For all $n \geq 0$, $\mathcal{A}_{2n+1}(\mathfrak{A}, \mathfrak{b}) = 0$.

Proposition 4.5 implies:

Proposition 2.6. Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. For all $n \geq 0$, there is a canonical surjective \mathbb{Q} -linear map:

$$\varphi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \twoheadrightarrow \mathcal{G}_n^{\mathfrak{b}}(\mathfrak{A}, \mathfrak{b}).$$

To get a similar surjective map onto $\mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$, we need more general diagrams. An $(\mathfrak{A}, \mathfrak{b})$ -augmented diagram is the union of an $(\mathfrak{A}, \mathfrak{b})$ -colored diagram (its *Jacobi part*) and of finitely many isolated vertices colored by prime integers. The *degree* of an $(\mathfrak{A}, \mathfrak{b})$ -augmented diagram is the number of its vertices of valence 0 or 3. Set:

$$\mathcal{A}_n^{\text{aug}}(\mathfrak{A}, \mathfrak{b}) = \frac{\mathbb{Q}\langle (\mathfrak{A}, \mathfrak{b})\text{-augmented diagrams of degree } n \rangle}{\mathbb{Q}\langle \text{AS, IHX, LE, OR, Hol, LV, EV, LD, Aut} \rangle} \quad \text{for } n \geq 0,$$

$$\mathcal{A}^{aug}(\mathfrak{A}, \mathfrak{b}) = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}).$$

In Section 5, we prove:

Theorem 2.7. *Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. For all $n \geq 0$, there is a canonical surjective \mathbb{Q} -linear map:*

$$\varphi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \twoheadrightarrow \mathcal{G}_n(\mathfrak{A}, \mathfrak{b}).$$

The following theorem is a consequence of Theorem 2.7 and Theorem 2.10.

Theorem 2.8. *Let \mathfrak{A}_0 be the trivial Blanchfield module. We have a graded space isomorphism $\mathcal{G}(\mathfrak{A}_0) \cong \mathcal{A}^{aug}(\mathfrak{A}_0)$.*

2.4 The Lescop invariant and the Kricker invariant

In order to introduce the Kricker invariant of [GK04] and the Lescop invariant of [Les11], we first define the graded space $\mathcal{A}(\delta)$ where they take values and we relate it to the graded space $\mathcal{A}(\mathfrak{A}, \mathfrak{b})$.

Let $\delta \in \mathbb{Q}[t^{\pm 1}]$. A δ -colored diagram is a trivalent graph whose vertices are oriented and whose edges are oriented and colored by $\frac{1}{\delta(t)}\mathbb{Q}[t^{\pm 1}]$. The degree of a δ -colored diagram is the number of its vertices. Set:

$$\mathcal{A}_n(\delta) = \frac{\mathbb{Q}\langle \delta\text{-colored diagrams of degree } n \rangle}{\mathbb{Q}\langle \text{AS, IHX, LE, OR, Hol, Hol}' \rangle},$$

where the relations AS, IHX, LE, OR, Hol are represented Figure 4 and the relation Hol' is represented Figure 5. Here the relations LE, OR and Hol hold with edges labelled in $\frac{1}{\delta(t)}\mathbb{Q}[t^{\pm 1}]$. Note that in the case of $\mathcal{A}_n(\mathfrak{A}, \mathfrak{b})$, the relation Hol' is induced by the relations Hol, EV and LD. Since any trivalent graph has an even number of vertices, we have

Figure 5: Relation Hol', with $f, g \in \frac{1}{\delta(t)}\mathbb{Q}[t^{\pm 1}]$.

$\mathcal{A}_{2n+1}(\delta) = 0$ for all $n \geq 0$.

With an $(\mathfrak{A}, \mathfrak{b})$ -colored diagram D of degree n , we associate a δ -colored diagram $\tilde{\psi}_n(D)$. Let V be the set of univalent vertices of D . A *pairing* of V is an involution of V with no fixed point. Let \mathfrak{p} be the set of pairings of V . Fix $p \in \mathfrak{p}$. Define a δ -colored diagram $p(D)$

in the following way. If $v \in V$ and $v' = p(v)$, replace in D the vertices v and v' , and their adjacent edges, by a colored edge, as indicated in Figure 6. Now set:

$$\tilde{\psi}_n(D) = \sum_{p \in \mathfrak{p}} p(D).$$

Note that $\tilde{\psi}_n(D) = 0$ when the number of univalent vertices is odd. We obtain a \mathbb{Q} -linear




Figure 6: Pairing of vertices

map $\tilde{\psi}_n$ from the rational vector space freely generated by the $(\mathfrak{A}, \mathfrak{b})$ -colored diagrams of degree n to $\mathcal{A}_n(\delta)$. One easily checks that $\tilde{\psi}_n$ induces a map:

$$\psi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\delta).$$

The disjoint union of diagrams defines a multiplicative operation on $\mathcal{A}(\delta) = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n(\delta)$ which endows it with a graded algebra structure. Denote by \exp_{\sqcup} the exponential map with respect to this multiplication.

The following result asserts the existence and the properties of an invariant Z which may be either the Lescop invariant or the Kricker invariant. Although it is not known whether they are equal or not, they both satisfy the properties of the theorem. In the sequel, we will refer to “the invariant Z ”.

Theorem 2.9 ([Les11, Les13, Kri00, GK04, Mou17]). *There is an invariant $Z = (Z_n)_{n \in \mathbb{N}}$ of $\mathbb{Q}SK$ -pairs with the following properties.*

- If (M, K) is a $\mathbb{Q}SK$ -pair with Blanchfield module $(\mathfrak{A}, \mathfrak{b})$, then $Z_n(M, K) \in \mathcal{A}_n(\delta)$, where δ is the annihilator of \mathfrak{A} .
- Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. Let δ be the annihilator of \mathfrak{A} . The \mathbb{Q} -linear extension of $Z_n : \mathcal{P}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\delta)$ to $\mathcal{F}_0(\mathfrak{A}, \mathfrak{b})$ vanishes on $\mathcal{F}_{n+1}(\mathfrak{A}, \mathfrak{b})$ and $Z_n \circ \varphi_n = \psi_n$, where $\varphi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n^b(\mathfrak{A}, \mathfrak{b})$ is the surjection of Proposition 2.6.
- Let $p_n^c : \mathcal{A}_n(\delta) \rightarrow \mathcal{A}_n(\delta)$ be the map which sends a connected diagram to itself and non-connected diagrams to 0. Set $Z_n^c = p_n^c \circ Z_n$ and $Z^c = \sum_{n > 0} Z_n^c$. Then Z^c is additive under connected sum and $Z = \exp_{\sqcup}(Z^c)$.

Note that, in particular, if the map ψ_n is injective, then the map φ_n is an isomorphism.

In order to take into account the whole quotient $\mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$, we extend the invariant Z . Define a δ -augmented diagram as the disjoint union of a δ -colored diagram with finitely many isolated vertices colored by prime integers. The *degree* of such a diagram is the number of its vertices. Set:

$$\mathcal{A}_n^{aug}(\delta) = \frac{\mathbb{Q}\langle \delta\text{-augmented diagrams of degree } n \rangle}{\mathbb{Q}\langle \text{AS, IHX, LE, OR, Hol, Hol}' \rangle}.$$

The map ψ_n naturally extends to a map $\psi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n^{aug}(\delta)$ preserving the isolated vertices. We now define an invariant $Z^{aug} = (Z_n^{aug})_{n \in \mathbb{N}}$ of QSK-pairs such that the \mathbb{Q} -linear extension of Z_n^{aug} to $\mathcal{F}_0(\mathfrak{A}, \mathfrak{b})$ takes values in $\mathcal{A}_n^{aug}(\delta)$, from which the invariant Z is recovered by forgetting the isolated vertices. For a prime integer p , define an invariant ρ_p by $\rho_p(M, K) = -v_p(|H_1(M; \mathbb{Z})|) \cdot \bullet_p$, where v_p is the p -adic valuation. Once again, the disjoint union makes $\mathcal{A}^{aug}(\delta) = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n^{aug}(\delta)$ a graded algebra. Set:

$$Z^{aug} = Z \sqcup \exp_{\sqcup} \left(\sum_{p \text{ prime}} \rho_p \right).$$

In Section 6, we prove:

Theorem 2.10. *Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. Let δ be the annihilator of \mathfrak{A} . Consider the surjection $\varphi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ of Theorem 2.7 and the map $\psi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n^{aug}(\delta)$. Then the \mathbb{Q} -linear extension of $Z_n^{aug} : \mathcal{P}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n^{aug}(\delta)$ to $\mathcal{F}_0(\mathfrak{A}, \mathfrak{b})$ vanishes on $\mathcal{F}_{n+1}(\mathfrak{A}, \mathfrak{b})$ and $Z_n^{aug} \circ \varphi_n = \psi_n$.*

If the Alexander module is trivial, the relations LV and LD allow to express the elements of $\mathcal{A}_n(\mathfrak{A}, \mathfrak{b})$ without diagrams with univalent vertices. Hence ψ_n is an isomorphism and Theorems 2.8 and 1.1 follow. In general, note that “the map $\psi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n^{aug}(\delta)$ is injective” is equivalent to “the map $\psi_k : \mathcal{A}_k(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_k(\delta)$ is injective for all $k \leq n$ ”. Hence we focus on the study of injectivity of the map $\psi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\delta)$.

2.5 About the map $\psi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\delta)$ and perspectives

We now state a result about the injectivity of the map $\psi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\delta)$ for n even.

In Section 7, we prove:

Theorem 2.11. *Let n be an even positive integer and $N \geq 3n/2$. Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. Let δ be the annihilator of \mathfrak{A} . Define the Blanchfield module $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ as the direct sum of N copies of $(\mathfrak{A}, \mathfrak{b})$. Then the map $\overline{\psi}_n : \mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}}) \rightarrow \mathcal{A}_n(\delta)$ is an isomorphism.*

This result provides a rewriting of the map ψ_n in the general case. We have a natural map $\iota_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ defined on a diagram by interpreting the labels of its

univalent vertices as elements of the first copy of $(\mathfrak{A}, \mathfrak{b})$ in $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$. The following diagram commutes.

$$\begin{array}{ccc}
 & & \mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}}) \\
 & \nearrow \iota_n & \downarrow \cong \overline{\psi}_n \\
 \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) & & \mathcal{A}_n(\delta) \\
 & \searrow \psi_n &
 \end{array}$$

We mention here results from [AM17] about the map $\psi_2 : \mathcal{A}_2(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_2(\delta)$ for small Alexander modules.

Proposition 2.12 ([AM17]). *If $\dim_{\mathbb{Q}}(\mathfrak{A}) = 2$, then ψ_2 is injective but not surjective.*

Proposition 2.13 ([AM17]). *If \mathfrak{A} is the direct sum of two isomorphic Blanchfield modules of \mathbb{Q} -dimension 2 with annihilator δ , then ψ_2 is injective if and only if $\delta \neq t + 1 + t^{-1}$. In this case, it is an isomorphism.*

Perspectives. As mentioned in the introduction, our main goal in this paper is to study Question 1 in order to determine if the Lescop/Kricker invariant Z is a universal finite type invariant of \mathbb{Q} SK-pairs up to degree 0 and 1 invariants. Theorem 2.11 provides the following rewriting of this question.

We have a map $\mathcal{A}_n((\mathfrak{A}, \mathfrak{b})^{\oplus k}) \rightarrow \mathcal{A}_n((\mathfrak{A}, \mathfrak{b})^{\oplus k+1})$ defined by viewing the labels of the univalent vertices in the direct sum of the first k copies of $(\mathfrak{A}, \mathfrak{b})$ in $(\mathfrak{A}, \mathfrak{b})^{\oplus k+1}$. We also have a map $C_n : \mathcal{G}_n^b((\mathfrak{A}, \mathfrak{b})^{\oplus k}) \rightarrow \mathcal{G}_n^b((\mathfrak{A}, \mathfrak{b})^{\oplus k+1})$ induced by the connected sum with a fixed \mathbb{Q} SK-pair $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Using Theorem 2.2, one can check that the map C_n is independent of the fixed pair (M, K) . These maps provide the following commutative diagram, where the vertical arrows are the maps φ_n and Z_n , for any integer N such that $N \geq \frac{3n}{2}$.

$$\begin{array}{ccccccc}
 \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) & \rightarrow & \cdots & \rightarrow & \mathcal{A}_n((\mathfrak{A}, \mathfrak{b})^{\oplus k}) & \rightarrow & \cdots & \rightarrow & \mathcal{A}_n((\mathfrak{A}, \mathfrak{b})^{\oplus N}) \\
 \downarrow & & & & \downarrow & & & & \downarrow \cong \\
 \mathcal{G}_n^b(\mathfrak{A}, \mathfrak{b}) & \rightarrow & \cdots & \rightarrow & \mathcal{G}_n^b((\mathfrak{A}, \mathfrak{b})^{\oplus k}) & \rightarrow & \cdots & \rightarrow & \mathcal{G}_n^b((\mathfrak{A}, \mathfrak{b})^{\oplus N}) \\
 & & & & \downarrow & & & & \downarrow \cong \\
 & & & & \mathcal{A}_n(\delta) & & & &
 \end{array}$$

Lemma 2.14. *The map $Z_n : \mathcal{G}_n^b((\mathfrak{A}, \mathfrak{b})^{\oplus k}) \rightarrow \mathcal{A}_n(\delta)$ is injective for all k if and only if $C_n : \mathcal{G}_n^b((\mathfrak{A}, \mathfrak{b})^{\oplus k}) \rightarrow \mathcal{G}_n^b((\mathfrak{A}, \mathfrak{b})^{\oplus k+1})$ is injective for all k . This assertion for all $(\mathfrak{A}, \mathfrak{b})$ and*

all n is equivalent to the assertion that the space of finite type invariants of \mathbb{Q} SK-pairs is generated as an algebra by degree 0 invariants and invariants that are additive under connected sum.

2.6 The case of knots in \mathbb{Z} -spheres

A great part of the results stated up to that point have an equivalent in the case of \mathbb{Z} SK-pairs. In this subsection, we adapt the definitions and state the results in this case.

Given a \mathbb{Z} SK-pair (M, K) and the infinite cyclic covering \tilde{X} of the exterior of K in M , define the *integral Alexander module* of (M, K) as the $\mathbb{Z}[t^{\pm 1}]$ -module $\mathfrak{A}_{\mathbb{Z}}(M, K) = H_1(\tilde{X}; \mathbb{Z})$ and the *Blanchfield form* $\mathfrak{b}_{\mathbb{Z}}$ on this module. The integral Alexander module of a \mathbb{Q} SK-pair (M, K) endowed with its Blanchfield form is its *integral Blanchfield module* denoted by $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})(M, K)$. In the sequel, by an *integral Blanchfield module*, we mean a pair $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ which can be realized as the integral Blanchfield module of a \mathbb{Z} SK-pair.

Replacing \mathbb{Q} by \mathbb{Z} in the definitions of Subsection 2.1, define *integral Lagrangians*, *integral LP-surgeries* and *integral null LP-surgeries*. Note that a Borromean surgery is an integral LP-surgery.

For diagram spaces, we have to adapt the relation Aut . Given $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$, set $(\mathfrak{A}, \mathfrak{b}) = \mathbb{Q} \otimes (\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$. Define the relation $\text{Aut}_{\mathbb{Z}}$ on $(\mathfrak{A}, \mathfrak{b})$ -colored diagrams as the relation Aut restricted to the action of the automorphisms in $\text{Aut}(\mathfrak{A}, \mathfrak{b})$ that are induced by automorphisms of the $\mathbb{Z}[t^{\pm 1}]$ -module $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$. Set:

$$\mathcal{A}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) = \tilde{\mathcal{A}}_n(\mathfrak{A}, \mathfrak{b}) / \langle \text{Aut}_{\mathbb{Z}} \rangle \quad \text{and} \quad \mathcal{A}^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}).$$

Since the opposite of the identity is an automorphism of $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$, we have:

Lemma 2.15. *For all $n \geq 0$, $\mathcal{A}_{2n+1}^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) = 0$.*

The filtration $(\mathcal{F}_n)_{n \in \mathbb{N}}$ of Subsection 2.1 generalizes the following filtration introduced by Garoufalidis and Rozansky in [GR04]. Let $\mathcal{F}_0^{\mathbb{Z}}$ be the rational vector space generated by all \mathbb{Z} SK-pairs, up to orientation-preserving homeomorphism. Define a filtration $(\mathcal{F}_n^{\mathbb{Z}})_{n \in \mathbb{N}}$ of $\mathcal{F}_0^{\mathbb{Z}}$ by means of null Borromean surgeries.

Remark: Habegger [Hab00a, Theorem 2.5] and Auclair and Lescop [AL05, Lemma 4.11] proved that two \mathbb{Z} -handlebodies whose boundaries are LP-identified can be obtained from one another by a finite sequence of Borromean surgeries. Therefore, the filtration defined on $\mathcal{F}_0^{\mathbb{Z}}$ by integral null LP-surgeries is equal to the filtration $(\mathcal{F}_n^{\mathbb{Z}})_{n \in \mathbb{N}}$.

Theorem 2.16 ([Mou15] Theorem 1.15). *An integral null LP-surgery induces a canonical isomorphism between the integral Blanchfield modules of the involved \mathbb{Z} SK-pairs. Conversely, for any isomorphism ζ from the integral Blanchfield module of a \mathbb{Z} SK-pair (M, K) to the integral Blanchfield module of a \mathbb{Z} SK-pair (M', K') , there is a finite sequence of integral null LP-surgeries from (M, K) to (M', K') which induces the composition of ζ by the multiplication by a power of t .*

This result provides a splitting of the filtration $(\mathcal{F}_n^{\mathbb{Z}})_{n \in \mathbb{N}}$ as the direct sum of filtrations $(\mathcal{F}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}))_{n \in \mathbb{N}}$ of subspaces $\mathcal{F}_0^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ of $\mathcal{F}_0^{\mathbb{Z}}$, where $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ runs along all isomorphism classes of integral Blanchfield modules. Set $\mathcal{G}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) = \mathcal{F}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) / \mathcal{F}_{n+1}^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ and $\mathcal{G}^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) = \bigoplus_{n \in \mathbb{N}} \mathcal{G}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$. Theorem 2.16 implies $\mathcal{G}_0^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) = \mathbb{Q}$. In [GR04], Garoufalidis and Rozansky identified the graded space $\mathcal{G}^{\mathbb{Z}}(\mathfrak{A}_0)$, where \mathfrak{A}_0 is the trivial Blanchfield module, with the graded space $\mathcal{A}^{\mathbb{Z}}(\mathfrak{A}_0)$. Theorem 2.8 generalizes this result.

Proposition 4.6 implies:

Theorem 2.17. *Fix an integral Blanchfield module $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$. For all $n \geq 0$, there is a canonical surjective \mathbb{Q} -linear map:*

$$\varphi_n^{\mathbb{Z}} : \mathcal{A}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) \twoheadrightarrow \mathcal{G}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}).$$

Corollary 2.18. *Fix an integral Blanchfield module $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ and an integer $n \geq 0$. Then $\mathcal{G}_{2n+1}^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) = 0$.*

As in Subsection 2.4, we have a map $\psi_n^{\mathbb{Z}} : \mathcal{A}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) \rightarrow \mathcal{A}_n(\delta)$, where δ is the annihilator of $\mathfrak{A} = \mathbb{Q} \otimes \mathfrak{A}_{\mathbb{Z}}$. Theorem 2.10 implies that the degree n part of the invariant Z provides a \mathbb{Q} -linear map $Z_n : \mathcal{F}_0^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) \rightarrow \mathcal{A}_n(\delta)$ such that $Z_n \circ \varphi_n^{\mathbb{Z}} = \psi_n^{\mathbb{Z}}$.

Set $\mathfrak{b} = id_{\mathbb{Q}} \otimes \mathfrak{b}_{\mathbb{Z}}$. We have a natural projection $\mathcal{A}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) \rightarrow \mathcal{A}_n(\mathfrak{A}, \mathfrak{b})$. The map $\psi_n^{\mathbb{Z}}$ is the composition of the map ψ_n with this projection. Hence we could adapt Theorem 2.11 and get a surjective map $\overline{\psi}_n^{\mathbb{Z}}$, but we would not get injectivity, which is what we are mostly interested in.

3 Equivariant clasper calculus

For a \mathbb{Q} SK-pair (M, K) , let $\mathcal{F}_0^b(M, K)$ be the rational vector space generated by all the \mathbb{Q} SK-pairs that can be obtained from (M, K) by a finite sequence of null Borromean surgeries, up to orientation-preserving homeomorphism. For $n > 0$, let $\mathcal{F}_n^b(M, K)$ be the subspace of $\mathcal{F}_0^b(M, K)$ generated by the $[(M, K); \Gamma]$ for all m -component null Y -links with $m \geq n$.

Lemma 3.1 ([GGP01] Lemma 2.2). *Let Γ be a Y -graph in a 3-manifold V , which has a 0-framed leaf that bounds a disk in V whose interior does not meet Γ . Then $V(\Gamma) \cong V$.*

Lemma 3.2 ([GGP01] Theorem 3.1, [Auc06] Lemma 5.1.1). *Let $\Gamma_0, \Gamma_1, \Gamma_2$ be the Y -graphs drawn in a genus 4 handlebody in Figure 7. Assume this handlebody is embedded in a 3-manifold V . Then $V(\Gamma_0) \cong V(\Gamma_1 \cup \Gamma_2)$.*

Lemma 3.3. *Let Γ be an n -component Y -link null in $M \setminus K$. Let J be a framed knot, rationally null-homologous in $M \setminus K$, and disjoint from Γ . Let Γ' be obtained from Γ by sliding an edge of Γ along J (see Figure 8). Then $[(M, K); \Gamma] = [(M, K); \Gamma'] \pmod{\mathcal{F}_{n+1}^b(M, K)}$.*




Figure 7: Topological equivalence for edge sliding




Figure 8: Sliding an edge

Proof. Let Γ'_0 be the component of Γ' that contains the slided edge and let Γ_0 be the corresponding component of Γ . By Lemma 3.2, the surgery on Γ'_0 is equivalent to the simultaneous surgeries on Γ_0 and on a null Y-graph $\hat{\Gamma}_0$ which has a leaf which is a meridian of a leaf of Γ_0 . It follows that $[(M, K); \Gamma] - [(M, K); \Gamma'] = [(M, K); \Gamma \cup \hat{\Gamma}_0] \in \mathcal{F}_{n+1}^b(M, K)$. \square

In particular, the above lemma shows that the class of $[(M, K); \Gamma] \bmod \mathcal{F}_{n+1}^b(M, K)$ is invariant under an integral number of twists of the edges.

Lemma 3.4 ([GGP01] Theorem 3.1). *Let $\Gamma_0, \Gamma_1, \Gamma_2$ be the Y-graphs drawn in a genus 4 handlebody in Figure 9. Assume this handlebody is embedded in a 3-manifold V . Then $V(\Gamma_0) \cong V(\Gamma_1 \cup \Gamma_2)$.*

Lemma 3.5. *Let Γ be an n -component Y-link null in $M \setminus K$. Let ℓ be a leaf of Γ . Let γ be a framed arc starting at the vertex incident to ℓ and ending in another point of ℓ , embedded in $M \setminus K$ as the core of a band glued to the associated surface of Γ as shown in Figure 10. The arc γ splits the leaf ℓ into two leaves ℓ' and ℓ'' . Denote by Γ' (resp. Γ'') the Y-link obtained from Γ by replacing the leaf ℓ by ℓ' (resp. ℓ''). If ℓ' and ℓ'' are rationally null-homologous in $M \setminus K$, then Γ' and Γ'' are null Y-links and $[(M, K); \Gamma] = [(M, K); \Gamma'] + [(M, K); \Gamma''] \bmod \mathcal{F}_{n+1}^b(M, K)$.*




Figure 9: Topological equivalence for leaf cutting




Figure 10: Cutting a leaf

Proof. Let Γ_0 (resp. Γ'_0, Γ''_0) be the component of Γ (resp. Γ', Γ'') that contains the leaf ℓ (resp. ℓ', ℓ''). By Lemma 3.4, the surgery on Γ_0 is equivalent to simultaneous surgeries on Γ'_0 and on a null Y-graph $\hat{\Gamma}'_0$ obtained from Γ'_0 by sliding an edge along ℓ'' . Set $\hat{\Gamma}' = (\Gamma \setminus \Gamma_0) \cup \hat{\Gamma}'_0$. We have $[(M, K); \hat{\Gamma}'] + [(M, K); \Gamma''] - [(M, K); \Gamma] = [(M, K); (\Gamma \setminus \Gamma_0) \cup \hat{\Gamma}'_0 \cup \Gamma''_0] \in \mathcal{F}_{n+1}^b(M, K)$. Conclude with Lemma 3.3. \square

The next lemma is a consequence of [GGP01, Lemma 4.8].

Lemma 3.6. *Let Γ be an n -component Y-link null in $M \setminus K$. If a leaf ℓ of Γ bounds a disk in $M \setminus (\Gamma \setminus \ell)$ and has framing 1, i.e. the linking number of ℓ with its parallel induced by the framing of Γ is 1, then $[(M, K); \Gamma] = 0 \text{ mod } \mathcal{F}_{n+1}^b(M, K)$.*

The above two lemmas imply that the class of $[(M, K); \Gamma] \text{ mod } \mathcal{F}_{n+1}^b(M, K)$ does not depend on the framing of the leaves of Γ .

Lemma 3.7. *Let Γ be an n -component Y-link null in $M \setminus K$. Let ℓ be a leaf of Γ . Assume $\Gamma \setminus \ell$ is fixed. Then $[(M, K); \Gamma] \text{ mod } \mathcal{F}_{n+1}^b(M, K)$ only depends on the homotopy class of ℓ in $(M \setminus K) \setminus (\Gamma \setminus \ell)$.*

Proof. If the leaf ℓ is modified by an isotopy in $(M \setminus K) \setminus (\Gamma \setminus \ell)$, then the homeomorphism class of $(M, K)(\Gamma)$ is preserved. If the leaf ℓ crosses itself during a homotopy, apply

Lemma 3.5, as shown in Figure 11, and conclude that $[(M, K); \Gamma] \bmod \mathcal{F}_{n+1}^b(M, K)$ is unchanged by applying Lemma 3.1. \square




Figure 11: Self-crossing of a leaf

Lemma 3.8. *Let Γ be an n -component Y -link null in $M \setminus K$. Let ℓ be a leaf of Γ . Let Γ' be an n -component null Y -link such that $\Gamma' \setminus \ell'$ coincides with $\Gamma \setminus \ell$, where ℓ' is a leaf of Γ' . Let $\widetilde{\Gamma \setminus \ell}$ be the preimage of $\Gamma \setminus \ell$ in the infinite cyclic covering \widetilde{X} associated with (M, K) . Let $\tilde{\ell}$ and $\tilde{\ell}'$ be lifts of ℓ and ℓ' respectively, with the same basepoint. If ℓ and ℓ' are homotopic in $M \setminus K$ and $\tilde{\ell}$ and $\tilde{\ell}'$ are rationally homologous in $\widetilde{X} \setminus (\widetilde{\Gamma \setminus \ell})$, then $[(M, K); \Gamma] = [(M, K); \Gamma'] \bmod \mathcal{F}_{n+1}^b(M, K)$.*

Proof. Consider a homotopy from ℓ to ℓ' in $M \setminus K$. Thanks to Lemma 3.7, it suffices to treat the case when the leaf crosses some edges or leaves of $\Gamma \setminus \ell$ during the homotopy. As shown in Figure 12, Lemma 3.5 implies that the bracket $[(M, K); \Gamma]$ is then added




Figure 12: Crossing of an edge or a leaf

brackets $[(M, K); \hat{\Gamma}]$, where $\hat{\Gamma}$ is the null Y -link obtained from Γ by adding the cutting arc to the edge adjacent to ℓ , and by replacing ℓ by a meridian of the crossed edge or leaf. In the case of a meridian of an edge, Lemmas 3.1 and 3.3 show that the added bracket vanishes.

Fix a leaf ℓ_0 of $\Gamma \setminus \ell$. Let the $[(M, K); \hat{\Gamma}_i]$, for $i \in I$, be the brackets added during the homotopy when the leaf ℓ crosses the leaf ℓ_0 . In each $\hat{\Gamma}_i$, pull the basepoint of the

leaf replacing the leaf ℓ onto the initial basepoint of ℓ . Let ℓ_i be the obtained leaf. Let $\tilde{\ell}_i$ be the lift of ℓ_i which has the same basepoint as $\tilde{\ell}$. Let Y be the complement in \tilde{X} of the preimage of ℓ_0 . In $H_1(Y; \mathbb{Q})$, we have $\tilde{\ell} = \sum_{i \in I} \tilde{\ell}_i + \tilde{\ell}'$. Since $\tilde{\ell}$ and $\tilde{\ell}'$ are homologous in $\tilde{X} \setminus (\widetilde{\Gamma \setminus \ell})$, it implies that $\sum_{i \in I} lk_e(\tilde{\ell}_i, \tilde{\ell}_0) = 0$, where $\tilde{\ell}_0$ is a lift of ℓ_0 . By construction of the $\tilde{\ell}_i$, each $lk_e(\tilde{\ell}_i, \tilde{\ell}_0)$ is equal to $\pm t^k$ for some $k \in \mathbb{Z}$. Thanks to Lemmas 3.1, 3.3 and 3.5, it follows that the $\hat{\Gamma}_i$ can be grouped by pairs with opposite corresponding brackets. Hence $[(M, K); \Gamma] = [(M, K); \Gamma'] \text{ mod } \mathcal{F}_{n+1}^b(M, K)$. \square

Lemma 3.9. *Let Γ be an n -component Y -link null in $M \setminus K$. Let ℓ be a leaf of Γ . Let $\widetilde{\Gamma \setminus \ell}$ be the preimage of $\Gamma \setminus \ell$ in the infinite cyclic covering \tilde{X} associated with (M, K) . Let $\tilde{\ell}$ be a lift of ℓ . If $\tilde{\ell}$ is trivial in $H_1(\tilde{X} \setminus (\widetilde{\Gamma \setminus \ell}); \mathbb{Q})$, then $[(M, K); \Gamma] = 0 \text{ mod } \mathcal{F}_{n+1}^b(M, K)$.*

Proof. Since $\tilde{\ell}$ has a multiple which is trivial in $H_1(\tilde{X}; \mathbb{Z})$, Lemma 3.5 allows us to assume $\tilde{\ell}$ itself is trivial in $H_1(\tilde{X}; \mathbb{Z})$. Hence $\tilde{\ell}$ is a product of commutators of loops in \tilde{X} . It follows that ℓ is homotopic to $\prod_{i \in I} [\alpha_i, \beta_i]$ in $M \setminus K$, where I is a finite set and the α_i and β_i satisfy $lk(\alpha_i, K) = 0$, $lk(\beta_i, K) = 0$. Construct a surface Σ in $(M \setminus K) \setminus \Gamma$ whose handles are bands around the α_i and β_i , so that $\partial\Sigma$ is homotopic to ℓ in $M \setminus K$. Let Γ' be the Y -link obtained from Γ by replacing ℓ by $\partial\Sigma$. Note that the lifts of $\partial\Sigma$ are null-homologous in $\tilde{X} \setminus (\widetilde{\Gamma \setminus \ell})$. Hence, by Lemma 3.8, $[(M, K); \Gamma] = [(M, K); \Gamma'] \text{ mod } \mathcal{F}_{n+1}^b(M, K)$.

Let us prove that $[(M, K); \Gamma'] = 0 \text{ mod } \mathcal{F}_{n+1}^b(M, K)$. Apply Lemma 3.5 to cut the leaf $\partial\Sigma$ into leaves α_i , β_i , α_i^{-1} , β_i^{-1} . Apply it again to reglue each leaf α_i with the corresponding leaf α_i^{-1} and each leaf β_i with the corresponding leaf β_i^{-1} . The obtained Y -links all have a leaf which is homotopically trivial in the complement of K and of the complement of the leaf in the Y -link. Then the result follows from Lemma 3.7. \square

Lemma 3.10. *Let Γ be an n -component Y -link null in $M \setminus K$. Let ℓ be a leaf of Γ . Let $\widetilde{\Gamma \setminus \ell}$ be the preimage of $\Gamma \setminus \ell$ in the infinite cyclic covering \tilde{X} associated with (M, K) . Let $\tilde{\ell}$ be a lift of ℓ . Fix $\Gamma \setminus \ell$. Then the class of $[(M, K); \Gamma] \text{ mod } \mathcal{F}_{n+1}^b(M, K)$ only depends on the class of $\tilde{\ell}$ in $H_1(\tilde{X} \setminus (\widetilde{\Gamma \setminus \ell}); \mathbb{Q})$ and this dependance is \mathbb{Q} -linear.*

Proof. Let Γ' be a null Y -link which has a leaf ℓ' such that $\Gamma' \setminus \ell'$ coincides with $\Gamma \setminus \ell$, and $\tilde{\ell}'$ is homologous to $\tilde{\ell}$ in $\tilde{X} \setminus (\widetilde{\Gamma \setminus \ell})$, where $\tilde{\ell}'$ is the lift of ℓ' which has the same basepoint as $\tilde{\ell}$. Construct another null Y -link Γ^δ by replacing the leaf ℓ by $\ell - \ell'$ in Γ (see Figure 13). By Lemma 3.9, $[(M, K); \Gamma^\delta] = 0 \text{ mod } \mathcal{F}_{n+1}^b(M, K)$. Thus Lemma 3.5 implies $[(M, K); \Gamma] = [(M, K); \Gamma'] \text{ mod } \mathcal{F}_{n+1}^b(M, K)$. Linearity follows from Lemma 3.5. \square

4 Colored diagrams and Y -links

In this section, we apply clasper calculus to obtain the maps from diagrams spaces to graded quotients of Proposition 2.6 and Theorem 2.17.




Figure 13: The leaf $\ell - \ell'$

Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. An $(\mathfrak{A}, \mathfrak{b})$ -colored diagram is an *elementary* ($(\mathfrak{A}, \mathfrak{b})$ -colored) *diagram* if its edges that connect two trivalent vertices are colored by powers of t and if its edges adjacent to univalent vertices are colored by 1. Below, we associate a null Y-link with some elementary diagrams that generate $\tilde{\mathcal{A}}_n(\mathfrak{A}, \mathfrak{b})$. Let $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $\xi : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M, K)$ be an isomorphism. Let $m(K)$ be a meridian of K .

Let D be an elementary diagram. An embedding of D in $M \setminus K$ is *admissible* if the following conditions are satisfied.

- The vertices of D are embedded in some ball $B \subset M \setminus K$.
- Consider an edge colored by t^k . The homology class in $H_1(M \setminus K; \mathbb{Z})$ of the closed curve obtained by connecting the extremities of this edge by a path in B is $k m(K)$.

Such an embedding always exists. It suffices to embed the diagram in B , and to let each edge colored by t^k turn around K , k times. With an admissible embedding of an elementary diagram, we wish to associate a null Y-link.

Let Γ be a Y-graph null in $M \setminus K$. Let p be the internal vertex of Γ . Let ℓ be a leaf of Γ . The curve $\hat{\ell}$ drawn in Figure 14 is the *extension of ℓ in Γ* .




Figure 14: Extension of a leaf in a Y-graph

Let D be an elementary diagram, equipped with an admissible embedding in $M \setminus K$. Equip D with the framing induced by an immersion in the plane which induces the fixed orientation of the trivalent vertices. If an edge connects two trivalent vertices, then insert a little Hopf link in this edge, as shown in Figure 15. At each univalent vertex v , glue a



Figure 15: Replacement of an edge

leaf ℓ_v , trivial in $H_1(M \setminus K; \mathbb{Q})$, in order to obtain a null Y-link Γ . Let V be the set of all univalent vertices of D . Let B be the ball in the definition of the admissible embedding of D . Let \tilde{B} be a lift of B in the infinite cyclic covering \tilde{X} of the exterior of K in M . For $v \in V$, let γ_v be the coloring of v , let $\hat{\ell}_v$ be the extension of ℓ_v in Γ and let $\tilde{\ell}_v$ be the lift of $\hat{\ell}_v$ in \tilde{X} defined by lifting the basepoint in \tilde{B} . The null Y-link Γ is a *realization of D in (M, K) with respect to ξ* if the following conditions are satisfied:

- for all $v \in V$, $\tilde{\ell}_v$ is homologous to $\xi(\gamma_v)$,
- for all $(v, v') \in V^2$, $lk_e(\tilde{\ell}_v, \tilde{\ell}_{v'}) = f_{vv'}$.

If such a realization exists, the elementary diagram D is ξ -realizable.

Lemma 4.1. *Let $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $\xi : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M, K)$ be an isomorphism. Let $D \in \tilde{\mathcal{A}}_n(\mathfrak{A}, \mathfrak{b})$ be an elementary diagram of degree $n > 0$, ξ -realizable. Let Γ be a realization of D in (M, K) with respect to ξ . Then the class of $[(M, K); \Gamma]$ mod $\mathcal{F}_{n+1}^b(M, K)$ does not depend on the realization of D .*

Proof. If the ball B and its lift \tilde{B} are fixed, then the result follows from Lemmas 3.3 and 3.10. Fix the ball B and consider another lift $\tilde{B}' = \tau^k(\tilde{B})$ of B , where τ is the automorphism of \tilde{X} which induces the action of t and $k \in \mathbb{Z}$. A realization of D with respect to \tilde{B}' can be obtained from Γ by letting the internal vertex of each Y-graph in Γ turn around K , k times, and come back into B , by an isotopy of (M, K, Γ) . This does not change the result of the surgeries on these Y-graphs, hence this does not modify the bracket $[(M, K); \Gamma]$. Now consider two balls B_1 and B_2 in $M \setminus K$. If $B_1 \subset B_2$, a realization of D with respect to B_1 is a realization of D with respect to B_2 . If $B_1 \cap B_2 \neq \emptyset$, there is a ball $B_3 \subset B_1 \cap B_2$. If $B_1 \cap B_2 = \emptyset$, there is a ball $B_3 \supset B_1 \cup B_2$. Hence the class of the bracket $[(M, K); \Gamma]$ does not depend on the chosen ball B . \square

In the sequel, if D is a ξ -realizable elementary diagram, $[(M, K); D]_\xi$ denotes the class of $[(M, K); \Gamma]$ modulo $\mathcal{F}_{n+1}^b(M, K)$.

Let D be any elementary diagram. Let V be the set of all univalent vertices of D . For any family of rational numbers $(q_v)_{v \in V}$, define an elementary diagram $D' = (q_v)_{v \in V} \cdot D$ from D in the following way. Keep the same graph and the same colorings of the edges. For $v \in V$, multiply the coloring of v by q_v . For $v \neq v' \in V$, set $f_{vv'}^{D'} = q_v q_{v'} f_{vv'}^D$.

Lemma 4.2. *Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. Let $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $\xi : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M, K)$ be an isomorphism. Let D be any elementary diagram. Let V be the set of all univalent vertices of D . Then there exists a family of positive integers $(s_v)_{v \in V}$ such that $(s_v)_{v \in V} \cdot D$ is ξ -realizable.*

Proof. Let \tilde{X} be the infinite cyclic covering associated with (M, K) . Since any homology class in \mathfrak{A} has a multiple which can be represented by a knot in \tilde{X} , we can assume that the color γ_v of each vertex v in V can be represented by a knot in \tilde{X} . From D , define as above a Y-link Γ , null in $M \setminus K$, with leaves ℓ_v which satisfy the condition that $\tilde{\ell}_v$ is homologous to $\xi(\gamma_v)$ for all $v \in V$. For $v \neq v' \in V$, set $P_{vv'} = lk_e(\tilde{\ell}_v, \tilde{\ell}_{v'}) - f_{vv'}$. We can assume that $P_{vv'} \in \mathbb{Z}[t^{\pm 1}]$ for all $v \neq v' \in V$. Add well chosen meridians of ℓ_v to $\ell_{v'}$ to get $P_{vv'} = 0$. \square

Lemma 4.3. *Let $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $\xi : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M, K)$ be an isomorphism. Let D be an elementary $(\mathfrak{A}, \mathfrak{b})$ -colored diagram. Let V be the set of all univalent vertices of D . Let $(s_v)_{v \in V}$ and $(s'_v)_{v \in V}$ be families of integers such that $(s_v)_{v \in V} \cdot D$ and $(s'_v)_{v \in V} \cdot D$ are ξ -realizable. Then:*

$$\prod_{v \in V} s'_v [(M, K); (s_v)_{v \in V} \cdot D]_{\xi} = \prod_{v \in V} s_v [(M, K); (s'_v)_{v \in V} \cdot D]_{\xi}.$$

Proof. Let Γ be a realization of $(s_v)_{v \in V} \cdot (s'_v)_{v \in V} \cdot D$ in (M, K) with respect to ξ . By Lemma 3.10, $[(M, K); \Gamma]$ is equal to both sides of the equality. \square

Let D be an elementary $(\mathfrak{A}, \mathfrak{b})$ -colored diagram. Let V be the set of all univalent vertices of D . The above result allows us to define:

$$[(M, K); D]_{\xi} = \prod_{v \in V} \frac{1}{s_v} [(M, K); (s_v)_{v \in V} \cdot D]_{\xi},$$

where $(s_v)_{v \in V}$ is any family of integers such that $(s_v)_{v \in V} \cdot D$ is ξ -realizable.

Lemma 4.4. *Let D be an elementary $(\mathfrak{A}, \mathfrak{b})$ -colored diagram. Let (M, K) and (M', K') be $\mathbb{Q}SK$ -pairs in $\mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $\xi : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M, K)$ and $\xi' : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M', K')$ be isomorphisms. Then $[(M', K'); D]_{\xi'} = [(M, K); D]_{\xi} \text{ mod } \mathcal{F}_{n+1}(\mathfrak{A}, \mathfrak{b})$.*

Proof. Set $\zeta = \xi' \circ \xi^{-1}$. By Theorem 2.2, (M', K') can be obtained from (M, K) by a finite sequence of null LP-surgeries which induces $\zeta \circ m_k$, for $k \in \mathbb{Z}$, where m_k is the multiplication by t^k . Assume the sequence contains a single surgery $(\frac{A'}{A})$. Let V be the set of all univalent vertices of D . Let $(s_v)_{v \in V}$ be a family of integers such that $(s_v)_{v \in V} \cdot D$ is ξ -realizable by a null Y-link Γ in $(M \setminus K) \setminus A$. Then:

$$\left[(M, K); \Gamma, \frac{A'}{A} \right] = [(M, K); \Gamma] - [(M', K'); \Gamma].$$

In (M', K') , Γ is a realization of $(s_v)_{v \in V} \cdot D$ with respect to $\xi' \circ m_k$. Hence it is also a realization of $(s_v)_{v \in V} \cdot D$ with respect to ξ' (it suffices to change the lift \tilde{B} of the ball B , see Lemma 4.1).

The case of several surgeries easily follows. \square

In the sequel, the class of $[(M, K); D]_\xi$ modulo $\mathcal{F}_{n+1}(\mathfrak{A}, \mathfrak{b})$ is denoted by $[D]$.

Proposition 4.5. *Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. Let $n > 0$. There is a canonical, \mathbb{Q} -linear and surjective map $\varphi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n^b(\mathfrak{A}, \mathfrak{b})$ given by $D \mapsto [D]$ for any elementary diagram D .*

Proof. Let \mathcal{D}_n be the rational vector space freely generated by the $(\mathfrak{A}, \mathfrak{b})$ -colored diagrams of degree n . If D is an elementary $(\mathfrak{A}, \mathfrak{b})$ -colored diagram, set $\tilde{\varphi}_n(D) = [D]$. Define $\tilde{\varphi}_n(D)$ for any $(\mathfrak{A}, \mathfrak{b})$ -colored diagram D so that the obtained \mathbb{Q} -linear map $\tilde{\varphi}_n : \mathcal{D}_n \rightarrow \mathcal{G}_n^b(\mathfrak{A}, \mathfrak{b})$ satisfies the relation LE and EV. Let us check that $\tilde{\varphi}_n$ satisfies the relations AS, IHX, OR, Hol, LV, LD and Aut. OR is trivial. LV follows from Lemma 3.10. Hol is obtained by letting the corresponding vertex of a realization of D turn around the knot K . AS and IHX respectively follow from [GGP01, Corollary 4.6] and [GGP01, Lemma 4.10]. Aut follows from Lemma 4.4. For the relation LD, it suffices to prove that $\tilde{\varphi}_n(D) = \tilde{\varphi}_n(D') + \tilde{\varphi}_n(D_0)$, where D , D' and D_0 are elementary diagrams identical, except for the part drawn in Figure 16. Note that the edges adjacent to v_1 and v_2 are colored by 1. Since the diagram D_0 and the diagram D'_0 drawn in Figure 17 can be realized by the same null Y-link, we have $\tilde{\varphi}_n(D_0) = \tilde{\varphi}_n(D'_0)$. To see that $\tilde{\varphi}_n(D'_0) = \tilde{\varphi}_n(D''_0)$, apply the relation LV at the vertex v_2 to obtain $\tilde{\varphi}_n(D''_0) = \tilde{\varphi}_n(D'_0) + \tilde{\varphi}_n(D_{00})$ and then apply the relation LV at the vertex v_1 to obtain $\tilde{\varphi}_n(D_{00}) = 0$. Apply again the relation LV at the vertex v_1 to get $\tilde{\varphi}_n(D) = \tilde{\varphi}_n(D') + \tilde{\varphi}_n(D''_0)$.




Figure 16: The diagrams D , D' , and D_0 , where $\gamma_1, \gamma_2 \in \mathfrak{A}$ and $k \in \mathbb{Z}$.

Finally, the map $\tilde{\varphi}_n$ induces a canonical \mathbb{Q} -linear map $\varphi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n^b(\mathfrak{A}, \mathfrak{b})$. For $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$, any n -component Y-link null in $M \setminus K$ is a realization of an elementary $(\mathfrak{A}, \mathfrak{b})$ -colored diagram, which is the disjoint union of n diagrams of degree 1. Hence φ_n is surjective. \square

In the setting of \mathbb{Z} SK-pairs, all the results of Section 3 apply since we work modulo $\mathcal{F}_{n+1}^b(M, K)$. All the results of the current section apply as well. In Lemma 4.4, note




Figure 17: The diagrams D'_0 , D''_0 and D_{00} , with $v \neq v_1, v_2$.

that we use Theorem 2.16 instead of Theorem 2.2. We finally get the similar result to the above proposition.

Proposition 4.6. *Fix an integral Blanchfield module $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$. Let $n > 0$. There is a canonical, \mathbb{Q} -linear and surjective map $\varphi_n^{\mathbb{Z}} : \mathcal{A}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) \rightarrow \mathcal{G}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ given by $D \mapsto [(M, K); D]$ for any elementary diagram D , where (M, K) is any \mathbb{Z} SK-pair with Blanchfield module $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$, $\xi : (\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) \rightarrow (\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})(M, K)$ is an isomorphism and Γ_D is a realization of D in (M, K) with respect to ξ .*

Fix $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ and set $(\mathfrak{A}, \mathfrak{b}) = (\mathbb{Q} \otimes_{\mathbb{Z}} \mathfrak{A}_{\mathbb{Z}}, Id_{\mathbb{Q}} \otimes \mathfrak{b}_{\mathbb{Z}})$. The corresponding map φ_n satisfies $\varphi_n \circ p_n = \omega_n \circ \varphi_n^{\mathbb{Z}}$, where $p_n : \mathcal{A}_n^{\mathbb{Z}}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\mathfrak{A}, \mathfrak{b})$ is the natural projection and $\omega_n : \mathcal{G}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) \rightarrow \mathcal{G}_n^b(\mathfrak{A}, \mathfrak{b})$ is the map induced by the inclusion $\mathcal{F}_n^{\mathbb{Z}}(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}}) \hookrightarrow \mathcal{F}_n(\mathfrak{A}, \mathfrak{b})$.

5 The surjective map $\varphi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$

In this section, we prove Theorems 2.7 and 2.3.

Fix a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$. Let (M, K) be a \mathbb{Q} SK-pair in $\mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $\xi : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M, K)$ be an isomorphism. Let $n > 0$. Let D be an $(\mathfrak{A}, \mathfrak{b})$ -augmented diagram of degree n whose Jacobi part D_J is elementary. With an isolated vertex colored by a prime integer p , we associate a surgery $\frac{B_p}{B^3}$, where B_p is a fixed \mathbb{Q} -ball such that $|H_1(B_p; \mathbb{Z})| = p$. Hence, if D_J is ξ -realizable, with a realization of D_J , we associate a family of n disjoint null LP-surgeries.

Lemma 5.1. *Let $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $\xi : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M, K)$ be an isomorphism. Let $n > 0$. Let D be an $(\mathfrak{A}, \mathfrak{b})$ -augmented diagram whose Jacobi part D_J is elementary. Let $(p_i)_{1 \leq i \leq n-k}$ be the labels of the isolated vertices of D . If D_J is ξ -realizable, let Γ be a realization of D_J in (M, K) with respect to ξ . Then:*

- $[(M, K); D]_{\xi} := \left[(M, K); \left(\frac{B_{p_i}}{B^3} \right)_{1 \leq i \leq n-k}, \Gamma \right] \in \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ does not depend on (M, K) , on ξ or on the realization Γ of D_J .

If D_J is any elementary diagram, set $[D] = \prod_{v \in V} \frac{1}{s_v} [(M, K); (s_v)_{v \in V} \cdot D]_\xi$, where $(s_v)_{v \in V}$ is a family of integers such that $(s_v)_{v \in V} \cdot D_J$ is ξ -realizable and $(s_v)_{v \in V} \cdot D$ is the disjoint union of $(s_v)_{v \in V} \cdot D_J$ with the 0-valent part of D . Then:

- $[D] \in \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ does not depend on $(s_v)_{v \in V}$, (M, K) or ξ .

Proof. Take $(M', K') \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$ and an isomorphism $\xi' : (\mathfrak{A}, \mathfrak{b}) \rightarrow (\mathfrak{A}, \mathfrak{b})(M, K)$ such that D_J is ξ' -realizable. Let Γ' be a realization of D_J with respect to ξ' . By Proposition 4.5:

$$[(M', K'); \Gamma'] = [(M, K); \Gamma] \text{ mod } \mathcal{F}_{k+1}(\mathfrak{A}, \mathfrak{b}).$$

Let p be a prime integer. Let $M_p = B^3 \cup_{\partial B^3} B_p$. In the equality in $\mathcal{F}_0(\mathfrak{A}, \mathfrak{b})$ corresponding to the above relation, make a connected sum of each QSK-pair with M_p . Then subtract the new equality from the original one, to obtain:

$$\left[(M', K'); \frac{B_p}{B^3}, \Gamma' \right] = \left[(M, K); \frac{B_p}{B^3}, \Gamma \right] \text{ mod } \mathcal{F}_{k+2}(\mathfrak{A}, \mathfrak{b}).$$

Applying this process $n - k$ times, we get:

$$\left[(M', K'); \left(\frac{B_{p_i}}{B^3} \right)_{1 \leq i \leq n-k}, \Gamma' \right] = \left[(M, K); \left(\frac{B_{p_i}}{B^3} \right)_{1 \leq i \leq n-k}, \Gamma \right] \text{ mod } \mathcal{F}_{n+1}(\mathfrak{A}, \mathfrak{b}).$$

If D_J is any elementary diagram, use Lemma 3.10, as in Lemma 4.3, to prove that $[(M, K); D]_\xi = \prod_{v \in V} \frac{1}{s_v} [(M, K); (s_v)_{v \in V} \cdot D]_\xi$ does not depend on the family of integers $(s_v)_{v \in V}$ such that $(s_v)_{v \in V} \cdot D_J$ is ξ -realizable. Conclude with the first assertion of the lemma. \square

The above result implies that the map $\varphi_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n^b(\mathfrak{A}, \mathfrak{b})$ extends to a canonical \mathbb{Q} -linear map $\varphi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ defined by $\varphi_n(D) = [D]$ for any diagram $D \in \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b})$ whose Jacobi part is elementary. To prove Theorem 2.7, it remains to show that the map $\varphi_n : \mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{G}_n(\mathfrak{A}, \mathfrak{b})$ is surjective. We first recall results from [Mou12a] and give consequences of them.

Definition 5.2. Let d be a positive integer. A d -torus is a \mathbb{Q} -torus T_d such that:

- $H_1(\partial T_d; \mathbb{Z}) = \mathbb{Z}\alpha \oplus \mathbb{Z}\beta$, with $\langle \alpha, \beta \rangle = 1$,
- $d\alpha = 0$ in $H_1(T_d; \mathbb{Z})$,
- $\beta = d\gamma$ in $H_1(T_d; \mathbb{Z})$, where γ is a curve in T_d ,
- $H_1(T_d; \mathbb{Z}) = \frac{\mathbb{Z}}{d\mathbb{Z}}\alpha \oplus \mathbb{Z}\gamma$.

Definition 5.3. An elementary surgery is an LP-surgery among the following ones:

1. connected sum (genus 0),
2. LP-replacement of a standard torus by a d -torus (genus 1),
3. Borromean surgery (genus 3).

Theorem 5.4 ([Mou12a] Theorem 1.15). *If A and B are two \mathbb{Q} -handlebodies with LP-identified boundaries, then B can be obtained from A by a finite sequence of elementary surgeries and their inverses in the interior of the \mathbb{Q} -handlebodies.*

Corollary 5.5. *The space $\mathcal{F}_n(\mathfrak{A}, \mathfrak{b})$ is generated by the $\left[(M, K); \left(\frac{E'_i}{E_i} \right)_{1 \leq i \leq n} \right]$ defined by a $\mathbb{Q}SK$ -pair $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$ and elementary null LP-surgeries $\left(\frac{E'_i}{E_i} \right)$.*

Proof. Consider $\left[(M, K); \left(\frac{A'_i}{A_i} \right)_{1 \leq i \leq n} \right] \in \mathcal{F}_n(\mathfrak{A}, \mathfrak{b})$. By Theorem 5.4, for each i , A_i and A'_i can be obtained from one another by a finite sequence of elementary surgeries or their inverses. Write $A'_1 = A_1 \left(\frac{E'_1}{E_1} \right) \dots \left(\frac{E'_k}{E_k} \right)$. For $0 \leq j \leq k$, set $B_j = A_1 \left(\frac{E'_1}{E_1} \right) \dots \left(\frac{E'_j}{E_j} \right)$. Then:

$$\left[(M, K); \left(\frac{A'_i}{A_i} \right)_{1 \leq i \leq n} \right] = \sum_{j=1}^k \left[(M, K) \left(\frac{B_{j-1}}{B_0} \right); \frac{E'_j}{E_j}, \left(\frac{A'_i}{A_i} \right)_{2 \leq i \leq n} \right].$$

Decompose each surgery $\left(\frac{A'_i}{A_i} \right)$ in this way and conclude with:

$$\left[(M, K); \frac{E'}{E}, \left(\frac{A'_i}{A_i} \right)_{2 \leq i \leq n} \right] = - \left[(M, K) \left(\frac{E'}{E} \right); \frac{E}{E'}, \left(\frac{A'_i}{A_i} \right)_{2 \leq i \leq n} \right].$$

□

Let $\mathcal{F}_0^{\text{QS}}$ be the rational vector space generated by all \mathbb{Q} -spheres up to orientation-preserving homeomorphism. Let $(\mathcal{F}_n^{\text{QS}})_{n \in \mathbb{N}}$ be the filtration of $\mathcal{F}_0^{\text{QS}}$ defined by LP-surgeries, as before Definition 2.1. Let $\mathcal{G}_n^{\text{QS}} = \frac{\mathcal{F}_n^{\text{QS}}}{\mathcal{F}_{n+1}^{\text{QS}}}$ be the associated quotients.

Lemma 5.6 ([Mou12a] Proposition 1.8). *For each prime integer p , let B_p be a \mathbb{Q} -ball such that $H_1(B_p; \mathbb{Z}) \cong \frac{\mathbb{Z}}{p\mathbb{Z}}$. Then $\mathcal{G}_1^{\text{QS}} = \bigoplus_{p \text{ prime}} \mathbb{Q} \left[S^3; \frac{B_p}{B^3} \right]$.*

Lemma 5.7. For each prime integer p , let B_p be a \mathbb{Q} -ball such that $H_1(B_p; \mathbb{Z}) \cong \frac{\mathbb{Z}}{p\mathbb{Z}}$. Let (M, K) be a $\mathbb{Q}SK$ -pair in $\mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let B be a \mathbb{Q} -ball. Let $\left(\frac{A'_i}{A_i}\right)_{1 \leq i < n}$ be disjoint null LP-surgeries in (M, K) . Then $\left[(M, K); \frac{B}{B^3}, \left(\frac{A'_i}{A_i}\right)_{1 \leq i < n}\right]$ is a rational linear combination of the $\left[(M, K); \frac{B_p}{B^3}, \left(\frac{A'_i}{A_i}\right)_{1 \leq i < n}\right]$ and elements of $\mathcal{F}_{n+1}(\mathfrak{A}, \mathfrak{b})$.

Proof. By Lemma 5.6, there is a relation:

$$\left[S^3; \frac{B}{B^3}\right] = \sum_{p \text{ prime}} a_p \left[S^3; \frac{B_p}{B^3}\right] + \sum_{j \in J} b_j \left[N_j; \frac{C'_j}{C_j}, \frac{D'_j}{D_j}\right],$$

where J is a finite set, the a_p and b_j are rational numbers, the a_p are all trivial except a finite number and, for $j \in J$, $\left[N_j; \frac{C'_j}{C_j}, \frac{D'_j}{D_j}\right] \in \mathcal{F}_2^{\text{QS}}$. For $I \subset \{1, \dots, n-1\}$, make the connected sum of each \mathbb{Q} -sphere in the relation with $M \left(\left(\frac{A'_i}{A_i}\right)_{i \in I}\right)$ to obtain:

$$\begin{aligned} & \left[(M, K) \left(\left(\frac{A'_i}{A_i}\right)_{i \in I}\right); \frac{B}{B^3}\right] = \\ & \sum_{p \text{ prime}} a_p \left[(M, K) \left(\left(\frac{A'_i}{A_i}\right)_{i \in I}\right); \frac{B_p}{B^3}\right] + \sum_{j \in J} b_j \left[(M \# N_j, K) \left(\frac{A'_i}{A_i}\right)_{i \in I}; \frac{C'_j}{C_j}, \frac{D'_j}{D_j}\right]. \end{aligned}$$

Summing these equalities for all $I \subset \{1, \dots, n-1\}$, with appropriate signs, we get:

$$\begin{aligned} & \left[(M, K); \frac{B}{B^3}, \left(\frac{A'_i}{A_i}\right)_{1 \leq i < n}\right] = \\ & \sum_{p \text{ prime}} a_p \left[(M, K); \frac{B_p}{B^3}, \left(\frac{A'_i}{A_i}\right)_{1 \leq i < n}\right] + \sum_{j \in J} b_j \left[(M \# N_j, K); \frac{C'_j}{C_j}, \frac{D'_j}{D_j}, \left(\frac{A'_i}{A_i}\right)_{1 \leq i < n}\right]. \end{aligned}$$

□

Corollary 5.8. Let $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$. Let $\left(\frac{E'_i}{E_i}\right)_{1 \leq i \leq n}$ be null elementary surgeries of genus 0 or 3. Then $\left[(M, K); \left(\frac{E'_i}{E_i}\right)_{1 \leq i \leq n}\right] \in \varphi_n(\mathcal{A}_n^{\text{aug}}(\mathfrak{A}, \mathfrak{b}))$.

Proof. Thanks to Lemma 5.7, it suffices to treat the case when the genus 0 surgeries are surgeries of type $\left(\frac{B}{B^3}\right)$ for a \mathbb{Q} -ball B such that $|H_1(B; \mathbb{Z})|$ is a prime integer. In this case, the considered bracket is the image of a diagram given as the disjoint union of 0-valent vertices and of $(\mathfrak{A}, \mathfrak{b})$ -colored diagrams of degree 1. □

To conclude the proof of Theorem 2.7, we need the next result about degree 1 invariants of framed \mathbb{Q} -tori, i.e. \mathbb{Q} -tori equipped with an oriented longitude. Note that any two framed \mathbb{Q} -tori have a canonical LP-identification of their boundaries, which identifies the fixed longitudes. LP-surgeries are well-defined on framed \mathbb{Q} -tori and we have an associated notion of finite type invariants.

Lemma 5.9 ([Mou12a] Corollary 5.10). *For any prime integer p , let M_p be a \mathbb{Q} -sphere such that $H_1(M_p; \mathbb{Z}) \cong \mathbb{Z}/p\mathbb{Z}$. Let T_0 be a framed standard torus. If μ is a degree 1 invariant of framed \mathbb{Q} -tori such that $\mu(T_0) = 0$ and $\mu(T_0 \sharp M_p) = 0$ for any prime integer p , then $\mu = 0$.*

Proof of Theorem 2.7. Take $\lambda \in (\mathcal{F}_n(\mathfrak{A}, \mathfrak{b}))^*$ such that $\lambda(\mathcal{F}_{n+1}(\mathfrak{A}, \mathfrak{b})) = 0$. Assume $\lambda(\varphi_n(\mathcal{A}_n^{aug}(\mathfrak{A}, \mathfrak{b}))) = 0$. In order to prove that φ_n is onto, it is enough to prove that $\lambda = 0$. Thanks to Corollary 5.5, it suffices to prove that λ vanishes on the brackets defined by elementary surgeries. For elementary surgeries of genus 0 and 3, this follows from Corollary 5.8.

Consider a bracket $\left[(M, K); \left(\frac{T_{d_i}}{T_i} \right)_{1 \leq i \leq k}, \left(\frac{E'_i}{E_i} \right)_{1 \leq i \leq n-k} \right]$, where $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$, the T_i are standard tori null in $M \setminus K$, the T_{d_i} are d_i -tori for some positive integers d_i and the $\left(\frac{E'_i}{E_i} \right)$ are null elementary surgeries of genus 0 or 3. By induction on k , we will prove that λ vanishes on this bracket. We have treated the case $k = 0$. Assume $k > 0$. Fix a parallel of T_1 . If T is a framed \mathbb{Q} -torus, set:

$$\bar{\lambda}(T) = \lambda \left(\left[(M, K); \frac{T}{T_1}, \left(\frac{T_{d_i}}{T_i} \right)_{2 \leq i \leq k}, \left(\frac{E'_i}{E_i} \right)_{1 \leq i \leq n-k} \right] \right),$$

where the LP-identification $\partial T \cong \partial T_1$ identifies the preferred parallels. Then $\bar{\lambda}$ is a degree 1 invariant of framed \mathbb{Q} -tori:

$$\bar{\lambda} \left(\left[T; \frac{B_1}{A_1}, \frac{B_2}{A_2} \right] \right) = -\lambda \left(\left[(M, K) \left(\frac{T}{T_1} \right); \frac{B_1}{A_1}, \frac{B_2}{A_2}, \left(\frac{T_{d_i}}{T_i} \right)_{2 \leq i \leq k}, \left(\frac{E'_i}{E_i} \right)_{1 \leq i \leq n-k} \right] \right) = 0.$$

Moreover, we have $\bar{\lambda}(T_1) = 0$ and, by induction, $\bar{\lambda} \left(T_1 \left(\frac{B_p}{B^3} \right) \right) = 0$. Thus, by Lemma 5.9, $\bar{\lambda} = 0$. \square

Proof of Theorem 2.3. Theorem 2.7 provides a surjective map $\varphi_1: \mathcal{A}_1^{aug}(\mathfrak{A}, \mathfrak{b}) \twoheadrightarrow \mathcal{G}_1(\mathfrak{A}, \mathfrak{b})$. Thanks to Lemma 2.5, we have $\mathcal{A}_1^{aug}(\mathfrak{A}, \mathfrak{b}) = \bigoplus_{p \text{ prime}} \mathbb{Q} \bullet_p$. Hence $\mathcal{G}_1(\mathfrak{A}, \mathfrak{b})$ is generated by the images of the diagrams \bullet_p , which are the brackets $\left[(M, K); \frac{B_p}{B^3} \right]$ for all prime integers p , with any $(M, K) \in \mathcal{P}(\mathfrak{A}, \mathfrak{b})$.

For any prime integer p , define a \mathbb{Q} -linear map $\nu_p: \mathcal{F}_0 \rightarrow \mathbb{Q}$ by setting $\nu_p(M, K) = v_p(|H_1(M; \mathbb{Z})|)$ for all QSK-pair (M, K) , where v_p denotes the p -adic valuation. By

[Mou12a, Proposition 1.9], the ν_p are degree 1 invariants of \mathbb{Q} -spheres, hence they are also degree 1 invariants of QSK-pairs. This implies that the family $\left(\left[(M, K); \frac{B_p}{B_3} \right] \right)_{p \text{ prime}}$ is free in $\mathcal{G}_1(\mathfrak{A}, \mathfrak{b})$. \square

6 Extension of the Lescop/Kricker invariant

In this section, we prove Theorem 2.10.

Given two invariants λ_1 and λ_2 of QSK-pairs, define their product on any QSK-pair (M, K) by $(\lambda_1 \lambda_2)(M, K) = \lambda_1(M, K) \lambda_2(M, K)$ and extend to \mathcal{F}_0 by linearity. The following lemma is classical and holds for any objects and any invariants with values in some ring (see for instance [Mou12a, Lemma 6.2]).

Lemma 6.1.

$$\begin{aligned} & \left(\prod_{j=1}^n \lambda_j \right) \left(\left[(M, K); \left(\frac{B_i}{A_i} \right)_{i \in I} \right] \right) \\ &= \sum_{\emptyset = J_0 \subset \dots \subset J_n = I} \prod_{j=1}^n \lambda_j \left(\left[(M, K) \left(\left(\frac{B_i}{A_i} \right)_{i \in J_{j-1}} \right); \left(\frac{B_i}{A_i} \right)_{i \in J_j \setminus J_{j-1}} \right] \right) \end{aligned}$$

This lemma implies in particular that a product of finite type invariants is a finite type invariant whose degree is at most the sum of the degrees of the factors.

Proof of Theorem 2.10. We begin with a preliminary remark about the invariant Z . It follows from the last point in Theorem 2.9 that $Z_n \circ \varphi_n$ vanishes on diagrams that contain isolated vertices. Now, the degree n part of Z^{aug} is given by:

$$Z_n^{aug} = \sum_{k=0}^n \sum_{\substack{p_1 < \dots < p_s \\ \text{prime integers}}} \sum_{\substack{t_1 + \dots + t_s = n - k \\ t_i > 0}} Z_k \sqcup \left(\prod_{i=1}^s \frac{1}{t_i!} (\rho_{p_i})^{t_i} \right).$$

That Z_n^{aug} vanishes on \mathcal{F}_{n+1} follows from Lemma 6.1.

Let us compute $Z_n^{aug} \circ \varphi_n$. Let D be an $(\mathfrak{A}, \mathfrak{b})$ -augmented diagram of degree n . Write D as the disjoint union of its Jacobi part D_J and its 0-valent part D_\bullet . Apply Lemma 6.1, noting that for a term in the right hand side of the obtained equality to be non trivial:

- each bracket must have exactly the order of the corresponding invariant,
- each invariant ρ_p must be evaluated on a bracket associated with the diagram \bullet_p ,
- the invariant Z_k must be evaluated on a bracket associated with a diagram without isolated vertices.

It follows that $Z_n^{aug} \circ \varphi_n(D) = (Z_k \circ \varphi_k(D_J)) \sqcup D_\bullet = \psi_k(D_J) \sqcup D_\bullet = \psi_n(D)$. \square

Proof. Apply the Aut relation with the automorphism of $(\mathfrak{A}, \mathfrak{b})$ given by the opposite of the identity on \mathfrak{A}_1 and the identity on $\mathfrak{A}_2 \oplus \mathfrak{A}''$. \square

Corollary 7.4. *Assume \mathfrak{A}' is the orthogonal direct sum of submodules \mathfrak{A}_i , $i = 1, \dots, k$. Let $\gamma, \eta \in \mathfrak{A}'$. Write $\gamma = \sum_{i=1}^k \gamma_i$ and $\eta = \sum_{i=1}^k \eta_i$ with $\gamma_i, \eta_i \in \mathfrak{A}_i$. Then:*

$$\gamma \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f} \bullet \eta = \sum_{i=1}^k \gamma_i \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f_i} \bullet \eta_i,$$

for all families of rational fractions f_i such that $\mathfrak{b}(\gamma_i, \eta_i) = f_i \bmod \mathbb{Q}[t^{\pm 1}]$ and $\sum_{i=1}^k f_i = f$.

Lemma 7.5. *If $\gamma, \eta \in \mathfrak{A}'$ and $P \in \mathbb{Q}[t^{\pm 1}]$, then:*

$$P\gamma \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f} \bullet \eta = \gamma \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f} \bullet \bar{P}\eta.$$

Proof. In the case where P is a power of t , apply the Aut relation with the automorphism of $(\mathfrak{A}, \mathfrak{b})$ given by multiplication by some power of t on \mathfrak{A}' and identity on \mathfrak{A}'' . Conclude with the LV relation. \square

Corollary 7.6. *Assume $\mathfrak{A}' = \frac{\mathbb{Q}[t^{\pm 1}]}{(\pi)}\gamma$. Then $D = \gamma \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f} \bullet P\gamma$ for some $P \in \mathbb{Q}[t^{\pm 1}]$, with $f = f_{vw}^D$.*

Lemma 7.7. *Assume $\mathfrak{A}' = \frac{\mathbb{Q}[t^{\pm 1}]}{((t+1)^m)}\rho \oplus \frac{\mathbb{Q}[t^{\pm 1}]}{((t+1)^m)}\rho'$ with m odd, $\mathfrak{b}(\rho, \rho) = 0$, $\mathfrak{b}(\rho', \rho') = 0$, $\mathfrak{b}(\rho, \rho') = \frac{1}{(t+1)^m}$. Then $D = \rho \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f} \bullet Q\rho'$ for some $Q \in \mathbb{Q}[t^{\pm 1}]$, with $f = f_{vw}^D$.*

Proof. Write $D = \nu \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f} \bullet \nu'$, with $\nu = A\rho + B\rho'$ and $\nu' = A'\rho + B'\rho'$. Applying the Aut relation with the automorphism given by $\rho \mapsto 2\rho$, $\rho' \mapsto \frac{1}{2}\rho'$, and identity on \mathfrak{A}'' , we see that the diagrams $A\rho \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{0} \bullet A'\rho$ and $B\rho' \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{0} \bullet B'\rho'$ are trivial. Hence we can decompose D as follows:

$$D = A\rho \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f_1} \bullet B'\rho' + B\rho' \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f_2} \bullet A'\rho.$$

Now the automorphism given by $\rho \mapsto \rho'$, $\rho' \mapsto t^m\rho$, and identity on \mathfrak{A}'' gives

$$B\rho' \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f_2} \bullet A'\rho = Bt^m\rho \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f_2} \bullet A'\rho'.$$

Thanks to Lemma 7.5, we get $D = \rho \bullet \xrightarrow{\mathfrak{D}} \xrightarrow{f} \bullet P\rho'$, with $P = \bar{A}B' + \bar{B}t^{-m}A'$. \square

Proof of Proposition 7.1. For $\pi \in \mathbb{Q}[t^{\pm 1}]$, the π -component of a $\mathbb{Q}[t^{\pm 1}]$ -module is the submodule of its elements of order some power of π . Any Blanchfield module is the direct sum of its π -components, where π runs through all prime symmetric polynomials (including $t + 1$) and all products of two dual prime non-symmetric polynomials. Thanks to Corollary 7.4, we can assume that \mathfrak{A}' is reduced to one π -component.

First case: $\pi(-1) \neq 0$. The module \mathfrak{A}' can be written as an orthogonal direct sum:

$$\mathfrak{A}' = \bigoplus_{i=1}^p \frac{\mathbb{Q}[t^{\pm 1}]}{(\pi^{n_i})} \gamma_i$$

with $\mathfrak{b}(\gamma_i, \gamma_i) = \frac{P_i}{\pi^{n_i}}$ for some symmetric polynomial P_i prime to π , and $n := n_1 = \dots = n_q > n_{q+1} \geq \dots \geq n_p$. Replacing γ_1 by some rational multiple if necessary, we can assume that $\sum_{i=1}^q P_i$ is prime to π . Set $\gamma = \sum_{i=1}^p \gamma_i$. Then $\mathfrak{b}(\gamma, \gamma) = \frac{P}{\pi^n}$ with P symmetric and prime to π . It follows that the submodule $\langle \gamma \rangle$ of \mathfrak{A}' generated by γ has a trivial intersection with its orthogonal $\langle \gamma \rangle^\perp$, thus:

$$\mathfrak{A}' = \langle \gamma \rangle \oplus^\perp \langle \gamma \rangle^\perp.$$

By Corollaries 7.4 and 7.6, we can decompose D as $D = \sum_{i=1}^p \gamma_i \bullet \overset{\curvearrowright}{\text{D}} \xrightarrow{f_i} \bullet Q_i \gamma_i$ for some polynomials Q_i . Corollary 7.4 gives $D = \gamma \bullet \overset{\curvearrowright}{\text{D}} \xrightarrow{f} \bullet \eta$ with $\eta = \sum_{i=1}^p Q_i \gamma_i$ and $f = \sum_{i=1}^p f_i$. Write $\eta = A\gamma + \mu$ with $\mu \in \langle \gamma \rangle^\perp$. Since $\gamma \bullet \overset{\curvearrowright}{\text{D}} \xrightarrow{0} \bullet \mu = 0$ by Lemma 7.3, we get $D = \gamma \bullet \overset{\curvearrowright}{\text{D}} \xrightarrow{f} \bullet A\gamma$. Similarly $D' = \gamma \bullet \overset{\curvearrowright}{\text{D}} \xrightarrow{f} \bullet B\gamma$. The condition on f implies $\frac{AP}{\pi^n} = \frac{BP}{\pi^n} \pmod{\mathbb{Q}[t^{\pm 1}]}$, thus $A = B \pmod{\pi^n}$ and $A\gamma = B\gamma$.

Second case: $\pi = t + 1$. In that case, the decomposition of \mathfrak{A}' may involve non-cyclic submodules. We have $\mathfrak{A}' = \mathfrak{A}_1 \oplus^\perp \mathfrak{A}_2$ with:

$$\mathfrak{A}_1 = \left(\bigoplus_{i=1}^p \frac{\mathbb{Q}[t^{\pm 1}]}{(t+2+t^{-1})^{n_i}} \gamma_i \right) \quad \mathfrak{A}_2 = \left(\bigoplus_{j=1}^k \left(\frac{\mathbb{Q}[t^{\pm 1}]}{(t+1)^{m_j}} \rho_j \oplus \frac{\mathbb{Q}[t^{\pm 1}]}{(t+1)^{m_j}} \rho'_j \right) \right)$$

with $\mathfrak{b}(\gamma_i, \gamma_i) = \frac{P_i}{(t+2+t^{-1})^{n_i}}$, $P_i(-1) \neq 0$, $\mathfrak{b}(\rho_j, \rho_j) = 0$, $\mathfrak{b}(\rho'_j, \rho'_j) = 0$, $\mathfrak{b}(\rho_j, \rho'_j) = \frac{1}{(t+1)^{m_j}}$, $n_1 = \dots = n_q > n_{q+1} \geq \dots \geq n_p$, $m_1 \geq \dots \geq m_k$, m_j odd. We can assume $\sum_{i=1}^q P_i$ prime to $(t+1)$. Set $\gamma = \sum_{i=1}^p \gamma_i$ and $\rho = \sum_{j=1}^k \rho_j$.

Proceeding as in the first case, applications of Corollaries 7.4 and 7.6 and Lemma 7.7 give:

$$D = \gamma \bullet \overset{\curvearrowright}{\text{D}} \xrightarrow{f_1} \bullet \alpha + \rho \bullet \overset{\curvearrowright}{\text{D}} \xrightarrow{f_2} \bullet \beta,$$

with $\alpha \in \mathfrak{A}_1$ and $\beta \in \mathfrak{A}_2$. Finally, $D = (\gamma + \rho) \bullet \xrightarrow{f} \mathbb{D} \xrightarrow{f} \bullet \eta$ with $\eta \in \mathfrak{A}'$ and $f = f_{vw}^D$.

Similarly $D' = (\gamma + \rho) \bullet \xrightarrow{f} \mathbb{D} \xrightarrow{f} \bullet \eta'$ with $\eta' \in \mathfrak{A}'$.

First assume $2n_1 > m_1$. We have $\mathfrak{b}(\gamma + \rho, \gamma + \rho) = \sum_{i=1}^p \frac{P_i}{(t+2+t^{-1})^{n_i}} = \frac{P}{(t+2+t^{-1})^{n_1}}$ with $P(-1) \neq 0$. We get $\mathfrak{A}' = \langle \gamma + \rho \rangle \oplus^\perp \langle \gamma + \rho \rangle^\perp$ and we conclude as in the first case.

Now assume $m_1 > 2n_1$. It is easily checked that $\langle \gamma + \rho, \rho'_1 \rangle \cap \langle \gamma + \rho, \rho'_1 \rangle^\perp = 0$. Hence $\mathfrak{A}' = \langle \gamma + \rho, \rho'_1 \rangle \oplus^\perp \langle \gamma + \rho, \rho'_1 \rangle^\perp$, and we can assume $\eta, \eta' \in \langle \gamma + \rho, \rho'_1 \rangle$. By Theorem 7.2, there is a basis (μ, μ') of $\langle \gamma + \rho, \rho'_1 \rangle$ such that $\mathfrak{b}(\mu, \mu) = 0$, $\mathfrak{b}(\mu', \mu') = 0$, and $\mathfrak{b}(\mu, \mu') = \frac{1}{(t+1)^{m_1}}$.

By Lemma 7.7, we have $D = \mu \bullet \xrightarrow{f} \mathbb{D} \xrightarrow{f} \bullet A\mu'$ and $D' = \mu \bullet \xrightarrow{f} \mathbb{D} \xrightarrow{f} \bullet B\mu'$. Since the linking f is the same, we get $A = B \pmod{(t+1)^{m_1}}$ and $A\mu' = B\mu'$. \square

Let us fix some notations. Let n be an even positive integer and $N \geq 3n/2$. For $i = 1, \dots, N$, let $(\mathfrak{A}_i, \mathfrak{b}_i)$ be a copy of $(\mathfrak{A}, \mathfrak{b})$ and fix an isomorphism $\xi_i : (\mathfrak{A}, \mathfrak{b}) \xrightarrow{\cong} (\mathfrak{A}_i, \mathfrak{b}_i)$. Let $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ be the orthogonal direct sum of the $(\mathfrak{A}_i, \mathfrak{b}_i)$. Define permutation automorphisms ξ_{ij} of $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ by $\xi_j \circ \xi_i^{-1}$ on \mathfrak{A}_i , $\xi_i \circ \xi_j^{-1}$ on \mathfrak{A}_j and identity on the other \mathfrak{A}_ℓ 's. Given a diagram D with set of univalent vertices V , denote by $D((\gamma_v)_{v \in V}, (f_{vw})_{v \neq w \in V})$ the diagram obtained from D by replacing the label of the vertex v by γ_v and the linking between v and w by f_{vw} . If all the linkings are the same as in D , we drop this part of the notation.

Definition 7.8. An $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ -colored diagram D is *distributed* if there are a decomposition of the set of univalent vertices of D as $V = \sqcup_{i=1}^{|V|/2} \{v_i, w_i\}$ and indices ℓ_i with $\ell_i \neq \ell_j$ if $i \neq j$ such that the labels of v_i and w_i are elements of \mathfrak{A}_{ℓ_i} for all i and the linkings between vertices in different pairs are trivial.

Proposition 7.9. *The space $\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ is generated by distributed $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ -colored diagrams.*

Proof. Let D be an $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ -colored diagram of degree n . First note that D has n trivalent vertices and each univalent vertex is related to a trivalent vertex by an edge since we avoid struts, hence D has at most $3n$ univalent vertices. We shall prove that D is a linear combination of distributed diagrams. Thanks to the LV relation, we can assume that all labels of univalent vertices of D are elements of the \mathfrak{A}_i 's. Thanks to the LD and LV relations, we can assume that all univalent vertices have non-trivial labels and the linking f_{vw} is trivial if v and w are labelled in different \mathfrak{A}_i 's. If D has an odd number of univalent vertices labelled in some \mathfrak{A}_i , application of the automorphism given by opposite identity on \mathfrak{A}_i and identity on the other \mathfrak{A}_j 's shows it is trivial. Assume D has an even number of univalent vertices labelled in each \mathfrak{A}_i . Let i be an index such that the number of univalent vertices of D labelled in \mathfrak{A}_i is maximum; denote $2s$ this number. If $s > 1$, there is an \mathfrak{A}_j that contains no labels of univalent vertices of D . Consider the following automorphism

χ_{ij} of $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$:

$$\chi_{ij}(\gamma) \mapsto \begin{cases} x\gamma + y\xi_j \circ \xi_i^{-1}(\gamma) & \text{if } \gamma \in \mathfrak{A}_i \\ y\xi_i \circ \xi_j^{-1}(\gamma) - x\gamma & \text{if } \gamma \in \mathfrak{A}_j \\ \gamma & \text{if } \gamma \in \mathfrak{A}_\ell \text{ with } \ell \neq i, j \end{cases},$$

where x and y are positive rational numbers such that $x^2 + y^2 = 1$. Apply the Aut relation with χ_{ij} to D and use the LV relation to express $D = D((\gamma_v)_{v \in V})$ as the sum of $x^{2s}D$, $y^{2s}D((\xi_{ij}(\gamma_v))_{v \in V})$ and a linear combination C of diagrams with strictly less than $2s$ vertices in \mathfrak{A}_i and in \mathfrak{A}_j . Now D and $D((\xi_{ij}(\gamma_v))_{v \in V})$ are equal thanks to the Aut relation with ξ_{ij} . It follows that D is a rational multiple of C . Conclude by iterating. \square

Remark. In the case of \mathbb{Z} SK-pairs, Proposition 7.9 is the point in this section which does not work. Indeed, this proposition uses automorphisms χ_{ij} whose definition is based on rational numbers x and y that are not integers. Thus it is not clear whether such isomorphisms are induced by isomorphisms of the underlying integral Blanchfield module. For instance, the next lemma provides an integral Blanchfield module $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ defined as an orthogonal direct sum of two isomorphic Blanchfield modules such that no isomorphism of $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ “mixes” the two summands.

Lemma 7.10. *Let $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ be an integral Blanchfield module defined by $\mathfrak{A}_{\mathbb{Z}} = \frac{\mathbb{Z}[t^{\pm 1}]}{(\delta)}\gamma \oplus \frac{\mathbb{Z}[t^{\pm 1}]}{(\delta)}\eta$ with $\delta(t) = t - 1 + t^{-1}$ and $\mathfrak{b}_{\mathbb{Z}}(\gamma, \gamma) = \mathfrak{b}_{\mathbb{Z}}(\eta, \eta)$. Then any isomorphism of $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ preserves this direct sum decomposition.*

Proof. Throughout the proof, the polynomials are considered in $\frac{\mathbb{Z}[t^{\pm 1}]}{(\delta)}$. An isomorphism of $(\mathfrak{A}_{\mathbb{Z}}, \mathfrak{b}_{\mathbb{Z}})$ has the following form:

$$\begin{cases} \gamma \mapsto P\gamma + Q\eta \\ \eta \mapsto R\gamma + S\eta \end{cases},$$

with $P\bar{P} + Q\bar{Q} = 1$, $R\bar{R} + S\bar{S} = 1$ and $P\bar{R} + Q\bar{S} = 0$. Since δ has degree 2, one can write $P(t) = at + b$ and $Q(t) = ct + d$ with $a, b, c, d \in \mathbb{Z}$. This gives:

$$P\bar{P} + Q\bar{Q} = a^2 + b^2 + ab + c^2 + d^2 + cd = \frac{1}{2}((a+b)^2 + a^2 + b^2 + (c+d)^2 + c^2 + d^2).$$

If $PQ \neq 0$, then $a \neq 0$ or $b \neq 0$, and $c \neq 0$ or $d \neq 0$. It follows that $P\bar{P} + Q\bar{Q} \geq 2$, contradicting the first condition on P and Q . Hence $PQ = 0$ and the conditions on the polynomials P, Q, R, S give $P = S = 0$ or $Q = R = 0$. \square

Recall the map $\iota_n : \mathcal{A}_n(\mathfrak{A}, \mathfrak{b}) \rightarrow \mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ is defined on diagrams by $\iota_n(D((\gamma_v)_{v \in V})) = D((\xi_1(\gamma_v))_{v \in V})$.

Proposition 7.11. *If D is an $(\mathfrak{A}, \mathfrak{b})$ -colored diagram of degree n with an even number of univalent vertices:*

$$\iota_n(D((\gamma_v)_{v \in V}, (f_{vw})_{v \neq w \in V})) = \frac{1}{s!} \sum_{\sigma \in \Upsilon} D((\xi_{\sigma(v)}(\gamma_v))_{v \in V}, (\delta_{\sigma(v)\sigma(w)} f_{vw})_{v \neq w \in V}),$$

where $s = |V|/2$ and $\Upsilon = \{\sigma : V \rightarrow \{1, \dots, s\} \text{ such that } |\sigma^{-1}(i)| = 2 \text{ for all } i = 1, \dots, s\}$.

Proof. We apply the method of the previous proposition with precise computations. We indeed prove a slightly more general result. Consider an $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ -colored diagram $D = D((\gamma_v)_{v \in V \sqcup W}, (f_{vw})_{v \neq w \in V \sqcup W})$ with $|V| = 2s$, $\gamma_v \in \mathfrak{A}_1$ if $v \in V$, $\gamma_w \in \mathfrak{A}_i$ with $i > s$ if $w \in W$ and $f_{vw} = 0$ if $v \in V$ and $w \in W$. We prove by induction on s that in $\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$:

$$D = \frac{1}{s!} \sum_{\sigma \in \Upsilon} D((\xi_{1\sigma(v)}(\gamma_v))_{v \in V} \cup (\gamma_w)_{w \in W}, (\delta_{\sigma(v)\sigma(w)} f_{vw})_{v \neq w \in V} \cup (f_{vw})_{v \neq w \in W}),$$

where the non-indicated linkings are trivial. We will use that our formulas remain valid when permuting the indices of the \mathfrak{A}_i 's, without mentioning it.

The result is trivial if $s = 1$. Take $s > 1$. Applying the Aut relation with χ_{12} to D , we get:

$$D = \sum_{k=0}^s \sum_{\substack{V=V_1 \sqcup V_2 \\ |V_1|=2k}} x^{2k} y^{2(s-k)} D((\xi_{11}(\gamma_v))_{v \in V_1} \cup (\xi_{12}(\gamma_v))_{v \in V_2} \cup (\gamma_w)_{w \in W}),$$

with, for the diagram in the right hand side, the linking $\delta_{\sigma(v)\sigma(w)} f_{vw}$ if $v \neq w$ are both in V_1 or both in V_2 , f_{vw} if $v \neq w$ are both in W and 0 otherwise. Now apply twice the induction hypothesis with V_1 and V_2 instead of V to obtain:

$$(1 - x^{2s} - y^{2s})D = \sum_{k=1}^{s-1} \sum_{\substack{V=V_1 \sqcup V_2 \\ |V_1|=2k}} \frac{x^{2k} y^{2(s-k)}}{k!(s-k)!} \sum_{\sigma \in \Upsilon_1} \sum_{\nu \in \Upsilon_2} D((\xi_{1\sigma(v)}(\gamma_v))_{v \in V_1} \cup (\xi_{1\nu(v)}(\gamma_v))_{v \in V_2} \cup (\gamma_w)_{w \in W}),$$

with the required linkings, where Υ_1 (resp. Υ_2) is defined as Υ with V_1 and $\{1, \dots, k\}$ (resp. V_2 and $\{k+1, \dots, s\}$) instead of V and $\{1, \dots, s\}$. To conclude, note that each diagram in the right hand side occurs once for each value of k . \square

Proof of Theorem 2.11. Define the inverse $\overline{\psi}_n$ of the map $\overline{\psi}_n : \mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}}) \rightarrow \mathcal{A}_n(\delta)$ in the following way. Given a δ -colored diagram D of degree n , denote by e_i , $i = 1, \dots, k$ its edges whose label is non-polynomial. ‘‘Open’’ each such edge e_i as represented in Figure 18, label the created vertices v and w by some γ_v and γ_w in \mathfrak{A}_i such that $\mathfrak{b}(\gamma_v, \gamma_w) = f \pmod{Q[t^{\pm 1}]}$, and fix the linking $f_{vw} = f$. Such γ_v and γ_w always exist: note that γ_v can be chosen




Figure 18: Opening an edge

of order δ , the annihilator of \mathfrak{A}_i , use the non-degeneracy of the Blanchfield form and the fact that the denominator of f has to divide δ . Fix the other linkings to 0, so that we obtain a distributed diagram $\Phi_n(D)$. It does not depend on the numbering of the edges of D thanks to the Aut relation in $\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ with the permutation automorphisms ξ_{ij} . It is also independent of the choice of labels $\gamma_v, \gamma_w \in \mathfrak{A}_i$ by Proposition 7.1. Note that these independence arguments imply that any distributed diagram in $\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ is a $\Phi_n(D)$.

We have to check that the relations defining $\mathcal{A}_n(\delta)$ are respected. It is immediate for AS and IHX. OR follows from the rule $f_{wv}(t) = f_{vw}(t^{-1})$ on linkings. Hol and Hol' are recovered *via* Hol and EV. LE follows from LE when the involved edges have polynomial labels, from LD when one of the involved edges has a polynomial label and from LV when the involved edges have non-polynomial labels. In this latter case, note that one can open this edge with the same label on one univalent vertex for the three diagrams. Finally, we have a well-defined map $\Phi_n : \mathcal{A}_n(\delta) \rightarrow \mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ satisfying by construction $\overline{\psi}_n \circ \Phi_n = \text{id}_{\mathcal{A}_n(\delta)}$. Now Φ_n is surjective by Proposition 7.9. Thus $\overline{\psi}_n$ and Φ_n are inverse isomorphisms. \square

We end with a few results that derive from the above argument and prove useful in the study of the structure of the diagram space $\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$, as it appears in [AM17]. The first one gives a simplified presentation of $\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$.

Proposition 7.12. *Keeping the notations fixed before Definition 7.8, we have:*

$$\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}}) \cong \frac{\mathbb{Q}\langle \text{degree } n \text{ distributed } (\overline{\mathfrak{A}}, \overline{\mathfrak{b}})\text{-colored diagrams} \rangle}{\mathbb{Q}\langle AS, IHX, Hol, OR, LE, LV, EV, LD, Aut_{res} \rangle}$$

where the relation Aut_{res} is the Aut relation restricted to the following automorphisms of $(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$: permutation automorphisms ξ_{ij} , automorphisms fixing one \mathfrak{A}_i setwise and the others pointwise. Moreover, if $(\mathfrak{A}, \mathfrak{b})$ is cyclic, we can further restrict the Aut relation to: permutation automorphisms ξ_{ij} , multiplication by t or -1 on one \mathfrak{A}_i and identity on the others.

Proof. We can see that the space defined by the given presentation is isomorphic to $\mathcal{A}_n(\delta)$ as we did for $\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ in the proof of Theorem 2.11. At the level of generators, the proof of Theorem 2.11 only uses distributed diagrams and at the level of relations, one has to check that the proof of Proposition 7.1 only uses the Aut relation with the allowed automorphisms. \square

In order to study $\mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$, it is natural and helpful to consider the filtration induced by the number of univalent vertices. For $k = 0, \dots, 3n$, let $\mathcal{A}_n^{(k)}(\mathfrak{A}, \mathfrak{b})$ be the subspace of $\mathcal{A}_n(\mathfrak{A}, \mathfrak{b})$ generated by diagrams with at most k univalent vertices and set:

$$\widehat{\mathcal{A}}_n^{(k)}(\mathfrak{A}, \mathfrak{b}) = \frac{\mathbb{Q}\langle (\mathfrak{A}, \mathfrak{b})\text{-colored diagrams of degree } n \text{ with at most } k \text{ univalent vertices} \rangle}{\mathbb{Q}\langle \text{AS, IHX, LE, OR, Hol, LV, EV, LD, Aut} \rangle}.$$

Similarly, let $\mathcal{A}_n^{(k)}(\delta)$ be the subspace of $\mathcal{A}_n(\delta)$ generated by diagrams with at most $k/2$ edges with a non-polynomial label and set:

$$\widehat{\mathcal{A}}_n^{(k)}(\delta) = \frac{\mathbb{Q}\left\langle \begin{array}{c} \delta\text{-colored diagrams of degree } n \\ \text{with at most } k/2 \text{ edges with a non-polynomial label} \end{array} \right\rangle}{\mathbb{Q}\langle \text{AS, IHX, LE, OR, Hol, Hol}' \rangle}.$$

Recall that all these diagram spaces are trivial when n is odd. Moreover, the number of trivalent vertices and the number of univalent vertices in a uni-trivalent graph have the same parity. So we are only interested in cases where n and k are even. Define a map $\widehat{\psi}_n^{(k)} : \widehat{\mathcal{A}}_n^{(k)}(\mathfrak{A}, \mathfrak{b}) \rightarrow \widehat{\mathcal{A}}_n^{(k)}(\delta)$ via pairings of vertices as ψ_n was defined in Section 2.4.

Proposition 7.13. *Let n, k and K be integers such that $0 \leq k \leq 3n$ and $k \leq 2K$. Then:*

- the isomorphism $\overline{\psi}_n : \mathcal{A}_n(\overline{\mathfrak{A}}, \overline{\mathfrak{b}}) \rightarrow \mathcal{A}_n(\delta)$ induces an isomorphism $\mathcal{A}_n^{(k)}(\overline{\mathfrak{A}}, \overline{\mathfrak{b}}) \cong \mathcal{A}_n^{(k)}(\delta)$,
- the map $\widehat{\psi}_n^{(k)} : \widehat{\mathcal{A}}_n^{(k)}((\mathfrak{A}, \mathfrak{b})^{\oplus K}) \rightarrow \widehat{\mathcal{A}}_n^{(k)}(\delta)$ is an isomorphism,
- the space $\widehat{\mathcal{A}}_n^{(k)}((\mathfrak{A}, \mathfrak{b})^{\oplus K})$ admits the following presentation:

$$\widehat{\mathcal{A}}_n^{(k)}((\mathfrak{A}, \mathfrak{b})^{\oplus K}) \cong \frac{\mathbb{Q}\left\langle \begin{array}{c} \text{degree } n \text{ distributed } ((\mathfrak{A}, \mathfrak{b})^{\oplus K})\text{-colored diagrams} \\ \text{with at most } k \text{ univalent vertices} \end{array} \right\rangle}{\mathbb{Q}\langle \text{AS, IHX, Hol, OR, LE, LV, EV, LD, Aut}_{res} \rangle}.$$

Proof. The first point is due to the fact that $\overline{\psi}_n$ identifies the $(\mathfrak{A}, \mathfrak{b})$ -colored diagrams with at most k univalent vertices with the δ -colored diagrams with at most $k/2$ edges with a non-polynomial label. The last two points follow from the same argument as Theorem 2.11 and Proposition 7.12. \square

In [AM17], a Blanchfield module $(\mathfrak{A}, \mathfrak{b})$ is explicated for which $\widehat{\mathcal{A}}_2^{(4)}(\overline{\mathfrak{A}}, \overline{\mathfrak{b}}) \not\cong \mathcal{A}_2^{(4)}(\overline{\mathfrak{A}}, \overline{\mathfrak{b}})$ and this provides the example with a negative answer to Question 2 mentioned in the introduction.

References

- [AL05] E. AUCLAIR & C. LESCOP – “Clover calculus for homology 3-spheres via basic algebraic topology”, *Algebraic & Geometric Topology* **5** (2005), p. 71–106.
- [AM17] B. AUDOUX & D. MOUSSARD – “On some beaded Jacobi diagrams spaces”, soon on arXiv, 2017.
- [Auc06] E. AUCLAIR – “Surfaces et invariants de type fini en dimension 3”, Ph.D. Thesis, Université Joseph Fourier, Grenoble, 2006.
- [Bla57] R. C. BLANCHFIELD – “Intersection theory of manifolds with operators with applications to knot theory”, *Annals of Mathematics (2)* **65** (1957), p. 340–356.
- [GGP01] S. GAROUFALIDIS, M. GOUSSAROV & M. POLYAK – “Calculus of clovers and finite type invariants of 3-manifolds”, *Geometry & Topology* **5** (2001), p. 75–108.
- [GK04] S. GAROUFALIDIS & A. KRICKER – “A rational noncommutative invariant of boundary links”, *Geometry & Topology* **8** (2004), p. 115–204.
- [GR04] S. GAROUFALIDIS & L. ROZANSKY – “The loop expansion of the Kontsevich integral, the null-move and S -equivalence”, *Topology* **43** (2004), no. 5, p. 1183–1210.
- [Hab00a] N. HABEGGER – “Milnor, Johnson, and tree level perturbative invariants”, preprint, 2000.
- [Hab00b] K. HABIRO – “Claspers and finite type invariants of links”, *Geometry and Topology* **4** (2000), p. 1–83.
- [Kri00] A. KRICKER – “The lines of the Kontsevich integral and Rozansky’s rationality conjecture”, arXiv:math/0005284, 2000.
- [Les11] C. LESCOP – “Invariants of knots and 3-manifolds derived from the equivariant linking pairing”, in *Chern-Simons gauge theory: 20 years after*, AMS/IP Stud. Adv. Math., vol. 50, AMS, Providence, RI, 2011, p. 217–242.
- [Les13] —, “A universal equivariant finite type knot invariant defined from configuration space integrals”, arXiv:1306.1705, 2013.
- [Mat87] S. V. MATVEEV – “Generalized surgeries of three-dimensional manifolds and representations of homology spheres”, *Matematicheskie Zametki* **42** (1987), no. 2, p. 268–278.
- [Mou12a] D. MOUSSARD – “Finite type invariants of rational homology 3-spheres”, *Algebraic & Geometric Topology* **12** (2012), no. 4, p. 2389–2428.

- [Mou12b] — , “On Alexander modules and Blanchfield forms of null-homologous knots in rational homology spheres”, *Journal of Knot Theory and its Ramifications* **21** (2012), no. 5, p. 1250042, 21.
- [Mou15] — , “Rational Blanchfield forms, S-equivalence, and null LP-surgeries”, *Bulletin de la Société Mathématique de France* **143** (2015), no. 2, p. 403–431.
- [Mou17] — , “Splitting formulas for the rational lift of the Kontsevich integral”, arXiv:1705.01315, 2017.
- [Oht96] T. OHTSUKI – “Finite type invariants of integral homology 3-spheres”, *J. Knot Theory Ramifications* **5** (1996), no. 1, p. 101–115.