



**HAL**  
open science

## Astrocladistics: Sixteen Years Later

Didier Fraix-Burnet

► **To cite this version:**

Didier Fraix-Burnet. Astrocladistics: Sixteen Years Later. 21st Evolutionary Biology Meeting at Marseilles, Sep 2017, Marseille, France. , 2017. hal-01623694

**HAL Id: hal-01623694**

**<https://hal.science/hal-01623694v1>**

Submitted on 25 Oct 2017


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


### Abstract

The introduction of phylogenetic tools in astrophysics started in 2001 and was presented in 2008 at the 12th Evolutionary Biology Meeting at Marseille. Many results have been obtained, essentially using Maximum Parsimony (cladistics), covering very diverse kinds of objects of the Universe : galaxies, dwarfs galaxies, gamma ray bursts, globular clusters, stellar populations, stars, jovian satellites, supernovae. These successes open some new challenges for astrocladistics.

### Achievements



2008 session of the Marseille conference



Fraix-Burnet, Choler, Douzery 2006

### Challenges and Perspectives

#### Cladistics and Classification

We are now facing several important questions:


- ▶ how to **synthetise several cladograms** obtained with different traits?
- ▶ distance-based approaches and networks: **which distance?**
- ▶ **phenetic classification** first then cladistics?

The PHYLOCODE project initially inspired us, similar difficulties?

#### Pre-classification with Hierarchical Clustering

1494 galaxies have been pre-clustered into 300 pre-groups with a **hierarchical clustering method** followed by a **cladistic analysis**.

Eleven parameters are used: *B-V*, *logRe*, *surface brightness*, *Hβ*, *D4000*, *Mass*, *Sersic index n*, *Hα+NII*, *Gband*, *Mg* and *Na*.




Most parsimonious tree on the 300 "pre-groups" (leaves). The nine groups ("clades") shown in colour are the larger ones with more than 30 galaxies.

#### Some results on galaxies

Initial motivation: renewal of the classification of galaxies.

#### First "general" multivariate classification of (≈1000) galaxies:




Fraix-Burnet et al 2012

But this is only for 1000 galaxies...

#### Why it works

Demonstration with the **evolutionary tracks of stellar evolution** using 11 parameters (traits).



Only phylogenetic tools like cladistics are able to reconstruct the families of stars related by evolution! This problem is called the **chemical tagging of stars**.

#### Many applications in astrophysics

- ▶ More than **16 papers** by our team
- ▶ 2 papers and 2 posters by four other teams

A blog: <https://astrocladistics.org/>



### Conclusion

- ▶ Phylogenetic tools: marvelous techniques to **reconstruct evolutionary paths**, whatever the nature of evolution.
- ▶ Great **success in astrophysics**, more and more applications.
- ▶ **Connections with physical models** to be improved.
- ▶ **Still no multivariate classification of galaxies**.
- ▶ **Networks** better suited to represent galaxy diversification?