



High-order hybrid discontinuous galerkin methods for diffusion-convection equations with evanescent diffusion

Loïc Dijoux, Vincent Fontaine

► To cite this version:

Loïc Dijoux, Vincent Fontaine. High-order hybrid discontinuous galerkin methods for diffusion-convection equations with evanescent diffusion. InterPore, May 2017, Rotterdam, Netherlands. hal-01625960

HAL Id: hal-01625960


<https://hal.science/hal-01625960v1>

Submitted on 30 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High-order hybrid discontinuous galerkin methods for diffusion-convection equations with evanescent diffusion



Loïc DIJOUX, Vincent FONTAINE
University of La Réunion



Physical context

- Diffusion-advection equation

$$\vec{\nabla} \cdot (-\alpha \vec{\nabla} u + \beta u) = f \quad \text{in } \Omega$$

where α : the conductivity tensor, u : the state variable, β : velocity advection field and f : source term.

Application field:

- Ground water flow
- Heat and Mass transfert, ...

A mixed formulation

We introduce an auxiliary variable \mathbf{Q}_d :

$$\begin{cases} \mathbf{Q}_d = -\alpha^{\frac{1}{2}} \vec{\nabla} u \\ \nabla \cdot \mathbf{q}_d + \nabla \cdot \beta u = f \end{cases}$$

where $\mathbf{q}_d = \alpha^{\frac{1}{2}} \mathbf{Q}_d$. We introduce two approximate variables u_h and \mathbf{Q}_h of respectively u and \mathbf{Q}_d :

$$\begin{aligned} \mathbf{Q}_h &= \sum_{i=1}^{n_q} Q_i \mathbf{w}_i \\ u_h &= \sum_{j=1}^{n_u} u_j v_j \end{aligned}$$

where n_q and n_u are the degrees of freedom of \mathbf{Q}_h and u_h respectively. \mathbf{w}_i and v_j are their corresponding interpolations functions.

Approximation spaces

We define finite element spaces:

$$\mathcal{W}_h^p = \{\mathbf{w} \in [L^2(\Omega)]^d : \mathbf{w}|_K \in \mathcal{W}^p(K), \quad \forall K \in \mathcal{T}_h\},$$

$$\mathcal{V}_h^p = \{v \in L^2(\Omega) : v|_K \in \mathcal{V}^p(K), \quad \forall K \in \mathcal{T}_h\},$$

and add a traced finite element space:

$$\mathcal{M}_h^p = \{\mu \in L^2(\varepsilon_h) : \mu|_e \in \mathcal{M}^p(e), \quad \forall e \in \varepsilon_h\}$$

Discrete weak formulation

We impose the continuity of $\widehat{\alpha^{\frac{1}{2}} \mathbf{Q}_h} + \widehat{\beta u_h}$ on each edge $e \in \varepsilon_h$ through one more equation and finally write :

Find $(\mathbf{Q}_h, u_h, \lambda_h) \in \mathcal{W}_h^p \times \mathcal{V}_h^p \times \mathcal{M}_h^p$ such that :

$$\begin{aligned} (\mathbf{Q}_h, \mathbf{w}_h)_{\mathcal{T}_h} - (u_h, \nabla \cdot (\alpha^{\frac{1}{2}} \mathbf{w}_h))_{\mathcal{T}_h} + \langle \lambda_h, \alpha^{\frac{1}{2}} \mathbf{w}_h \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h} &= -\langle g_D, \alpha^{\frac{1}{2}} \mathbf{w}_h \cdot \mathbf{n} \rangle_{\Gamma_D} \quad \forall \mathbf{w}_h \in \mathcal{W}_h^p \\ \langle \alpha^{\frac{1}{2}} \mathbf{Q}_h + \widehat{\beta u_h}, v_h \rangle_{\partial \mathcal{T}_h} - \langle \alpha^{\frac{1}{2}} \mathbf{Q}_h + \beta u_h, \nabla v_h \rangle_{\mathcal{T}_h} &= (f, v_h)_{\mathcal{T}_h} \quad \forall v_h \in \mathcal{V}_h^p \\ \langle [\alpha^{\frac{1}{2}} \mathbf{Q}_h + \widehat{\beta u_h}], \mu_h \rangle_{\varepsilon_h} &= -\langle g_N, \mu_h \rangle_{\Gamma_N} \quad \forall \mu_h \in \mathcal{M}_h^p \end{aligned}$$

Definition of $\hat{\mathbf{q}}_h$ (see [1, 3]) on each face $e \in \partial \mathcal{T}_h$:

$$\hat{\mathbf{q}}_h \cdot \mathbf{n}_e + \widehat{\beta u_h} \cdot \mathbf{n}_e = \mathbf{q}_h \cdot \mathbf{n}_e + \beta \cdot \mathbf{n}_e \lambda_h + \tau_e (P u_h - \lambda_h)$$

where $\tau_e = \sqrt{\det(\alpha)} h^{-1} + \text{MAX}(\beta \cdot \mathbf{n}_e, 0)$, see [2] and [1]

Examples of HdG methods:

Methods	$\mathcal{W}_h^p(K)$	$\mathcal{V}_h^p(K)$	$\mathcal{M}_h^p(e)$	τ_e	$\ u - u_h\ ^{+}$	$\ \mathbf{q} - \mathbf{q}_h^*\ ^{+}$
RT_p	$(P^p)^d + x P^p$	P^p	P^p	0	$p+1$	$p+1$
HdG_p	$(P^p)^d$	P^p	P^p	$kh^{-1} + \beta \cdot \mathbf{n}_e $	$p+1$	$p+1$
HdG_p^b	$(P^p)^d$	P^{p+1}	P^p	$kh^{-1} + \beta \cdot \mathbf{n}_e $	$p+2$	$p+1$
HdG_p^m	$(P^p)^d + x P^p$	P^{p+1}	P^p	$\sqrt{\det(\alpha)} h^{-1} + \text{MAX}(\beta \cdot \mathbf{n}_e, 0)$	$p+2$	$p+1$

+ : for $\beta = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

Numerical results : degenerate case


We propose to study a heterogeneous and anisotropic case with the exact solution defined in $\Omega \in (0, 1) \times (0, \frac{1}{3})$ as :

$$\begin{cases} u(x, y) = 1 & \text{for } \Omega_1 \in (0, \frac{1}{3}) \times (0, \frac{1}{3}) \\ u(x, y) = 1 & \text{for } \Omega_2 \in (\frac{1}{3}, \frac{2}{3}) \times (0, \frac{1}{3}) \\ u(x, y) = \frac{e - e^x}{e} & \text{for } \Omega_3 \in (\frac{2}{3}, 1) \times (0, \frac{1}{3}) \end{cases}$$



For Ω_1, Ω_3 we have the conductivity tensor $\alpha = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and for Ω_2 we have $\alpha = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$. The velocity field is defined in Ω and we have $\beta = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$.

Numerical results : degenerate case

Triangular mesh (GMSH : 7547 faces) & HdG_p^m :



Discrete solution : $u_h \in \mathcal{W}_h^2$:



Bibliography

References

- C. LEHRENFELD, *Hybrid discontinuous galerkin methods for solving incompressible flow problems*, PhD thesis, RWTH Aachen University, 2010.
- N. C. NGUYEN, J. PERAIRE, AND B. COCKBURN, *An implicit high-order hybridizable discontinuous Galerkin method for linear convection-diffusion equations*, Journal of Computational Physics, 228 (2009), pp. 3232–3254.
- I. OIKAWA, *A hybridized discontinuous galerkin method with reduced stabilization*, Journal of Scientific Computing, 65 (2015), pp. 327–340.

Contact

Loïc DIJOUX : loic.dijoux2@univ-reunion.fr
Vincent FONTAINE : vincent.fontaine@univ-reunion.fr