



**HAL**  
open science

## Towards a genomic monitoring of intra-breed sheep variability

Liesbet François, Grégoire Leroy, Guillaume Baloche, Jérôme Raoul, Coralie Danchin-Burge, Denis Laloë

### ► To cite this version:

Liesbet François, Grégoire Leroy, Guillaume Baloche, Jérôme Raoul, Coralie Danchin-Burge, et al.. Towards a genomic monitoring of intra-breed sheep variability. 64. Annual Meeting of the European Federation of Animal Science (EAAP), Aug 2013, Nantes, France. Wageningen Academic Publishers, Annual Meeting of the European Association for Animal Production, 19 (1ère Ed.), 665 p., 2013, Annual Meeting of the European Association for Animal Production. hal-01628092

**HAL Id: hal-01628092**

**<https://hal.science/hal-01628092>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Whole genome association study for reproductive seasonality trait in the Rasa Aragonesa sheep breed**A. Martínez-Royo<sup>1</sup>, J.L. Alabart<sup>1</sup>, J. Folch<sup>1</sup>, B. Lahoz<sup>1</sup>, E. Fantova<sup>2</sup> and J.H. Calvo<sup>1,3</sup><sup>1</sup>CITA, Unidad de Tecnología en Producción Animal, Avda Montañana 930, 50059 Spain, <sup>2</sup>Oviaragón, Grupo Pastores, Camino Cogullada s/n, 50014 Spain, <sup>3</sup>Fundación ARAID, C/ María de Luna 11, 1<sup>a</sup>, 50018 Spain; amartinezroyo@aragon.es

Many sheep breeds from the Mediterranean area have seasonal patterns of oestrous behaviour and ovulation. Maximal reproductive activities occur from August to March. This reproductive seasonality induces great variation in lamb production and, therefore, in the market price of lamb meat. This spring ovulatory activity is under genetic control, yet to date only a small proportion of the total variation has been explained by genes identified through linkage analysis and candidate gene association studies. To address the lack of information at the genomic level we present a results from a preliminary whole genome association study that was performed using data on reproductive seasonality for 141 ewes of the Rasa Aragonesa Spanish sheep breed that were genotyped using the OvineSNP50K BeadChip platform of Illumina®. Biannual registration data of seasonality was measured in an experimental flock as total days of anestrus (TDA) based on weekly blood progesterone levels from January to August, and corrected for age and body condition score. TDA was defined as the sum of periods in which three or more consecutive samplings having plasmatic progesterone levels lower than 0.5 ng/ml. Results have pointed to new regions across the genome not previously described that are potentially associated with natural cyclicality in this breed. Use of a greater number of samples and next generation high density BeadChips could help to accurately identify loci that are involved in controlling the seasonality of sheep. Financed by Ministerio de Economía y Competitividad and FEDER (INNPACT Project IPT-010000-2010-33) and INIA (B. Lahoz grant).

**Towards a genomic monitoring of intra-breed sheep variability**L. François<sup>1,2,3,4</sup>, G. Leroy<sup>1,5</sup>, G. Baloche<sup>6</sup>, J. Raoul<sup>2,3</sup>, C. Danchin-Burge<sup>2,3</sup> and D. Laloë<sup>1,5</sup><sup>1</sup>AgroParis Tech, UMR1313 GABI, 16 Rue Claude Bernard, 75231 Paris, France, <sup>2</sup>IDELE, Animal Genetics, BP 42118, 31321 Castanet-Tolosan, France, <sup>3</sup>IDELE, Animal Genetics, 149 Rue de Bercy, 75595 Paris, France, <sup>4</sup>BOKU, Institut für Nutztierwissenschaften, Gregor Mendel Straße 33, 1180 Vienna, Austria, <sup>5</sup>INRA, UMR1313 GABI, CRJ, 78352 Jouy-en-Josas, France, <sup>6</sup>INRA, SAGA, BP 52627, 31326 Castanet-Tolosan, France; francois.liesbeth@gmail.com

Limiting the number of breeding stock used can result in the loss of genetic variability in most breeds, and may restrict the capacity to improve and adapt breeds to future needs. The objective of the VARUME project (genetic Variability of RUMinants and Equine species) was to quantify the genetic variability in French ruminant and equine species, based both on pedigree and molecular data. The selection program of the four main dairy sheep breeds in France (Lacaune, Manech Tête rousse, Manech Tête noire and Basco-Béarnaise) is based on the selection of a few elite breeding animals. To assess the variability in genomic information, males were genotyped using the SNP50 chip. The genetic variability in these four breeds will be determined by calculating the effective population size based on linkage disequilibrium over multiple generations. As the effective population size may depend on parameters, data source and methods used, a comparison will be made with the effective population size based on pedigree information, the differences will be interpreted considering the size of the sample and structure of the population. This information can be used to observe the evolution of genetic variability in the breeds and in turn improve the breed's capability for adaptation.

# **Book of Abstracts of the 64<sup>th</sup> Annual Meeting of the European Federation of Animal Science**



**Book of abstracts No. 19 (2013)  
Nantes, France  
26 - 30 August 2013**