



**HAL**  
open science

## **A decision support tool to predict mass transfer and microorganism growth in modified atmosphere packaging of fresh food products**

Estelle Chaix, Bertrand Broyart, Carole Guillaume, Olivier Couvert, Valérie Guillard

### ► To cite this version:

Estelle Chaix, Bertrand Broyart, Carole Guillaume, Olivier Couvert, Valérie Guillard. A decision support tool to predict mass transfer and microorganism growth in modified atmosphere packaging of fresh food products. 2012 EFFoST Annual Meeting - A Lunch Box for Tomorrow: An interactive combination of integrated analysis and specialized knowledge of food, Nov 2012, Montpellier, France. 2012. hal-01628134

**HAL Id: hal-01628134**

**<https://hal.science/hal-01628134>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



a8e67

# A Decision Support Tool to Predict Mass Transfer and Microorganism Growth in Modified Atmosphere Packaging of Fresh Food Products

E. Chaix <sup>(1)</sup>, B. Broyart <sup>(3)</sup>, C. Guillaume <sup>(1)</sup>, O. Couvert <sup>(2)</sup>, V. Guillard\* <sup>(1)</sup>

\* Contact: guillard@univ-montp2.fr

(1) UMR 1208 IATE Agropolymers Engineering and Emerging Technologies: UM2 – CIRAD – INRA – Montpellier SupAgro, Montpellier, France

(2) EA3882, Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, UMT 08.3 PHYSI'Opt, IFR148 ScInBioS, Quimper, France

(3) UMR 1145 GENIAL Food Engineering: INRA – AgroParisTech – CNAM, Massy, France

## Context of the study:

In **Modified Atmosphere Packaging** system (MAP) of fresh food products, one of the main roles assigned to the packaging material is to control gas transfer from the surrounding atmosphere toward headspace. Beyond gas losses toward external atmosphere, the gases dissolve and diffuse in the product leading to gas modification in headspace. Most of the studies recently carried out to evaluate the impact of gases on microorganism growth used the "pack and pray" empirical methodology. Different gas mixtures (mainly O<sub>2</sub>/CO<sub>2</sub>/N<sub>2</sub>) are successively tested on the product until the right combination (O<sub>2</sub>/CO<sub>2</sub> ratio, package surface/volume, mass of food, etc.) is found. In this approach, mass transfer phenomena through the packaging material are seldom taken into account.

The **objective** of this work is to develop a **Decision Support Tool** aiming at predicting the shelf life of fresh food. This tool is based on mathematical models coupling mass transfer models and predictive microbiology. The originality of the approach is that it takes into account, in a dynamic system, the impact of gradients of dissolved gases within the product (due to mass transfer through packaging and within headspace and food matrix) on microorganism growth, that was never done until now in predictive microbiology. It would be helpful to simplify the package design steps by being able to predict in advance the required window of packaging permeability for maintaining the quality of the packed food.

## Theory

### Mass transfer

The mass transfer is supposed monodirectional along the [O x] abscissa and modeled using Fick's laws. In headspace, the concentration of species is considered uniform along the [-e<sub>F</sub> 0] axis. The tray in which is placed the food is supposed to be impermeable to any gas flow except through the lid film (called 'packaging film' on the schema). The total pressure inside the headspace is supposed constant and **variations of headspace volume (V<sub>HS</sub>)** due to mass transfer are thus taken into account as follows:

$$\frac{dV_{HS}}{dt} = \frac{RT}{P_i} \left( \frac{\phi_{O_2} + \phi_{O_2,F}}{M_{O_2}} + \frac{\phi_{D_2} + \phi_{D_2,F}}{M_{D_2}} + \frac{\phi_{N_2,F}}{M_{N_2}} \right)$$

Variation of [gases] in headspace

$$\frac{dV_{HS} C_{i,k}}{dt} = \phi_{j,I} + \phi_{j,F}$$


Mass flow of gases through the packaging material

$$\phi_{j,F} = M_j P_j (S_F / e_F) (p_{j,e} - p_{j,i})$$

Mass flow of gases through food/packaging interface

$$\begin{cases} \phi_{j,I} = \frac{kM_j}{RT} (p_{j,i} - p_{j,e}) S_i \\ D_j \frac{\partial C_{j,i}(0,t)}{\partial x} = \phi_{j,I} / S_i = \frac{kM_j}{RT} (p_{j,i} - p_{j,e}) \end{cases}$$

with  $p_{j,e} = \frac{C_{j,i}}{M_j K_j}$



Microorganism growth (Rosso's model)

$$\begin{cases} \frac{dN}{dt} = 0 & \text{for } t \leq \text{lag} \\ \frac{dN}{dt} = \mu_{max} N \left( 1 - \frac{N}{N_{max}} \right) & \text{for } t > \text{lag} \end{cases}$$

Nomenclature

- $j$ =[O<sub>2</sub>,N<sub>2</sub>] represents respectively oxygen, carbon dioxide and nitrogen
- $\phi$  = gaseous phase;  $\phi_i$  interface and  $\phi_F$  = film
- $V_{HS}$  = volume of headspace (m<sup>3</sup>)
- $M_j$  = molar mass (kg.mol<sup>-1</sup>) of species
- $P_i$  = permeability (mol.m<sup>-1</sup>.s<sup>-1</sup>.Pa<sup>-1</sup>) of gas
- $S_i$  = film surface (m<sup>2</sup>)
- $e_F$  = film thickness (m)
- $p_{j,e}$  and  $p_{j,i}$  = external (atmospheric) and headspace gas partial pressure (Pa) of species
- $S$  = external mass transfer coefficient (m.s<sup>-1</sup>)
- $R$  = perfect gas constant (J.mole<sup>-1</sup>.K<sup>-1</sup>)
- $T$  = temperature (°K)
- $p_{j,i}$  = partial pressure of gas  $j$  at the interface/headspace/food (Pa)
- $S_i$  = interface headspace/food surface
- $K_j$  = Henry's coefficient of species  $j$  (in mole.Pa<sup>-1</sup>.m<sup>3</sup>)
- $D_j$  = diffusivity of the species  $j$  in the food (m<sup>2</sup>.s<sup>-1</sup>)
- $x$  = distance from headspace/food interface (m)
- $t$  = time (s)
- $S_j$  = net volume production rate of species  $j$  in the food due to reactions (kg.m<sup>-3</sup>.s<sup>-1</sup>)
- $N_{max}$  = maximal number of microorganisms (CFU/g) at the end of the growth curve
- $N$  = number of microorganisms at time  $t$  (CFU/g)
- $\mu_{max}$  = maximal growth rate
- $\mu$  = inhibitor or activator parameters
- $\mu_{opt}$  = optimal growth rate
- $S_2$  = net volume production rate of species  $j$  in the food due to reactions (kg.m<sup>-3</sup>.s<sup>-1</sup>)
- MIC = Minimal Inhibitory Concentration
- CO<sub>2</sub> = concentration of CO<sub>2</sub> (in %) imposed to the microorganism culture as a function of time and position in the food

### Predictive microbiology

Gamma-concept (Zwietering et al., 1992) was chosen to simulate microbial growth  $\mu_{max}$  rate as:


$$\mu_{max} = \mu_{opt} \cdot \gamma_T \cdot \gamma_{pH} \cdot \gamma_a \cdot \gamma_{im} \cdot \gamma_{CO_2}$$

Where  $\mu_{opt}$  is the optimal growth rate and  $\gamma_i$  are functions corresponding to each environmental factors effects (i.e. temperature, pH, water activity, interactions and CO<sub>2</sub>). To take into account the inhibitor effect of CO<sub>2</sub>, we consider that:

$$\gamma_{CO_2} = 1 - \left( \frac{\%CO_2(t,x)}{MIC} \right)^a$$

Finally, microbial concentration  $N$  is given according to the primary growth model of Rosso (1995).

## Preliminary simulations



**Conclusion:** This Decision Support Tool appears very interesting from a practical point of view, to help food manufacturer to choose the packaging material that best suits to a given food product to ensure food safety.

**Perspectives:** To go further, this modeling tool should be experimentally validated. An update version taking into account the respiration of microorganisms and its impact on gases concentrations is foreseen. A global sensitivity analysis will be performed to identify the less influential factors among the existing input parameters in order to simplify the model and reduce the experimental cost of parameters determination.

This work is done in the project MAP'Opt (2011-2015), funded by the French National Research Agency, whose full name is "Equilibrium gas composition in modified atmosphere packaging and food quality".

Zwietering M.H., Wijnen T., De Wit, J.C. & Van't Riet K., 1992. A decision support system for prediction of the microbial spoilage in foods. *J. Food Protection*, 55, 979-979.  
Rosso L., Lohry J.R., Beard S. and Rånby J.P. (1995). A convenient model to describe the combined effects of temperature and pH on microbial growth. *Appl. Env. Microbiol.* 61, 610-616.