



HAL
open science

On the stable Andreadakis problem

Jacques Darné

► **To cite this version:**

| Jacques Darné. On the stable Andreadakis problem. 2017. hal-01635093v2

HAL Id: hal-01635093

<https://hal.science/hal-01635093v2>

Preprint submitted on 27 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the stable Andreadakis problem

Jacques DARNÉ

February 27, 2018

Abstract

Let F_n be the free group on n generators. Consider the group IA_n of automorphisms of F_n acting trivially on its abelianization. There are two canonical filtrations on IA_n : the first one is its lower central series Γ_* ; the second one is the Andreadakis filtration \mathcal{A}_* , defined from the action on F_n . In this paper, we establish that the canonical morphism between the associated graded Lie rings $\mathcal{L}(\Gamma_*)$ and $\mathcal{L}(\mathcal{A}_*)$ is stably surjective. We then investigate a p -restricted version of the Andreadakis problem. A calculation of the Lie algebra of the classical congruence group is also included.

Introduction

Automorphisms of free groups have been widely studied over the years, from many different points of view. They are linked to the *mapping class groups* of surfaces and braid groups [FM12a]; they also act on a moduli space of graphs, called the *outer space*, introduced in [CV86], which is still actively studied nowadays (see, for instance, [BBM07], or [FM12b]). Recently, several results have also been obtained regarding the stable homology of these groups [Gal11, RWW17, DV15, Dja16a].

One way to try and understand the structure of these automorphism groups is to cut them into pieces, by considering a family of subgroups and studying how these interact with each other. Such families of subgroups can arise from the action on the free group F_n and related geometric objects, as is the case with the *automorphisms with boundaries* (see for instance [JW04] or [DP12]), and for the *Andreadakis subgroups*, which we now focus on.

The first Andreadakis subgroup of $\text{Aut}(F_n)$ is the IA -group. Precisely, we can first look at how automorphisms act on $F_n^{ab} \cong \mathbb{Z}^n$. That is, we can consider the projection from $\text{Aut}(F_n)$ onto $GL_n(\mathbb{Z})$. We then put aside this linear part by considering only IA_n , the subgroup of automorphisms acting trivially on \mathbb{Z}^n , which is an algebraic analogue of the *Torelli subgroup* of the mapping class group. An explicit finite set of generators of IA_n has been known for a long time [Nie24] – see also [BBM07, 5.6]. Nevertheless, the structure of IA_n remains largely mysterious. For instance, IA_3 is not finitely presented [KM97], and it is not known if IA_n is finitely presented for $n > 3$. Recent results about the IA -groups include the finite L -presentation of IA_n given in [DP16], or finiteness results on the lower central series of IA_n obtained in [CP17].

The IA -group is the first step of the *Andreadakis filtration* $IA_n = \mathcal{A}_1 \supseteq \mathcal{A}_2 \supseteq \dots$, in which \mathcal{A}_j is the group of automorphisms acting trivially on $F_n/\Gamma_{j+1}(F_n)$, where F_n is filtered by its lower central series $F_n = \Gamma_1(F_n) \supseteq \Gamma_2(F_n) \supseteq \dots$. The Andreadakis filtration is an N -series. As such, it contains the minimal N -series on IA_n , its lower

central series: for all k , $\mathcal{A}_k \supseteq \Gamma_k(IA_n)$. We are thus led to the problem of comparing these filtrations, that we call the *Andreadakis problem*.

Since the two filtrations are N -series, the associated graded objects are graded Lie rings (that is, Lie algebras over \mathbb{Z}), the Lie bracket being induced by the commutator map $(x, y) \mapsto [x, y] = xyx^{-1}y^{-1}$. The inclusion $i : \Gamma_*(IA_n) \subseteq \mathcal{A}_*$ induces a morphism of Lie rings:

$$i_* : \mathcal{L}(\Gamma_*(IA_n)) = \bigoplus_{j \geq 1} \Gamma_j(IA_n)/\Gamma_{j+1}(IA_n) \longrightarrow \mathcal{L}(\mathcal{A}_*) = \bigoplus_{j \geq 1} \mathcal{A}_j/\mathcal{A}_{j+1}. \quad (0.0.1)$$

Thus, the Andreadakis problem translates into the following question:

Problem 1 (Andreadakis). *How close is the morphism (0.0.1) to be an isomorphism?*

Andreadakis conjectured that the filtrations were the same [And65, p. 253]. In [Bar13], Bartholdi disproved the conjecture, using computer calculations. He then tried to prove that the two filtrations were the same up to finite index, but in the erratum [Bar16], he showed that even this weaker statement cannot be true. His proof uses the L -presentation of IA_n given in [DP16], to which he applies algorithmic methods described in [BEH08] to calculate (using the software GAP) the first degrees of the Lie algebra associated to each filtration.

In this paper, we are interested in the difference between $\mathcal{A}_k(F_n)$ and $\Gamma_k(IA_n)$ for $n \gg k$, that is, in the stable range. We thus ask the following question:

Problem 2 (Andreadakis - stable version). *How close is the morphism*

$$i_* : \mathcal{L}_k(\Gamma_*(IA_n)) = \Gamma_k(IA_n)/\Gamma_{k+1}(IA_n) \longrightarrow \mathcal{L}_k(\mathcal{A}_*) = \mathcal{A}_k/\mathcal{A}_{k+1} \quad (0.0.2)$$

to be an isomorphism when $n \gg k$?

Our main goal here is to show the following partial answer to this question.

Theorem 2.38 (Stable surjectivity). *When $n \geq k + 2$, the morphism (0.0.2) is surjective.*

A (weaker) rational version of this theorem has been obtained independently by Massuyeau and Sakasai [MS17, th. 5.1]. Like them, we prove it by building on results from [Sat12], but using quite different methods. These methods include a description of Andreadakis-like filtrations *via* a categorical framework, allowing us to state and study a p -restricted version of the problem. We answer the questions asked in [HM17] about this problem, and use our answers to study the stable p -restricted Andreadakis problem. Also, we solve the stable q -torsion Andreadakis problem for \mathbb{Z}^n , getting a complete calculation of the Lie ring of the congruence group $\mathrm{GL}_n(p\mathbb{Z})$ for $n \geq 5$.

Let us now describe in more detail the methods we use and the results contained in the present paper. In **section 1**, we set up a general framework for understanding N -series and their associated Lie algebras. We introduce a category \mathcal{SCF} of N -series. We remark that the categorical definition of an *action* of an object on another makes sense in this category. This allows us to interpret an old construction of Kaloujnine (see theorem 1.16) as the construction of universal actions in \mathcal{SCF} . This category is thus *action-representative*, a situation studied in [BJK05, BB07, Bou08]. Using this language, we are able to recover and generalize several classical constructions:

- Taking the graded rings associated to N -series gives a functor \mathcal{L} from \mathcal{SCF} to the category of Lie rings. This functor preserves actions, and the *Johnson morphism* admits a nice generalisation as the classifying morphism associated to an action between Lie rings obtained from an action in \mathcal{SCF} .
- *Lazard's classical construction* of N -series from algebra filtrations described in [Laz54] is recovered as a particular case of Kaloujnine's construction.
- We also obtain the *filtrations on congruence groups* studied in [Lop14].

In particular, we show that the filtration given by the last construction on the classical congruence group $GL_n(q\mathbb{Z})$ coincides with its lower central series when $n \geq 5$. As a consequence, we get an explicit calculation of this group's Lie ring (generalizing [LS76, Th. 1.1], which is the degree-one part):

Corollary 1.46. *For all $n \geq 5$ and all $q \geq 3$, there is a canonical isomorphism of graded Lie rings (in degrees at least one):*

$$\mathcal{L}(GL_n(q\mathbb{Z})) \cong \mathfrak{sl}_n(\mathbb{Z}/q)[t],$$

where the degree of t is 1, and the Lie bracket of Mt^i and Nt^j is $[M, N]t^{i+j}$.

Section 2 deals with the proof of our stable surjectivity result (Theorem 2.38). The proof relies on the constructions of the first section, applied to Fox's free differential calculus. The *Jacobian matrix* map $D : f \mapsto Df$ turns out to be a *derivation* from $\text{Aut}(F_n)$ to $GL_n(\mathbb{Z}F_n)$, sending the Andreadakis filtration to the congruence filtration $GL_n((IF_n)^*)$ (the group algebra $\mathbb{Z}F_n$ being filtered by the powers of its augmentation ideal IF_n). We then study such derivations, and the maps they induce on the graded Lie rings associated to N -series they preserve. We thus show that the *trace map* defined by $\text{Tr}(f) = \text{Tr}(Df - 1_n)$ induces a well-defined map:

$$\text{Tr} : \mathcal{L}(\mathcal{A}_*) \longrightarrow \text{gr}(\mathbb{Z}F_n).$$

The graded algebra $\text{gr}(\mathbb{Z}F_n)$ is in fact the tensor algebra TV over $V = F_n^{ab}$. A result from [BLGM90] implies that this trace map takes values in $[TV, TV]$. Studying the links between free differential calculus and differential calculus in TV , we show that this trace map is exactly the one introduced by Morita [Mor93, Def. 6.4], getting the explicit description in terms of contraction maps notably used by Satoh in [Sat12]. Denoting the Johnson morphisms by τ and τ' , we get a commutative diagram of graded linear maps:

$$\begin{array}{ccc} \mathcal{L}(IA_n) & & \\ i_* \downarrow & \searrow^{\tau'} & \\ \mathcal{L}(\mathcal{A}_*) & \xleftarrow{\tau} \text{Der}(\mathfrak{L}V) & \xrightarrow{\text{Tr}_M} TV, \end{array}$$

where $\text{Tr}_M \circ \tau = \text{Tr}$. This gives the following inclusions of subspaces of $\text{Der}(\mathfrak{L}V)$:

$$\text{Im}(\tau') \subseteq \mathcal{L}(\mathcal{A}_*) \subseteq \text{Tr}_M^{-1}([TV, TV]).$$

We observe that calculations from [Sat12] work over \mathbb{Z} . From this, we deduce that the subspaces $\text{Im}(\tau')$ and $\text{Tr}_M^{-1}([TV, TV])$ are stably the same, so these inclusions are equalities in the stable range, and i_* must be stably surjective. We close the section by

investigating some of the consequences of this result for automorphisms of free nilpotent groups.

In **section 3**, we turn to the *p*-restricted version of the Andreadakis problem. Precisely, we can do the same construction as above, replacing the lower central series $\Gamma_*(F_n)$ by the mod-*p* lower central series $\Gamma_*^{[p]}(F_n)$, which is an N_p -series:

$$(\Gamma_i^{[p]})^p \subseteq \Gamma_{pi}^{[p]}.$$

Kaloujnine’s construction gives an associated Andreadakis filtration $\mathcal{A}_*^{[p]}$ on the group $IA^{[p]}$ of automorphisms of F_n acting trivially on $F_n^{ab} \otimes \mathbb{F}_p$. This filtration was shown in [HM17] to be an N_p -series. It then contains the minimal N_p -series $\Gamma_*^{[p]}(IA^{[p]})$. Whence the natural question:

Problem 3 (Andreadakis – *p*-restricted version). *What is the difference between the N_p -series $\mathcal{A}_*^{[p]}$ and $\Gamma_*^{[p]}(IA_n^{[p]})$?*

Answering to [HM17, rk. 8.6], we show that these two filtrations fit in the same kind of nice machinery as their classical counterparts, but they turn out to always differ. The paper ends on a quantification of the lack of stable surjectivity in this *p*-restricted case (see Proposition 3.20).

Acknowledgements: This work is part of the author’s PhD thesis. The author is indebted to his advisors, Antoine Touzé and Aurélien Djament, for their constant support, countless helpful discussions, and numerous remarks and comments on earlier versions of the present paper. He also thanks Gwenaél Massuyeau and Takao Satoh, who kindly agreed to be the reviewers of his thesis, for their useful observations and comments.

Contents

Introduction	1
1 Generalities on strongly central series	5
1.1 Notations and reminders	5
1.2 Strongly central filtrations and Lie algebras	6
1.3 Actions in the category of strongly central filtrations	7
1.4 Johnson’s morphisms	11
1.5 The Andreadakis problem	12
1.6 Lazard’s theorem	14
1.7 Congruence groups	15
1.8 Comparison between filtrations obtained from an action	17
2 Traces and stable surjectivity	19
2.1 Free differential calculus	19
2.2 Derivations and strongly central filtrations	21
2.3 Algebras, actions and derivations	22
2.4 Traces	25
2.4.1 The induced map between Lie algebras	26
2.4.2 Introducing the contraction map	27

2.5	Stable surjectivity	28
2.5.1	Vanishing of the trace map	28
2.5.2	Linear algebra	30
2.5.3	Stable cokernel of τ' and stable surjectivity	30
2.6	Automorphisms of free nilpotent groups	31
3	The case of positive characteristic	33
3.1	Dark's theorem	33
3.2	p -Restricted strongly central series	34
3.3	The p -restricted Andreadakis problem	35
3.4	The stable p -restricted Andreadakis problem	37
3.4.1	Vanishing of the trace map	37
3.4.2	Linear algebra	38
3.4.3	Stable cokernel of i_*	39

1 Generalities on strongly central series

1.1 Notations and reminders

Throughout the paper, G will denote an arbitrary group, and \mathbb{k} a commutative unitary ring. The left and right action of G on itself by conjugation are denoted respectively by $x^y = y^{-1}xy$ and ${}^yx = yxy^{-1}$. The *commutator* of two elements x and y in G is denoted by:

$$[x, y] := xyx^{-1}y^{-1}.$$

If A and B are subsets of G , we denote by $[A, B]$ the subgroup generated by the commutators $[a, b]$ with $(a, b) \in A \times B$. If A and B are stable by conjugation by elements of G (resp. by all automorphisms of G), then $[A, B]$ is a normal (resp. characteristic) subgroup of G . For instance, $[G, G]$ is a characteristic subgroup of G , called the *derived subgroup* of G . The quotient $G^{ab} := G/[G, G]$ is the *abelianization* of G , its bigger abelian quotient. The derived subgroup is the second step of a filtration of G by characteristic subgroups:

Definition 1.1. The *lower central series* of G , denoted by $\Gamma_*(G)$, or shortly Γ_* , is the filtration of G defined by:

$$\begin{cases} \Gamma_1 := G, \\ \Gamma_{k+1} := [G, \Gamma_k]. \end{cases}$$

Definition 1.2. A group G is said to be *nilpotent* if its lower central series stops. The least integer c such that $\Gamma_{c+1}(G) = \{1\}$ is then G 's *nilpotency class*. More generally, G is said to be *residually nilpotent* if its lower central series is *separated*, i.e. if:

$$\bigcap_i \Gamma_i(G) = \{1\}.$$

One can easily check the following formulas:

Proposition 1.3. For all $x, y, z \in G$,

- $[x, x] = 1$,
- $[x, y]^{-1} = [y, x]$,
- $[x, yz] = [x, y] ({}^y[x, z])$,

- $[[x, y], {}^y z] \cdot [[y, z], {}^z x] \cdot [[z, x], {}^x y] = 1,$
- $[[x, y^{-1}], z^{-1}]^x \cdot [[z, x^{-1}], y^{-1}]^z \cdot [[y, z^{-1}], x^{-1}]^y = 1.$

The last ones are two versions of the *Witt-Hall identity*, which implies the following:

Lemma 1.4 (3-subgroups lemma). *Let A, B and C be three subgroups of a group G . If two of the three following subgroups are trivial, then so is the third:*

$$[A, [B, C]], \quad [B, [C, A]], \quad [C, [A, B]].$$

Equivalently, one of them is contained in the normal closure of the two others.

1.2 Strongly central filtrations and Lie algebras

The theory of strongly central series has notably been studied by M. Lazard [Laz54].

Definition 1.5. Let G be a group. A *strongly central filtration* of G (also called *strongly central series* or *N -series*) is a filtration

$$G = G_1 \supseteq \cdots \supseteq G_i \supseteq \cdots$$

of G by subgroups, satisfying:

$$\forall i, j \geq 1, [G_i, G_j] \subseteq G_{i+j}.$$

Remark that indexation has to begin from $G = G_1$. In particular, $[G, N_i] \subseteq N_{i+1} \subseteq N_i$, which means exactly that the N_i are *normal* subgroups of G .

Proposition 1.6. *Let G be a group. The lower central series $\Gamma_*(G)$ is a strongly central series on G , and it is the minimal one.*

Proof. The strong centrality is shown by induction, using the 3-subgroup lemma 1.4. Given a strongly central filtration G_* of $G = G_1$, a straightforward induction then gives: $G_i \supseteq \Gamma_i(G)$ for any $i \geq 1$. \square

Let $G = G_1 \supseteq \cdots \supseteq G_k \supseteq \cdots$ be any strongly central filtration of a group G . The quotients $\mathcal{L}_i(N_*) := N_i/N_{i+1}$ are abelian (for any $i \geq 1$), since $[N_i, N_i] \subseteq N_{2i} \subseteq N_{i+1}$. The graded abelian group

$$\mathcal{L}(N_*) := \bigoplus_{i \geq 1} \mathcal{L}_i(N_*),$$

is endowed with a bracket induced by the commutator map $(x, y) \mapsto [x, y]$ of G . Using the formulas 1.3, one easily checks that this defines a Lie bracket: $\mathcal{L}(G_*)$ is a Lie ring (*i.e.* a Lie algebra over \mathbb{Z}).

Notation 1.7. We denote $\mathcal{L}(\Gamma_*(G))$ by $\mathcal{L}(G)$ (that is, if we do not specify a strongly central filtration on a group, it is understood to be filtered by its lower central series).

Example 1.8. If G is a free group, then $\mathcal{L}(G)$ is the free Lie algebra over the \mathbb{Z} -module G^{ab} [Laz54, th. 4.2].

As products of commutators become sums of brackets inside the Lie algebra, the following fundamental property follows from the definition of the lower central series:

Proposition 1.9. *The Lie ring $\mathcal{L}(G)$ is generated in degree 1. Precisely, it is generated (as a Lie ring) by $\mathcal{L}_1(G) = G^{ab}$. As a consequence, if G is of finite type, then each $\mathcal{L}_n(G)$ is too.*

1.3 Actions in the category of strongly central filtrations

Let \mathcal{SCF} be the category whose objects are the strongly central filtrations, where morphisms between G_* and H_* are the group morphisms from G_1 to H_1 preserving filtrations. There is a forgetful functor $\omega_1 : \mathcal{SCF} \rightarrow \mathcal{Grps}$ defined by $G_* \mapsto G_1$. This functor admits a left adjoint $\Gamma : G \mapsto \Gamma_*(G)$ (see Proposition 1.6). It also admits a right adjoint $G \mapsto (G, G, \dots)$.

Proposition 1.10. *The category \mathcal{SCF} is complete and cocomplete, and is homological (but not semi-abelian).*

A general reference on homological categories is [BB04]. The reader can also consult [HL11] for a simple version of the axioms defining homological cocomplete and semi-abelian categories.

Proof of Proposition 1.10. The forgetful functor ω_1 admits both a left and a right adjoint, so it has to commute to limits and colimits. It does not create either of them (in the sense of [ML98], V.1), but it almost does.

Precisely, let $F : \mathcal{D} \rightarrow \mathcal{SCF}$ be a diagram. The colimit G_∞ of the group diagram $\omega_1 F$ is in general endowed with several strongly central filtrations making $\omega_1 F(d) \rightarrow G_\infty$ into filtration-preserving morphisms (for instance the trivial one). One checks easily that the minimal such filtration (which is the intersection of all those) is the colimit of F .

Similarly, the limit G^∞ of the group diagram $\omega_1 F$ is endowed with several strongly central filtrations making $\varphi_d : G^\infty \rightarrow \omega_1 F(d)$ into filtration-preserving morphisms (for instance its lower central series). However, the maximal such filtration is the limit of F . It is explicitly described as:

$$G_*^\infty = \bigcap_d \varphi_d^{-1}(F(d)).$$

To check that \mathcal{SCF} is homological, one can check the axioms given in [HL11]. It is not semi-abelian, because there are equivalence relation $R_* \subseteq G_*^2$ for which R_* is not the induced filtration on R_1 (like in topological groups – or more generally in the categories of topological algebras considered in [BC05] – where an equivalence relation does not have to be endowed with the induced topology). \square

In a homological category, we need to distinguish between usual epimorphisms (resp. monomorphisms) and *regular* ones, that is, the ones obtained as coequalizers (resp. equalizers). In \mathcal{SCF} , the former are the u such that $u_1 = \omega_1(u)$ is an epimorphism (resp. a monomorphism), whereas the latter are *surjections* (resp. *injections*):

Definition 1.11. Let $u : G_* \rightarrow H_*$ be a morphism in \mathcal{SCF} . It is called an *injection* (resp. a *surjection*) when u_1 is injective (resp. surjective) and $u^{-1}(H_i) = G_i$ (resp. $u(G_i) = H_i$) for all i .

Examples of homological categories include abelian categories, the category \mathcal{Grps} of groups, or the category \mathcal{Lie} of Lie algebras. The usual lemmas of homological algebra (the nine lemma, the snake lemma, the five lemma...) are true in these categories. Homological categories differ from abelian ones notably by the fact that in general, two split extensions between the same objects are not isomorphic (by an isomorphism preserving the splittings). This allows us to define an action of an object on another.

Definition 1.12. Let \mathcal{C} be a homological category. If X and Z are two objects of \mathcal{C} , we define an *action* of Z on X as a split extension (with a given splitting):

$$X \longrightarrow Y \overset{\longleftarrow}{\rightleftarrows} Z.$$

When such an action is given, we will say that Z *acts on* X , and write: $Z \circlearrowleft X$.

This definition (which needs only the weaker setting of *pointed protomodular categories* to make sense) is motivated by the situation in $\mathcal{G}rps$, where an action of a group K on a group G is encoded by a semi-direct product structure $G \rtimes K$.

Remark 1.13. The choice of splitting is crucial here. For instance, the canonical extension:

$$X \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} X \times X \xrightarrow{\begin{pmatrix} 0 & 1 \end{pmatrix}} X$$

can be split by $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$, or by the diagonal $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$. The first choice gives the trivial action, whereas the second one gives the *adjoint action*, which is highly non-trivial: in $\mathcal{L}ie$, this gives the adjoint representation; in $\mathcal{G}rps$, we get the action of a group on itself by conjugation.

The set $\text{Act}(Z, X)$ of actions of Z on X is a contravariant functor in Z : the restriction of an action along a morphism is defined *via* a pullback. In $\mathcal{G}rps$, as in $\mathcal{L}ie$, this functor is representable, for any X . Indeed, an action of a group K on a group G is given by a morphism $K \rightarrow \text{Aut}(G)$. Similarly, an action of a Lie algebra \mathfrak{k} on a Lie algebra \mathfrak{g} is given by a morphism $\mathfrak{k} \rightarrow \text{Der}(\mathfrak{g})$, where $\text{Der}(\mathfrak{g})$ is the Lie algebra of *derivations* from \mathfrak{g} to itself. Recall that a derivation ∂ is a linear map satisfying:

$$\partial([x, y]) = [\partial x, y] + [x, \partial y].$$

The situation when actions are representable has notably been studied in [BJK05], and in several subsequent papers [BB07, Bou08]. The following terminology was introduced in [BB07, Def. 1.1]:

Definition 1.14. A homological category \mathcal{C} is said to be *action-representative* when the functor $\text{Act}(-, X)$ is representable, for any object $X \in \mathcal{C}$.

Our goal for the rest of this section is to construct universal actions in \mathcal{SCF} , getting in particular the following result:

Proposition 1.15. *The category \mathcal{SCF} is action-representative.*

A representative for $\text{Act}(-, X)$ is a universal action on X . Explicitly, it is an action of an object $\mathcal{A}(X)$ on X such that any action $Z \circlearrowleft X$ is obtained by restriction along a unique morphism $Z \rightarrow \mathcal{A}(X)$. For instance, in $\mathcal{G}rps$, the universal action on G is:

$$G \hookrightarrow G \rtimes \text{Aut}(G) \twoheadrightarrow \text{Aut}(G),$$

where the group $G \rtimes \text{Aut}(G)$ is the *holomorph* of G . Its underlying set is $G \times \text{Aut}(G)$, endowed with the product defined by $(g, \sigma) \cdot (h, \tau) := (g\sigma(h), \sigma\tau)$.

The construction of universal actions in \mathcal{SCF} is given by a theorem of Kaloujnine [Kal50a, Kal50b], quoted by Lazard in [Laz54, p. 117]:

Theorem 1.16 (Kaloujnine). *Let G_* be a strongly central series. Let $j \geq 1$ be an integer. Let $\mathcal{A}_j(G_*) \subseteq \text{Aut}(G_*)$ be the subgroup of automorphisms acting trivially on every quotient G_i/G_{i+j} . Then $\mathcal{A}_*(G_*)$ is a strongly central series.*

Notation 1.17. As for Lie rings (Notation 1.7) we abbreviate $\mathcal{A}_*(\Gamma_*(G))$ to $\mathcal{A}_*(G)$.

We can rewrite the definition of $\mathcal{A}_j(G_*)$ given in the theorem as:

$$\mathcal{A}_j(G_*) = \ker \left(\text{Aut}(G_*) \longrightarrow \prod_i \text{Aut}(G_i/G_{i+j}) \right).$$

Identifying G and $\text{Aut}(G)$ to the subgroups $G \times 1$ and $1 \times \text{Aut}(G)$ of the holomorph $G \rtimes \text{Aut}(G)$, we can define the commutator of an automorphism with an element of G :

$$[\sigma, g] = \sigma(g)g^{-1}.$$

Note that $[\text{Aut}(G), G] \subseteq G$. Using this point of view, we can rephrase the previous definition:

$$\mathcal{A}_j(G_*) = \{ \sigma \in \text{Aut}(G_*) \mid \forall i \geq 1, [\sigma, G_i] \subseteq G_{i+j} \} \subseteq G_1 \rtimes \text{Aut}(G_*). \quad (1.17.1)$$

Proof of Theorem 1.16. We abbreviate $\mathcal{A}_j(G_*)$ to \mathcal{A}_j . Obviously, $\mathcal{A}_{j+1} \subseteq \mathcal{A}_j$. We show the strong centrality using the 3-subgroup lemma (Lemma 1.4). Precisely, let $\alpha, \beta \geq 1$ be two integers. For all $i \geq 1$, the group $G_{i+\alpha+\beta}$ is normal in $G \rtimes \text{Aut}(G_*)$ (it is normal in G and $\text{Aut}(G_*)$ -stable). Lemma 1.4 thus implies:

$$[[\mathcal{A}_\alpha, \mathcal{A}_\beta], G_i] \subseteq G_{i+\alpha+\beta}.$$

This says exactly that $[\mathcal{A}_\alpha, \mathcal{A}_\beta] \subseteq \mathcal{A}_{\alpha+\beta}$, which is the desired conclusion. \square

Remark 1.18. The group $\mathcal{A}_1(G_*)$ is the group of automorphisms of G_1 preserving G_* and acting trivially on $\mathcal{L}(G_*)$.

Example 1.19. Let $G_* = \Gamma_*(G)$. Then $\mathcal{L}(G_*)$ is generated in degree one as a Lie algebra. As a consequence $\mathcal{A}_1(G_*)$ is the subgroup of automorphisms acting trivially on the abelianization $G^{ab} = \mathcal{L}_1(G)$, denoted by IA_G .

In order to show that the filtration $\mathcal{A}_*(G_*)$ acts universally on G_* , we need to investigate actions in \mathcal{SCF} .

Proposition 1.20. *An action $K_* \curvearrowright G_*$ in \mathcal{SFC} consists of a group action of $K = K_1$ on $G = G_1$ such that :*

$$\forall i, j, [K_i, G_j] \subseteq G_{i+j}.$$

Proof. Let an action of K_* on G_* be given:

$$G_* \hookrightarrow H_* \overset{\leftarrow}{\twoheadrightarrow} K_*.$$

The forgetful functor $\omega_1 : G_* \mapsto G_1$ from \mathcal{SCF} to groups has a left adjoint $G \mapsto \Gamma_*G$. Hence, it commutes with kernels. Applying this functor to the given action, we get a split extension of groups:

$$G_1 \hookrightarrow H_1 \overset{\leftarrow}{\twoheadrightarrow} K_1.$$

The group H_1 thus decomposes as a semi-direct product $G_1 \rtimes K_1$ of G_1 by K_1 .

In fact, there are other forgetful functors $\omega_i : G_* \mapsto G_i$, each one with a left adjoint $G \mapsto \Gamma_{\lceil \frac{\cdot}{i} \rceil} G$ (where $\lceil - \rceil$ is the usual ceiling function). We then get split extensions of groups:

$$G_i \hookrightarrow H_i \overset{\leftarrow}{\twoheadrightarrow} K_i.$$

The groups H_i thus decomposes as semi-direct products $G_i \rtimes K_i$ of G_i by K_i .

As H_* is a strongly central filtration, we can apply Lemma 1.21 below to get the desired relation. Conversely, let a group action be given as in the statement of the proposition. Using the same lemma, we see that $H_* = G_* \rtimes K_*$ is a strongly central filtration on $H = K \rtimes G$, and the corresponding split sequence in \mathcal{SCF} is exact. \square

Lemma 1.21. *Let $K \curvearrowright G$ be an action in \mathcal{Grps} , encoded in a semi-direct product structure $H = G \rtimes K$. Let G_* and K_* be given filtrations on $G = G_1$ and $K = K_1$ respectively. Then the $H_i := G_i \rtimes K_i$ are subgroups of H defining a strongly central filtration of H if and only if:*

$$\begin{cases} K_* \text{ is a strongly central series on } K, \\ G_* \text{ is a strongly central series on } G, \\ \forall i, j, [K_j, G_i] \subseteq G_{i+j}. \end{cases}$$

Proof. Suppose first that $(G_i \rtimes K_i)_i$ is strongly central. Then its projection K_* on K also is. So is its intersection G_* with G . Hence, the conclusion follows from:

$$[K_j, G_i] \subseteq (G_{i+j} \rtimes K_{i+j}) \cap G = G_{i+j}.$$

Conversely, under the hypothesis listed above, G_i is stable under the action of K_i , so the $H_i = G_i \rtimes K_i$ are subgroups of H . We then use the formulas 1.3 to compute $[kg, k'g']$ with k, g, k', g' in K_i, G_i, K_j and G_j respectively:

$$\begin{aligned} [kg, k'g'] &= [kg, k'] \cdot^{g'} [kg, g'] \\ &= {}^k [g, k'] \cdot [k, k'] \cdot {}^{k'k} [g, g'] \cdot {}^{k'} [k, g'] \\ &\in G_{i+j} \cdot K_{i+j} \cdot G_{i+j} \cdot G_{i+j} = G_{i+j} \rtimes K_{i+j}. \end{aligned}$$

This completes the proof. \square

We are now ready to show the result announced in Proposition 1.15:

Proposition 1.22. *Let G_* be a strongly central series. The strongly central series $\mathcal{A}_*(G_*)$ acts canonically on G_* , and this action is universal.*

Proof. That $\mathcal{A}_*(G_*)$ acts on G_* follows from the formula (1.17.1), Theorem 1.16 and Proposition 1.20.

Given an action of a strongly central series K_* on G_* , the underlying group action is described by a unique morphism from K_1 to $\text{Aut}(G_1)$. From Proposition 1.20, we deduce that this morphism sends K_j into $\mathcal{A}_j(G_*)$. Conversely, any morphism from K_* to $\mathcal{A}_*(G_*)$ in \mathcal{SCF} gives a group action lifting to an action in \mathcal{SCF} by Proposition 1.20. \square

Remark 1.23. If a group K acts on a group G , and G_* is a strongly central filtration on $G = G_1$, we can pull back the canonical filtration $\mathcal{A}_*(G_*)$ by the associated morphism:

$$K \longrightarrow \text{Aut}(G).$$

This gives a strongly central filtration $\mathcal{A}_*(K, G_*)$, maximal amongst strongly central filtrations on subgroups of K which act on G_* via the given action $K \circlearrowright G$. It can be described explicitly as:

$$\mathcal{A}_j(K, G_*) = \{ k \in K \mid \forall i \geq 1, [k, G_i] \subseteq G_{i+j} \} \subseteq K.$$

1.4 Johnson's morphisms

The construction of the Johnson morphism associated with an action in \mathcal{SCF} relies on the following:

Proposition 1.24. *The Lie functor $\mathcal{L} : \mathcal{SCF} \rightarrow \text{Lie}_{\mathbb{Z}}$ is exact, i.e. it preserves short exact sequences [BB04, Def 4.1.5].*

Proof. The exactness of \mathcal{L} is equivalent to the exactness of each $\mathcal{L}_i : \mathcal{SCF} \rightarrow \text{Ab}$. Consider the forgetful functors $\omega_i : G_* \mapsto G_i$ from \mathcal{SCF} to \mathcal{Grps} . Each ω_i has a left adjoint $G \mapsto \Gamma_{[\frac{*}{i}]}G$, so these functors preserve kernels. Moreover, they preserve regular epimorphisms, which are surjections in \mathcal{SCF} (cf. Definition 1.11). Hence, they are exact. Since \mathcal{L}_i is the cokernel of the injection $\omega_{i+1} \hookrightarrow \omega_i$, its exactness follows from the nine lemma in \mathcal{Grpes} . Precisely, if \mathcal{E} is any short exact sequence in \mathcal{SCF} , apply the nine lemma to the diagram $\omega_{i+1}(\mathcal{E}) \hookrightarrow \omega_i(\mathcal{E}) \rightarrow \mathcal{L}_i(\mathcal{E})$ to get the exactness of $\mathcal{L}_i(\mathcal{E})$ from that of $\omega_i(\mathcal{E})$ and $\omega_{i+1}(\mathcal{E})$. \square

As a consequence of Proposition 1.24, the functor \mathcal{L} preserves actions. Precisely, from an action in \mathcal{SCF} :

$$G_* \hookrightarrow H_* \overset{\curvearrowright}{\twoheadrightarrow} K_*,$$

we get an action in the category of graded Lie rings:

$$\mathcal{L}(G_*) \hookrightarrow \mathcal{L}(H_*) \overset{\curvearrowright}{\twoheadrightarrow} \mathcal{L}(K_*).$$

Such an action is given by a morphism of graded Lie rings:

$$\mathcal{L}(K_*) \longrightarrow \text{Der}_*(\mathcal{L}(G_*)). \quad (1.24.1)$$

The target is the (graded) Lie algebra of graded derivations: a derivation is of degree k when it raises degrees of homogeneous elements by k .

Definition 1.25. The morphism (1.24.1) is called the *Johnson morphism* associated to the given action $K_* \circlearrowright G_*$.

We can give an explicit description of this morphism: for $k \in K$, the derivation associated to \bar{k} is induced by $[\bar{k}, -]$ inside $\mathcal{L}(G_* \rtimes K_*) = \mathcal{L}(G_*) \rtimes \mathcal{L}(K_*)$, so it is induced by $[k, -]$ inside $G_* \rtimes K_*$.

Example 1.26. The Johnson morphism associated to the universal action $\mathcal{A}_*(G_*) \circlearrowright G_*$ is the Lie morphism:

$$\tau : \mathcal{L}(\mathcal{A}_*(G_*)) \longrightarrow \text{Der}_*(\mathcal{L}(G_*))$$

induced by $\sigma \mapsto (x \mapsto \sigma(x)x^{-1})$.

If K_1 is in fact a normal subgroup of a group K_0 , such that each K_i is normal in K_0 , and such that the action of K_1 on G_* can be extended to an action of K_0 , then K_0 acts on $\mathcal{L}(G_* \rtimes K_*)$ and this action factorizes through K_0/K_1 . Moreover, as this action is by automorphisms of the Lie ring:

Lemma 1.27. *Let $K_0 \triangleright K_1$ be given as above. Then the Johnson morphism τ is K_0/K_1 -equivariant.*

The action of K_0 on derivations is by conjugation. Precisely:

$$\tau(k \cdot x) = [k \cdot x, -] = k \cdot [x, k^{-1} \cdot -] = k \circ \tau(x) \circ k^{-1}.$$

Lemma 1.28. *Let $K_* \curvearrowright H_*$ be an action in \mathcal{SCF} . The associated Johnson morphism $\tau : \mathcal{L}(K_*) \rightarrow \text{Der}(\mathcal{L}(G_*))$ is injective if and only if K_* is the filtration $\mathcal{A}_*(K, G_*)$ defined in Remark 1.23.*

Proof. Every non-trivial element in $\ker(\tau_j)$ lifts to an element in $\mathcal{A}_j(K, G_*) - K_j$. Conversely, an element σ in $\mathcal{A}_j(K, G_*) - K_j$ is in $K_k - K_{k+1}$ for some $k < j$. Then $\bar{\sigma} \in K_k/K_{k+1} - \{0\}$ is a non-trivial element in $\ker(\tau)$. \square

Remark 1.29. The definition of $\mathcal{A}_j(K, G_*)$ makes sense for $j = 0$, giving a subgroup $K_0 = \mathcal{A}_0(K, G_*)$ of K acting as above. The morphism τ is then $\mathcal{A}_0(K, G_*)$ -equivariant. In fact, τ can then be extended to a morphism of *extended Lie algebras*, in the sense of [HM17]. Indeed, their construction of an algebra of *extended derivations* is exactly a construction of universal actions in the category of extended Lie algebras, and their version of the Johnson morphism is exactly the one we find if we replace N -series and Lie algebras by their extended version in the constructions above.

1.5 The Andreadakis problem

Let G be a group. To study the structure of $\text{Aut}(G)$, we can consider first how automorphisms act on G^{ab} . Then we can put aside this linear part by considering the kernel of the projection from $\text{Aut}(G)$ to $GL(G^{ab})$. This kernel IA_G is (residually) nilpotent when G is, and is endowed with two strongly central filtrations: its lower central series, and the *Andreadakis filtration* $\mathcal{A}_*(G)$. We are thus led to the problem of comparing these filtrations, which we call the *Andreadakis problem* (Problem 1).

Recall from Theorem 1.16 and Example 1.19 that $\mathcal{A}_*(G)$ is a strongly central filtration on $\mathcal{A}_1(G) = IA_G$. Since $\mathcal{A}_1(G)$ is the kernel of the canonical action of $\text{Aut}(G)$ on $\mathcal{L}(G)$, we get an induced faithful action of $\text{Aut}(G)/IA_G (\leq GL(G^{ab}))$ on $\mathcal{L}(G)$. The next lemma gives a similar concrete description of all the $\mathcal{A}_j(G)$:

Lemma 1.30. *The group $\mathcal{A}_j(G)$ is the subgroup of automorphisms acting trivially on $G/\Gamma_{j+1}(G)$, i.e. $\mathcal{A}_j(G) = \{ \sigma \in \text{Aut}(G) \mid [\sigma, G] \subseteq \Gamma_{j+1}(G) \}$.*

Proof. Let us denote by K_j the right-hand side of the equality. We only need to show the inclusion $K_j \subseteq \mathcal{A}_j(G)$. The case $j = 1$ is given in Example 1.19. Suppose it true for $j - 1$, and let $\sigma \in K_j$. In particular, σ is inside $K_{j-1} = \mathcal{A}_{j-1}(G)$, so it induces a derivation $[\sigma, -]$ of degree $j - 1$ of $\mathcal{L}(G)$, via the Johnson morphism. By definition, $[\sigma, G] \subseteq \Gamma_{j+1}$. This says exactly that the derivation $[\sigma, -]$ is trivial on $\mathcal{L}_1(G)$. Since $\mathcal{L}(G)$ is generated in degree one (Proposition 1.9), $[\sigma, -]$ has to be trivial on all of $\mathcal{L}(G)$: for all i , $[\sigma, \Gamma_i] \subseteq \Gamma_{i+j}$, which is the desired conclusion. \square

The filtration $\mathcal{A}_*(G)$ is strongly central on $\mathcal{A}_1(G) = IA_G$. As a consequence, $\mathcal{A}_k(G)$ contains $\Gamma_k(IA_G)$. A first consequence of this inclusion is that the (residual) nilpotency of G implies the (residual) nilpotency of IA_G . Precisely, let G be a c -nilpotent group. Then $\mathcal{A}_c(G) = \{1_G\}$. Accordingly, $\Gamma_c(IA_G) = \{1_G\}$, so IA_G is $(c-1)$ -nilpotent. In a similar fashion, one can check that IA_G has to be residually nilpotent when G is.

The following question is crucial for trying to understand the structure of automorphism groups of residually nilpotent groups, in particular for trying to understand the structure of $\text{Aut}(F_n)$:

Problem 1 (Andreadakis). *What is the difference between $\mathcal{A}_*(G)$ and $\Gamma_*(IA_G)$?*

Example 1.31. Consider the alternating group A_n . When $n \neq 2, 6$, $\text{Aut}(A_n) = \Sigma_n$ (acting by conjugation), as is easily deduced from [Rot95, cor. 7.5]. But if $n \geq 5$, then A_n is perfect: $[A_n, A_n] = A_n$. On the one hand, the Andreadakis filtration is thus constant equal to $IA(A_n) = \text{Aut}(A_n) = \Sigma_n$. On the other hand, the lower central series of Σ_n is $\Sigma_n, A_n, A_n, \dots$, so the two filtrations differ in this case.

Recall from the Introduction that we are interested in a stable form of the problem for $G = F_n$:

Problem 2 (Andreadakis - stable version). *What is the difference between $\mathcal{A}_k(F_n)$ and $\Gamma_k(IA_n)$ for $n \gg k$?*

The *Johnson morphism* turns out to be a powerful tool in the study of the Andreadakis filtration (which is the analogous of *Johnson's filtration* on the mapping class group, defined by Johnson in [Joh83]).

Recall the $\mathcal{A}_*(G)$ acts on $\Gamma_*(G)$ (see Proposition 1.22). The associated Johnson morphism τ is described in Example 1.26. The $\Gamma_i(G)$ are characteristic subgroups of G , so $\mathcal{A}_0(G) = \text{Aut}(G)$. Lemmas 1.27 and 1.28 then give:

Lemma 1.32. *The morphism τ is injective and $\text{Aut}(G)/IA_G$ -equivariant.*

The filtration $\Gamma_*(IA_G)$ also acts on $\Gamma_*(G)$, by pulling back the universal action along the inclusion $i : \Gamma_*(IA_G) \subseteq \mathcal{A}_*(F_n)$. The associated Johnson morphism is denoted by: $\tau' = \tau \circ i_*$. The morphism τ' is still equivariant, but is injective if and only if G satisfies Andreadakis' conjecture (*i.e.* if $i_* = 1$).

Example 1.33. If G is a free group, then $\mathcal{L}(F_n)$ is the free Lie algebra $\mathfrak{L}V$ on $V = G^{ab}$. If $\mathfrak{g} \curvearrowright \mathfrak{h}$ is an action in *Lie*, then \mathfrak{g} also acts on \mathfrak{h}^o , which is obtained by considering the module \mathfrak{h} as an abelian Lie algebra. Derivations from \mathfrak{g} to \mathfrak{h} then identify with sections of the projection $\mathfrak{h}^o \rtimes \mathfrak{g} \rightarrow \mathfrak{g}$. Hence, the free Lie algebra is also free with respect to derivations. In particular:

$$\text{Der}_k(\mathfrak{L}V) \cong \text{Hom}_{\mathbb{k}}(V, \mathfrak{L}_k V) \cong V^* \otimes \mathfrak{L}_k V.$$

The following result is well-known [Kaw06, th. 6.1]:

Proposition 1.34. *In degree one, the Johnson morphism τ' is a $\text{GL}_n(\mathbb{Z})$ -equivariant isomorphism:*

$$\tau'_1 : IA_n^{ab} \cong V^* \otimes \Lambda^2 V.$$

Proof. The group IA_n is generated by the following elements [Nie24] – see also [BBM07, 5.6]:

$$K_{ij} : x_t \mapsto \begin{cases} x_j x_i x_j^{-1} & \text{if } t = i \\ x_t & \text{else} \end{cases} \quad \text{and} \quad K_{ijk} : x_t \mapsto \begin{cases} [x_j, x_k] x_i & \text{if } t = i \\ x_t & \text{else.} \end{cases} \quad (1.34.1)$$

One can check by a direct calculation that these generators are sent to a basis of the free abelian group $V^* \otimes \Lambda^2 V$. \square

1.6 Lazard's theorem

Definition 1.35. A *filtered algebra* A_* is an associative \mathbb{k} -algebra A_0 endowed with a filtration by ideals: $A = A_0 \supseteq A_1 \supseteq \dots$ such that: $\forall i, j, A_i A_j \subseteq A_{i+j}$. We denote by $fAlg$ the category of filtered algebras (and filtration-preserving morphisms).

Example 1.36. Let $\mathbb{k}G$ be the *group algebra* of G with coefficients in the (commutative) ring \mathbb{k} . We denote by $\varepsilon : \mathbb{k}G \rightarrow \mathbb{k}$ its canonical *augmentation* and by $I_{\mathbb{k}}G := \ker(\varepsilon)$ its *augmentation ideal* (we will sometimes write IG , or even I , for short). Then $\mathbb{k}G$ is filtered by the powers I^*G of its augmentation ideal. If G is a free group, then $\text{gr}(\mathbb{k}G)$ is the tensor algebra over $G^{ab} \otimes \mathbb{k}$ [Pas79, th. 6.2].

From Theorem 1.16, we can deduce the useful corollary [Laz54, th. 3.1]:

Theorem 1.37 (Lazard). *Let $A = A_0 \supseteq A_1 \supseteq \dots$ be a filtered algebra. Then $A_*^\times := A^\times \cap (1 + A_*)$ is a strongly central filtration on $A_1^\times \subseteq A^\times$, and $(-) - 1$ induces an embedding of graded Lie algebras:*

$$\mathcal{L}(A_*^\times) \hookrightarrow \text{gr}_*(A_1).$$

Remark 1.38. Fix a morphism $G \xrightarrow{\alpha} A^\times$. We can pull back the filtration given by the theorem to get a strongly central filtration $\alpha^{-1}(1 + A_*)$ on $G_1 = \alpha^{-1}(1 + A_1)$.

Proof of Theorem 1.37. The filtration $A = A_0 \supseteq A_1 \supseteq \dots$ can be seen as a filtration of the abelian group A . As A is abelian, it has to be strongly central. Consider the action by left multiplication:

$$\rho : A^\times \longrightarrow \text{Aut}(A, +).$$

Let $a \in A^\times$. One can easily check that $\rho(a) \in \mathcal{A}_j(A_*)$ (where $\mathcal{A}_j(A_*)$ is the filtration defined in Kaloujnine's theorem 1.16) if and only if $a \in 1 + A_j$. Thus, $A^\times \cap (1 + A_i) = \rho^{-1}(\mathcal{A}_j(A_*))$ is a strongly central filtration, as announced. It remains to show that $\partial = \alpha - 1$ induces a morphism of Lie ring (necessarily injective). This can be checked directly from the formula:

$$[g, h] - 1 = [g - 1, h - 1]g^{-1}h^{-1}.$$

We will give a slightly different proof later on, using the concept of derivations (see Paragraph 2.2). \square

Example 1.39. Applying Remark 1.38 to the inclusion of G in $\mathbb{k}G$ filtered by the powers of the augmentation algebra, we get the *dimension series* of G :

$$D_*^{\mathbb{k}}G = G \cap (1 + I_{\mathbb{k}}^*G).$$

It is a strongly central series on G , so it contains Γ_*G . The question of the equality of $D_*^{\mathbb{Z}}G$ and Γ_*G was known as the *dimension subgroup problem* during a long time, until an example of a group for which the two filtration differ was given in [Rip72]. See [MP09, chap. 2] for more on this subject.

If G is a free group, then $\mathcal{L}(D_*^{\mathbb{Z}}G)$ is the sub-Lie ring generated in degree one in the tensor algebra $\text{gr}(\mathbb{Z}G) \cong TV$. Hence (by the PBW theorem), it is the free Lie ring. It then has to coincide with $\mathcal{L}(G)$, so $D_*^{\mathbb{Z}}G = \Gamma_*G$. Thus, free groups have the *dimension property*.

Lazard's theorem gives a construction of a strongly central filtration from a filtered algebra. Conversely, we can define a filtered algebra from a strongly central filtration G_* on $G = G_1$. Indeed, let $\mathbb{k}G$ be filtered by:

$$F_i := \mathbb{k}G \cdot (N_i - 1) = \ker(\mathbb{k}G \longrightarrow \mathbb{k}(G/H_i)).$$

This filtration does not make $\mathbb{k}G$ into a filtered algebra, but it generates a filtration which does:

$$\mathfrak{a}_j^{\mathbb{k}}(N_*) := \sum_{i_1 + \dots + i_n \geq j} \mathbb{k}G \cdot (N_{i_1} - 1) \cdots (N_{i_n} - 1).$$

One can easily check that these constructions are universal:

Proposition 1.40. *The above constructions define an adjunction:*

$$SCF \begin{array}{c} \xrightarrow{\mathfrak{a}_*^{\mathbb{k}}} \\ \xleftarrow{(-)_*^{\times}} \end{array} fAlg_{\mathbb{k}}.$$

1.7 Congruence groups

If I is an (associative) ring without unit, recall that its *congruence group* $GL_n(I)$ is defined as:

$$GL_n(I) := \ker(GL_n(A) \longrightarrow GL_n(A/I)),$$

where A is any unitary (associative) ring containing I as a (two-sided) ideal, (e.g. $A = I \rtimes \mathbb{Z}$). This group depends only on I , as it is exactly $(1 + M_n(I))^{\times}$.

If $A = A_0 \supseteq A_1 \supseteq \dots$ is a filtered algebra (see Definition 1.35), then so is the matrix algebra $M_n(A)$, endowed with the filtration $M_n(A_*)$. Theorem 1.37 gives us a strongly central filtration of the congruence group:

$$GL_n(A_1) = GL_n(A) \cap (1 + M_n(A_1)) = \ker(GL_n(A) \longrightarrow GL_n(A/A_1)).$$

by congruence subgroups:

$$GL_n(A_j) = GL_n(A) \cap (1 + M_n(A_j)) = \ker(GL_n(A) \longrightarrow GL_n(A/A_j)),$$

and an embedding of the associated Lie ring into a matrix algebra:

$$\mathcal{L}(GL_n(A_*)) \hookrightarrow \text{gr}_*(M_n(A_*)) \cong M_n(\text{gr}_*(A_*)).$$

As in the proof of Theorem 1.37, this filtration can be interpreted as:

$$GL_n(A_*) = \mathcal{A}_*(GL_n(A), A_*^n).$$

We also can recover Lazard's theorem as the case $n = 1$ of this construction.

Suppose that A_* is commutative. Then the usual determinant defines a filtration-preserving morphism:

$$\det : GL_n(A_*) \longrightarrow GL_1(A_*) = A_*^{\times}.$$

Indeed, if $M \in M_n(A_j)$, then $\det(1+M) \in 1+A_j$. The following proposition determines the associated graded morphism:

Proposition 1.41. *The following square commutes:*

$$\begin{array}{ccc} \mathcal{L}(GL_n(A_*)) & \xrightarrow{(-)-1} & M_n(\text{gr}(A_*)) \\ \downarrow \det & & \downarrow \text{Tr} \\ \mathcal{L}(A_*^\times) & \xrightarrow{(-)-1} & \text{gr}(A_*). \end{array}$$

Moreover, it is Cartesian, that is:

$$\mathcal{L}(GL_n(A_*)) \cong \text{Tr}^{-1}(\mathcal{L}(A_*^\times) - 1).$$

The kernels of \det and Tr then coincide. We thus recover (an generalize slightly) a result of [Lop14]:

Corollary 1.42. *Let $SL_n(A_*)$ be the kernel of the determinant. Then:*

$$\mathcal{L}(SL_n(A_*)) \cong \ker(\text{Tr}) = \mathfrak{sl}_n(\text{gr}(A_*)).$$

Remark 1.43. If $\mathcal{L}(A_*^\times) = 0$, then $\mathcal{L}(GL_n(A_*)) = \mathcal{L}(SL_n(A_*))$, and this Lie ring identifies to $\mathfrak{sl}_n(\text{gr}(A_*))$. This happens for example when $GL_1(A_1) = \{1\}$ (implying $SL_n(A_*) = GL_n(A_*)$), which is verified for $A_* = q^*\mathbb{Z}$ (if $q > 2$), or $A_* = t^*\mathbb{k}[t]$.

Proof of Proposition 1.41. Let $M \in M_n(A_j)$. Then:

$$\det(1 + M) \equiv 1 + \text{Tr}(M) \pmod{A_j^2}.$$

When $j \geq 1$, then $A_j^2 \subseteq A_{j+1}$, so this formula gives the commutativity of the above square. The module $\text{Tr}^{-1}(\mathcal{L}_j(A_*^\times) - 1)$ is additively generated by the matrices:

$$\bar{a}\mathbf{e}_{\alpha\beta}, \bar{a}(\mathbf{e}_{11} - \mathbf{e}_{\alpha\alpha}) \text{ and } \bar{b}\mathbf{e}_{11}, \text{ for } \alpha \neq \beta, a \in A_j \text{ and } 1 + b \in A_j^\times.$$

Replacing $\bar{a}(\mathbf{e}_{11} - \mathbf{e}_{\alpha\alpha})$ by $\bar{a}(\mathbf{e}_{11} + \mathbf{e}_{1\alpha} - \mathbf{e}_{\alpha 1} - \mathbf{e}_{\alpha\alpha})$, we can lift these to $GL_n(A_j)$ as follows:

$$\begin{cases} 1 + a\mathbf{e}_{\alpha\beta} & \text{lifts } \bar{a}\mathbf{e}_{\alpha\beta}, \\ 1 + a(\mathbf{e}_{11} + \mathbf{e}_{1\alpha} - \mathbf{e}_{\alpha 1} - \mathbf{e}_{\alpha\alpha}) & \text{lifts } \bar{a}(\mathbf{e}_{11} + \mathbf{e}_{1\alpha} - \mathbf{e}_{\alpha 1} - \mathbf{e}_{\alpha\alpha}), \\ 1 + b\mathbf{e}_{11} & \text{lifts } \bar{b}\mathbf{e}_{11}. \end{cases}$$

This completes the proof. \square

Let $n \geq 3$. If A is a "non-totally-imaginary Dedekind ring of arithmetic type" and \mathfrak{q} is an ideal of A , then we have [BMS67, cor. 4.3 (b)] that $SL_n(\mathfrak{q})$ is normally generated in $SL_n(A)$ (in fact in $E_n(A)$) by the shear mappings $1 + t\mathbf{e}_{\alpha\beta}$ with $\alpha \neq \beta$ and $t \in \mathfrak{q}$. This applies for instance to $A = \mathbb{Z}$ and $\mathfrak{q} = (q)$. From this we deduce:

Proposition 1.44. *If $n \geq 5$, under the above hypothesis:*

$$\Gamma_*(SL_n(\mathfrak{q})) = SL_n(\mathfrak{q}^*).$$

Proof. The filtration $SL_n(\mathfrak{q}^*)$ is strongly central on $SL_n(\mathfrak{q})$, so it contains its lower central series. Conversely, using that:

$$\begin{cases} 1 + (a + b)\mathbf{e}_{\alpha\beta} = (1 + a\mathbf{e}_{\alpha\beta})(1 + b\mathbf{e}_{\alpha\beta}) & \text{if } \alpha \neq \beta, \\ 1 + ab\mathbf{e}_{\alpha\beta} = [1 + a\mathbf{e}_{\alpha\gamma}, 1 + b\mathbf{e}_{\gamma\beta}] & \text{if } \alpha, \beta \text{ and } \gamma \text{ are pairwise distinct,} \end{cases}$$

one can easily check that for any t in \mathfrak{q}^k and any $\alpha \neq \beta$, if $n \geq 5$:

$$1 + t\mathbf{e}_{\alpha\beta} \in \Gamma_k(SL_n(\mathfrak{q})).$$

Using the result from [BMS67], we see that these generate $SL_n(\mathfrak{q}^k)$ as a normal subgroup of $SL_n(A)$. Hence $SL_n(\mathfrak{q}^k) \subseteq \Gamma_k(SL_n(\mathfrak{q}))$, as required. \square

Remark 1.45 (On the q -torsion Andreadakis problem for \mathbb{Z}^n). Fix an integer q . A q -torsion strongly central filtration is a strongly central filtration G_* such that $G_i^q \subseteq G_{i+1}$ for every $i \geq 1$ (this means exactly that $q\mathcal{L}(G_*) = 0$). On any group G , there is a minimal q -torsion strongly central filtration $\Gamma_*^{(q)}(G)$. Moreover, $\mathcal{A}_*(G_*)$ is q -torsion whenever G_* is, because $\mathcal{L}(\mathcal{A}_*(G_*))$ embeds into a q -torsion Lie algebra *via* the Johnson morphism. Let us denote $\mathcal{A}_*(\Gamma_*^{(q)}(G))$ by $\mathcal{A}_*^{(q)}(G)$ and $\mathcal{A}_1^{(q)}(G)$ by $IA^{(q)}(G)$ (it is the group of automorphisms acting trivially on $G^{ab} \otimes (\mathbb{Z}/q)$). Thus we get an inclusion of $\Gamma_*^{(q)}(IA^{(q)})$ into $\mathcal{A}_*^{(q)}$ and a corresponding q -torsion Andreadakis problem.

Apply this with $G = \mathbb{Z}^n$ and $q \geq 3$. Then $\Gamma_*^{(q)}(G) = q^*\mathbb{Z}$, $\text{Aut}(G) = GL_n(\mathbb{Z})$, and $\mathcal{A}_*^{(q)}(G) = GL_n(q^*\mathbb{Z}) = SL_n(q^*\mathbb{Z})$ Proposition 1.44 then gives an answer to the q -torsion Andreadakis problem for \mathbb{Z}^n in the stable range ($n \geq 5$): $\mathcal{A}_*^{(q)}(\mathbb{Z}^n)$ is the lower central series of $IA^{(q)}(\mathbb{Z}^n) = SL_n(q\mathbb{Z})$.

When $A = \mathbb{Z}$ and $\mathfrak{q} = (q)$, the graded ring $\text{gr}(q^*\mathbb{Z})$ is $(\mathbb{Z}/q)[t]$. Moreover, $GL_n(q\mathbb{Z}) = SL_n(q\mathbb{Z})$ (see Remark 1.43). Proposition 1.44 and Corollary 1.42 thus give:

Corollary 1.46. *For all $n \geq 5$ and all $q \geq 3$, there is a canonical isomorphism of graded Lie rings (in degrees at least one):*

$$\mathcal{L}(GL_n(q\mathbb{Z})) \cong \mathfrak{sl}_n(\mathbb{Z}/q)[t],$$

where the degree of t is 1, and the Lie bracket of Mt^i and Nt^j is $[M, N]t^{i+j}$.

Remark 1.47. This generalizes [LS76, Th. 1.1], which is the degree-one part.

Remark 1.48. For $n = 2$ and $q \geq 5$ a prime number, the group $SL_2(q\mathbb{Z})$ is free on $1 + q(q^2 - 1)/12$ generators [Gro52, Fra33]. Its Lie ring is then a free Lie ring on the same number of generators. The author does not know a complete calculation for $n = 3$ or 4 : the above calculus does give the abelianization, but it fails to determine the whole lower central series.

1.8 Comparison between filtrations obtained from an action

Let G be a group. Suppose that G acts on two strongly central series H_* et K_* (by automorphisms in \mathcal{SCF}). We then can ask what link exists between $\mathcal{A}_*(G, H_*)$ and $\mathcal{A}_*(G, K_*)$ (as defined in Remark 1.23), depending on the links between H_* and K_* .

The next proposition describes the behaviour of the construction $\mathcal{A}_*(G, -)$ with respect to injections, surjections (see Definition 1.11) and semi-direct products in the category $G\text{-SCF}$ of strongly central series endowed with a G -action (where morphisms respect this action):

Proposition 1.49. *Let G be a group acting on strongly central series N_* , H_* and K_* . Let $u : N_* \rightarrow H_*$ and $v : H_* \rightarrow K_*$ be G -equivariant morphisms.*

If $u : N_ \rightarrow H_*$ is an injection, then $\mathcal{A}_*(G, H_*) \subseteq \mathcal{A}_*(G, N_*)$.*

If $v : H_ \rightarrow K_*$ is a surjection, then $\mathcal{A}_*(G, H_*) \subseteq \mathcal{A}_*(G, K_*)$.*

If $N_ \hookrightarrow H_* \overset{\leftarrow}{\twoheadrightarrow} K_*$ is a split exact sequence in $G\text{-SCF}$, then:*

$$\mathcal{A}_*(G, H_*) = \mathcal{A}_*(G, K_*) \cap \mathcal{A}_*(G, N_*).$$

Proof. In the first case, we identify $N = N_1$ to a subgroup of $H = H_1$. Let $g \in \mathcal{A}_j(G, H_*)$. We write:

$$[g, N_i] = [g, N \cap H_i] \subseteq [g, N] \cap [g, H_i] \subseteq N \cap H_{i+j} = N_{i+j}.$$

Similarly, to show the second assertion, let us take $g \in \mathcal{A}_j(G, H_*)$. We write:

$$[g, K_i] = [g, v(H_i)] = \varphi([g, H_i]) \subseteq v(H_{i+j}) = K_{i+j}.$$

The third assertion's hypothesis comes down to require that H_* decompose as a semi-direct product $K_* \rtimes N_*$, the action of G on H_* being factor-wise. We then get G -equivariant isomorphisms:

$$H_i/H_{i+j} \cong K_i/K_{i+j} \rtimes N_i/N_{i+j}.$$

An element g of G acts trivially on the left hand side if and only if it does on the right hand side. Whence the result. \square

Remark 1.50. Let $1 \rightarrow N_* \xrightarrow{u} H_* \xrightarrow{v} K_* \rightarrow 1$ be a non-split short exact sequence in $G - \mathcal{SCF}$, the first part of Proposition 1.49 gives:

$$\mathcal{A}_*(G, H_*) \subseteq \mathcal{A}_*(G, K_*) \cap \mathcal{A}_*(G, N_*).$$

Nevertheless, equality is not true in general. Indeed, the sequences:

$$1 \rightarrow N_i/N_{i+j} \xrightarrow{u} H_i/H_{i+j} \xrightarrow{v} K_i/K_{i+j} \rightarrow 1$$

are exact, but $g \in G$ can act trivially on the kernel and quotient without acting trivially on the middle term. For instance, $H^1(G) = \text{Ext}_G^1(\mathbb{Z}^{triv}, \mathbb{Z}^{triv})$ is non-trivial in general.

We can get a little more about semi-direct products:

Proposition 1.51. Let $N \hookrightarrow H \overset{\leftarrow}{\twoheadrightarrow} K$ be a split exact sequence in $G - \text{Grps}$. Suppose that N is filtered by a strongly central series N_* . Then:

$$\mathcal{A}_*(G, N_*) \subseteq \mathcal{A}_*(G, \mathcal{A}_*(K, N_*)).$$

Proof. Let us denote by K_* the filtration $\mathcal{A}_*(K, N_*)$ and by G_* the filtration $\mathcal{A}_*(G, N_*)$. A straightforward application of the 3-subgroup lemma (Lemma 1.4) in $(N \rtimes K) \rtimes G$ provides the inclusion:

$$[[G_\alpha, K_\beta], N_\gamma] \subseteq N_{\alpha+\beta+\gamma}.$$

Thus $[G_\alpha, K_\beta] \subseteq \mathcal{A}_{\alpha+\beta}(K, N_*) = K_{\alpha+\beta}$, which means that $G_\alpha \subseteq \mathcal{A}_\alpha(G, K_*)$. \square

Corollary 1.52. Let G be a group acting on a filtered algebra A_* by automorphisms of filtered algebras. Then:

$$\mathcal{A}_*(G, A_*) \subseteq \mathcal{A}_*(G, A_*^\times).$$

Proof. Apply Proposition 1.51 to the G -equivariant split exact sequence:

$$A \hookrightarrow A \rtimes A^\times \overset{\leftarrow}{\twoheadrightarrow} A^\times$$

and the given filtration on A . \square

Here is an interesting case when the filtrations of Corollary 1.52 are equal:

Proposition 1.53. *Let G be a group, and \mathbb{k} a commutative ring. Then:*

$$\mathcal{A}_*(\text{Aut}(G), D_*^{\mathbb{k}}G) = \mathcal{A}_*(\text{Aut}(G), I_{\mathbb{k}}^*G).$$

Proof. The algebra $\mathbb{Z}G$ is filtered by $I_{\mathbb{k}}^*G$, the powers of its augmentation ideal. The group $\text{Aut}(G)$ acts on $\mathbb{Z}G$ by automorphisms of filtered algebras. As $\mathcal{A}_*(G, I_{\mathbb{k}}^*G) = D_*^{\mathbb{k}}G$, Proposition 1.51, applied to the action of $\text{Aut}(G)$ on $\mathbb{k}G \rtimes G$, gives an inclusion:

$$\mathcal{A}_*(\text{Aut}(G), I_{\mathbb{k}}^*G) \subseteq \mathcal{A}_*(\text{Aut}(G), D_*^{\mathbb{k}}G).$$

To show that it is in fact an equality, take $\varphi \in \mathcal{A}_*(\text{Aut}(G), D_*^{\mathbb{k}}G)$. Then:

$$\forall g \in G = D_1G, \quad [\varphi, g] = (\varphi(g)g^{-1} - 1)g \in (D_{j+1} - 1)G \subseteq I^{j+1}.$$

We then show that $[\varphi, I^i] \subseteq I^{i+j}$ by induction on $i \geq 1$, using the formula:

$$[\varphi, uv] = [\varphi, u]\varphi(v) + u[\varphi, v] \quad (= (\varphi(u) - u) \cdot \varphi(v) + u \cdot (\varphi(v) - v)).$$

Let us remark that, in the language of paragraph 2.3, the last formula states that $[\varphi, -]$ is a $(1, \varphi)$ -derivation, so we have in fact used lemma 2.18. \square

2 Traces and stable surjectivity

2.1 Free differential calculus

We recall some basic concepts of free differential calculus. A detailed account can be found in [Fox53].

Definition 2.1. Let G be a group, and M a $\mathbb{k}G$ -module. A *derivation* from G to M is a map $\partial : G \rightarrow M$ such that:

$$\forall g, h \in G, \quad \partial(gh) = \partial g + g \cdot \partial h.$$

It can be extended to a linear map $\partial : \mathbb{k}G \rightarrow M$, which verifies:

$$\forall u, v \in \mathbb{k}G, \quad \partial(uv) = \partial(u)\varepsilon(v) + u \cdot \partial(v).$$

We denote by $\text{Der}(G, M)$ or $\text{Der}(\mathbb{k}G, M)$ the space of derivations from G to M . We will often write $\text{Der}(\mathbb{k}G)$ for $\text{Der}(\mathbb{k}G, \mathbb{k}G)$.

Remark 2.2. Let $G = F_S$ be the free group over a set S . As derivations identify with sections of $M \rtimes G \rightarrow G$, we get: $\text{Der}(\mathbb{k}F_S, M) \cong M^S$ for any $\mathbb{k}G$ -module M .

Definition 2.3. Let $S = (x_i)_i$ be a chosen basis of a free group F . The following requirement defines a derivation of $\mathbb{k}F$:

$$\frac{\partial}{\partial x_i} : x_t \mapsto \begin{cases} 1 & \text{if } t = i, \\ 0 & \text{else.} \end{cases}$$

Let us give a first version of the chainrule:

Proposition 2.4. *Let $\lambda : F_Y \rightarrow G$ a group morphism, where F_Y is the free group on a set $Y = \{y_j\}$. Then, for u in $\mathbb{k}Y$ and ∂ in $\text{Der}(\mathbb{k}G)$:*

$$\partial(\lambda u) = \sum_j \lambda \left(\frac{\partial u}{\partial y_j} \right) \partial(\lambda y_j).$$

Remark 2.5. The sums involved here are finite, because only a finite number of letters appear in a given element u .

Proof of proposition 2.4. One can check that each member of this equality defines a derivation from $\mathbb{k}F_Y$ to $\mathbb{k}G$, where $\mathbb{k}F_Y$ acts on $\mathbb{k}G$ by $y \cdot g = \lambda(y)g$. As these formulas give the same result when evaluated at elements of the basis, the corresponding derivations are equal. \square

If $G = F_X = \langle x_i \rangle$ is free too, we can apply proposition 2.4 with $\partial = \frac{\partial}{\partial x_i}$ to get:

$$\frac{\partial(\lambda u)}{\partial x_i} = \sum_j \lambda \left(\frac{\partial u}{\partial y_j} \right) \frac{\partial(\lambda y_j)}{\partial x_i}.$$

For $X = Y$ and $\lambda = 1_{F_X}$, this gives a change-of-base formula similar to the usual one.

The following definition keeps on with our analogy with classical differential calculus:

Definition 2.6. Let $F_Y = \langle y_j \rangle$ and $F_X = \langle x_i \rangle$ be free groups as above. Let f be a morphism from F_Y to F_X . We define its *Jacobian matrix* (with respect to the chosen basis) by:

$$D(f) := \left(\frac{\partial f(y_j)}{\partial x_i} \right)_{ji} \in M_{YX}(\mathbb{k}X).$$

Remark 2.7. The morphism f is determined by Df . Indeed, proposition 2.4, applied with $\lambda = 1$, $\partial : v \mapsto v - \varepsilon(v)$ and $u = f(x_i)$, gives:

$$f(x_i) - 1 = \sum_j \frac{\partial f(x_i)}{\partial x_j} (x_j - 1).$$

Let $F_Z = \langle z_k \rangle$, $F_Y = \langle y_j \rangle$ and $F_X = \langle x_i \rangle$ be free groups, and let $F_Z \xrightarrow{g} F_Y \xrightarrow{f} F_X$ be morphisms between them. We can use proposition 2.4 to get a chainrule for Jacobian matrices:

$$\frac{\partial f(g(z_k))}{\partial x_i} = \sum_j f \left(\frac{\partial g(z_k)}{\partial y_j} \right) \frac{\partial f(y_j)}{\partial x_i}.$$

This can be restated as:

Corollary 2.8. Let $F_Z \xrightarrow{g} F_Y \xrightarrow{f} F_X$ be morphisms between free groups with fixed basis, as above. Then:

$$D(fg) = f(Dg)D(f).$$

Remark 2.9. The reader may have noticed that this formula seems to come "in the wrong way". This can be explained as follows: a morphism $f : F_Y \rightarrow F_X$ is in fact a r -tuple of monomials $f(y_j) = f_j(x_i) \in \mathbb{k}F_X$, and should as such be considered as a "polynomial function from F_X to F_Y ", whose coordinates would be the f_j s. From this point of view, fg would be a "polynomial function from F_X to F_Z ", whose coordinates would be given by $fg(z_k) = g_k(f_j(x_i))$: it looks more like " $g \circ f$ " !

This seems to be completely analogous to the classical setting of algebraic geometry. To get more accurate statements, one would have to interpret $\mathbb{k}F_X$ as an algebra of functions over a geometric object associated to F_X . Which should look like a one-point object with some local structure (as Df determines f).

2.2 Derivations and strongly central filtrations

We introduce the notion of a derivation from a group G to a group H on which it acts. Our aim here is to describe a general framework which will be useful to study Jacobian matrices and their interactions with Lie brackets: the chainrule formula 2.8 tells us that D is a derivation. We will also get back to Lazard's theorem 1.37 in this framework.

Definition 2.10. Let H be a group, on which another group G acts. A map $\partial : G \rightarrow H$ is a *derivation* if:

$$\forall x, y \in G, \partial(xy) = \partial x \cdot {}^x\partial y.$$

Remark 2.11. If $H = M$ is an abelian group, *i.e.* a representation of G , then we recover the usual definition of a derivation from G to M (see Definition 2.1).

To give a derivation $\partial : G \rightarrow H$ is exactly the same as giving a section $\sigma = (\partial, 1_G)$ of the canonical projection:

$$H \rtimes G \begin{array}{c} \xleftarrow{\sigma} \\ \xrightarrow{p} \end{array} G.$$

Keeping this in mind, the following lemma follows immediately:

Lemma 2.12. Let ∂ be a derivation from G to H . Then ∂^{-1} sends G -stable subgroups of H on subgroups of G .

Let H_* be strongly central filtration on a subgroup H_1 of H . Let G_* be a strongly central filtration on a subgroup G_1 of G , which acts on H_* through the given action of G on H (see Proposition 1.20). A derivation ∂ from G to H being given, we can use the morphism $\sigma = (\partial, 1_G)$ to pull back the filtration $H_* \rtimes G_*$. We thus get a strongly central filtration on $G_1 \cap \partial^{-1}(H_1)$:

$$\sigma^{-1}(H_i \rtimes G_i) = G_i \cap \partial^{-1}(H_i).$$

Remark 2.13. For instance, if H_* is given, we can let G_* be $\mathcal{A}_*(G, H_*)$, the maximal filtration acting on H_* , as described in Remark 1.23. The above construction then gives a strongly central filtration on $\mathcal{A}_1 \cap \partial^{-1}(H_1)$. This subgroup is all of G if and only if:

$$\begin{cases} G \text{ stabilises } H_*, \\ G \text{ acts trivially on } \mathcal{L}(H_*), \\ \partial(G) \subseteq H_1. \end{cases} \quad (2.13.1)$$

Under these conditions, $\mathcal{A}_* \cap \partial^{-1}(H_*)$ is a strongly central series on G . In particular, it then contains $\Gamma_*(G)$.

Keeping the above notations, the morphism $\sigma = (\partial, 1_G)$ induces a Lie ring morphism (which is injective by definition of the filtration on the domain):

$$\bar{\sigma} : \mathcal{L}(G_* \cap \partial^{-1}(H_*)) \hookrightarrow \mathcal{L}(H_*) \rtimes \mathcal{L}(G_*).$$

This ensures that ∂ induces a well-defined linear map $\bar{\partial}$ between the Lie algebras. Moreover, the map $\bar{\sigma} = (\bar{\partial}, \bar{1}) : x \mapsto \bar{\partial}x + x$ preserves Lie brackets, hence:

$$\bar{\partial}([x, y]) = [\bar{\partial}x, y] + [x, \bar{\partial}y] + [\bar{\partial}x, \bar{\partial}y].$$

If $\mathcal{L}(H_*)$ is an abelian Lie algebra, then the last term is zero, and $\bar{\partial}$ is a Lie derivation. This happens in particular when H is an abelian group.

Back to the proof of Lazard's theorem 1.37. Take the filtered (abelian) group $(A, +)$ as H , and the group A^\times as G acting by left multiplication ρ . We already know that:

$$\mathcal{A}_j(A^\times, A_*) = A_*^\times = A^\times \cap (1 + A_j).$$

Let ∂ be the derivation from A^\times to A defined by $g \mapsto g - 1$. Obviously, $\partial^{-1}(A_j) = A_j^\times$, which is exactly $\mathcal{A}_j(A^\times, A_*)$. It is then equal to $\mathcal{A}_j(A^\times, A_*) \cap \partial^{-1}(A_j)$.

The Lie ring $\mathcal{L}(A_*)$ is abelian (because A is an abelian group). Hence, the induced map $\bar{\partial}$ is a derivation (with respect to the canonical action of $\mathcal{L}(A_*^\times)$ on $\mathcal{L}(A_*)$):

$$\bar{\partial} : \mathcal{L}(A_*^\times) \longrightarrow \mathcal{L}(A_*).$$

Let us remark that the Lie algebra $\mathcal{L}(A_*)$ is quite different from $\text{gr}(A_*)$: the associative structure of A has been completely forgotten. Nevertheless, some part of this structures is encoded by the action of $\mathcal{L}(A_*^\times)$, which is inherited from left multiplication. The map $\bar{\partial}$ is a derivation with respect to this action, that is:

$$\bar{\partial}([x, y]) = [\bar{\partial}x, y] + [x, \bar{\partial}y].$$

These brackets are described through the action of $\mathcal{L}(A_*^\times)$ on $\mathcal{L}(A_*)$, as induced by commutators in $A \rtimes A^\times$:

$$\forall g \in A_1^\times, \forall x \in A, [g, x] = gx - x.$$

As a consequence:

$$\bar{\partial}([x, y]) = -(y(x - 1) - (x - 1)) + (x(y - 1) - (y - 1)) = xy - yx,$$

so $\bar{\partial}$ is in fact a Lie morphism to $\text{gr}(A_*)$. □

2.3 Algebras, actions and derivations

We now turn to studying derivations of algebras. In particular, we get a precise link between free differential calculus and differential calculus in the tensor algebra (see Proposition 2.25). We will use this in Paragraph 2.4.2 to get an explicit description of the trace map.

Let \mathcal{Alg}_- be the category of associative non-unitary algebras over a fixed commutative ring \mathbb{k} . This category is pointed (by 0) and protomodular. We can define actions there, as in paragraph 1.3. Actions in \mathcal{Alg}_- turn out to be representable. Precisely, for any algebra I , let $\text{End}_r(I)$ (resp. $\text{End}_l(I)$) be the algebra of right (resp. left) I -linear endomorphisms of I , *i.e.* \mathbb{k} -linear maps u from I to I satisfying:

$$\forall x, y \in I, u(xy) = u(x)y \quad (\text{resp. } u(xy) = xu(y)).$$

Define $\text{End}_{r,l}(I)$ as the kernel of :

$$\alpha : \begin{cases} \text{End}_r(I) \times \text{End}_l(I) & \longrightarrow & \text{End}_{\mathbb{k}}(I) \\ (u, v) & \longmapsto & u \circ v - v \circ u. \end{cases}$$

Proposition 2.14. *An action $A \circlearrowleft I$ in \mathcal{Alg}_- can be represented by a (unique) morphism:*

$$A \xrightarrow{(\lambda, \rho)} \text{End}_{r,l}(I).$$

Proof. If an action

$$I \longleftarrow B \overset{\curvearrowright}{\longrightarrow} A$$

is given, $\lambda(a)$ and $\rho(a)$ are obtained from left and right multiplications by a in B . Conversely, a morphism (λ, ρ) as above can be used to define an (associative) algebra structure on $I \times A$ defining an action of A on I . \square

Let us remark that if $I^2 = 0$ (that is, I is endowed with a trivial algebra structure), then an action of A on I is just a A -bimodule structure.

Remark 2.15. The same construction works in the category of (non-unitary) filtered algebras $fAlg_-$, where we also get a representation of actions. The algebras $\text{End}_r(I)$ and $\text{End}_l(I)$ are then filtered by the usual requirement: a morphism u is of degree at least j if $u(I_i) \subseteq I_{i+j}$ for all i . The same requirement will be used to define a filtration on any module of morphisms between filtered modules or algebras.

Definition 2.16. Let A act on I as above. A *derivation* from A to I is a \mathbb{k} -linear map $\partial : A \rightarrow I$ satisfying:

$$\partial(ab) = \partial a \cdot b + a \cdot \partial b.$$

The \mathbb{k} -module of derivations from A to I is denoted by $\text{Der}(A, I)$.

Remark 2.17. The relation defining derivations depends only on the A -bimodule structure on I . We are thus led to consider I° , the algebra obtained by taking the same underlying \mathbb{k} -module as I , endowed with the trivial product. The action of A on I induces an action of A on I° in the obvious manner, and a derivation from A to I is then the same as a section of the projection:

$$I^\circ \rtimes A \twoheadrightarrow A$$

in the category of algebras.

When we work in the category of filtered algebras, $\text{Der}(A, I)$ is a filtered module, a derivation ∂ being of degree at least j if $\partial(A_i) \subseteq I_{i+j}$ for all i . If A is filtered by its powers A^i , we just have to check this in degree one:

Lemma 2.18. *Let A be an algebra, filtered by its powers $A_i := A^i$, acting on a filtered algebra I_* . Let $\partial \in \text{Der}(A, I)$. Then ∂ is of degree at least j if and only if:*

$$\partial(A) \subseteq I_{j+1}.$$

Proof. An action of A_* on I_* is given by a left and a right multiplication which are filtered, meaning that $A_i I_j \subseteq I_{i+j}$ and $I_j A_i \subseteq I_{i+j}$. Use the formula:

$$\partial(a_1 \cdots a_i) = \sum_k a_1 \cdots a_{k-1} \partial(a_k) a_{k+1} \cdots a_i$$

to get the desired result. \square

We can get examples of actions from algebras acting on themselves. Precisely, the adjoint action of A on itself is just the obvious A -bimodule structure on A . Derivations are then the usual ones.

Given an action of A on I represented by (ρ, λ) , we can twist it by choosing endomorphisms φ and ψ of A and letting A act on I through $(\rho \circ \varphi, \lambda \circ \psi)$. This means that we let $a \in A$ act on I by $\varphi(a) \cdot -$ on the left, and by $- \cdot \psi(a)$ on the right. We give a name to derivations from A to the the twisted A -bimodule I .

Definition 2.19. Let φ and ψ be endomorphisms of A . A (φ, ψ) -derivation is a linear map $\partial : A \rightarrow I$ satisfying:

$$\partial(ab) = \partial a \cdot \psi b + \varphi a \cdot \partial b.$$

We denote by $\text{Der}_{(\varphi, \psi)}(A, I)$ the \mathbb{k} -module of such derivations.

Example 2.20. Let A be a group algebra $\mathbb{k}G$ and M a $\mathbb{k}G$ -module. We can make M into a bimodule by making $\mathbb{k}G$ act trivially on the right (that is, through ε). Then, $\text{Der}(\mathbb{k}G, M)$ is exactly the usual module of derivations (see Definition 2.1). If $M = \mathbb{k}G$, it is already a bimodule, but the above structure can be obtained through twisting the right action by $\eta\varepsilon : g \mapsto \varepsilon(g) \cdot 1$. Then $\text{Der}(\mathbb{k}G)$ (defined in Definition 2.1) is exactly $\text{Der}_{(id, \eta\varepsilon)}(\mathbb{k}G, \mathbb{k}G)$.

We can apply Lemma 2.18 to $A = I = IG$, to get:

Corollary 2.21. Let ∂ be a derivation of $\mathbb{Z}G$ such that $\partial(\mathbb{Z}G) \subseteq (IG)^{l+1}$ (which is always true for $l = -1$). For all integer k , we have:

$$\partial((IG)^k) \subseteq (IG)^{k+l}.$$

Remark 2.22. Let us stress that the proof given here is fairly direct. In fact, it gets even shorter in this case, the result following from: $\forall v \in I, \partial(uv) = u \cdot \partial v$.

Remark 2.23. The classical inclusion of $\Gamma_k - 1$ into IG^k can be shown by a direct induction, or follows from Lazard's theorem (as $D_*(G)$ is a strongly central series, it contains Γ_*G). Under the hypothesis of Proposition 2.21, it implies $\partial(\Gamma_k) \subseteq (IG)^{k+l}$, using that $\partial(1) = 0$.

Remark 2.24. Some sets of derivations obtained from actions of A on itself can have more structure than just a module structure. Precisely, if we twist the adjoint action of A by some (φ, ψ) as above, and if we add the requirement that φ and ψ are idempotents, then the set of (φ, ψ) -derivations from A to A commuting to φ and ψ is a sub-Lie algebra of $\text{End}_{\mathbb{k}}(A)$:

$$\begin{aligned} [\partial, \partial'](ab) &= \partial \partial' a \cdot \psi^2 b + \varphi \partial' a \cdot \partial \psi b + \partial \varphi a \cdot \psi \partial' b + \varphi^2 a \cdot \partial \partial' b \\ &\quad - \partial' \partial a \cdot \psi^2 b - \varphi \partial a \cdot \partial' \psi b - \partial' \varphi a \cdot \psi \partial b - \varphi^2 a \cdot \partial' \partial b \\ &= [\partial, \partial'](a) \cdot \psi(b) + \varphi(a) \cdot [\partial, \partial'](b). \end{aligned}$$

In particular, if $(\varphi, \psi) = (1, 1)$, we get that $\text{Der}(A)$ is a sub-Lie algebra of $\text{End}_{\mathbb{k}}(A)$. Another example is given by $A = \mathbb{k}G$ and $(\varphi, \psi) = (1, \eta\varepsilon)$. But more is true in this last case. Let $\mathbb{k}G$ be filtered by the powers of the augmentation ideal. If $\partial, \partial' \in \text{Der}(\mathbb{k}G)$ are such that ∂' has degree at least 0, then $\partial \circ \partial' \in \text{Der}(\mathbb{k}G)$, because $\varepsilon(\partial'v) = 0$ for any v .

Let $A_* \circlearrowleft I_*$ be an action of filtered algebras. Since the functor $\text{gr} : f\text{Alg} \rightarrow \text{gr Alg}$ from filtered algebras to graded ones is exact (the same proof as that of Proposition 1.24 works), this action is sent to an action $\text{gr}(A_*) \circlearrowleft \text{gr}(I_*)$ of graded algebras. Moreover, gr also commutes with $(-)^o$ (the definition of $(-)^o$ being extended to graded algebras in the obvious way), so we get a morphism:

$$\text{gr}(\text{Der}(A_*, I_*)) \hookrightarrow \text{Der}_*(\text{gr}(A_*), \text{gr}(I_*)),$$

where the target is the graded module of graded derivations. This morphism is obviously injective. It is in fact a restriction of the natural injection:

$$\mathrm{gr}(\mathrm{Hom}_{\mathbb{k}}(M_*, N_*)) \hookrightarrow \mathrm{Hom}_*(\mathrm{gr}(M_*), \mathrm{gr}(N_*))$$

between bifunctors on graded modules. As such, it preserves all algebraic structure inherited from the additive bifunctor structure (see Remark 2.24).

When A_* is $\mathbb{k}G$, filtered by the powers of IG , acting on itself, we thus get a morphism preserving the structure induced by composition:

$$\mathrm{gr}(\mathrm{Der}(\mathbb{k}G)) \hookrightarrow \mathrm{Der}_*(\mathrm{gr}(\mathbb{k}G)).$$

Proposition 2.25. *If G is a free group, and M_* is a filtered $\mathbb{k}G$ -module (considered as a bimodule with trivial right action), the canonical map:*

$$\mathrm{gr}(\mathrm{Der}(\mathbb{k}G, M_*)) \hookrightarrow \mathrm{Der}_*(\mathrm{gr}(\mathbb{k}G), \mathrm{gr}(M_*))$$

is an isomorphism. Here, by derivations, we mean (id, ε) -ones.

Proof. Let S be a free set of generators for G . Then $V = G^{ab}$ is free abelian on S , and $\mathrm{gr}(\mathbb{k}G) \cong TV$ is the tensor algebra. Identifying derivations with sections as above (see Remark 2.17), we see that a derivation is completely determined by the choice of its values on S :

$$\mathrm{Der}_*(TV, N_*) = \mathcal{F}_*(S, N_*),$$

for any graded TV -bimodule N_* , where $\mathcal{F}_*(S, N_*)$ is the set of graded maps from S (concentrated in degree 0) to N_* . The same is true for the other side. Indeed, a derivation from $\mathbb{k}G$ to M is a section of the projection $M \rtimes G \rightarrow G$, so is determined by a map $S \rightarrow M$:

$$\mathrm{Der}(\mathbb{k}G, M_*) = \mathrm{Der}(G, M_*) \cong M_*^S.$$

The second member is the set of maps from S to M , with the filtration inherited from the one on M . The desired isomorphism is then exactly: $\mathrm{gr}(M_*^S) \cong \mathcal{F}_*(S, \mathrm{gr}(M_*))$. \square

Remark 2.26. If M is a G -module, we can endow it with the universal $\mathbb{k}G$ -filtration $(IG)^* \cdot M$.

Remark 2.27. The isomorphism $\mathrm{gr}(\mathrm{Der}(\mathbb{k}G)) \cong \mathrm{Der}_*(TV)$ thus obtained preserves the algebraic structure obtained from the composition of derivations.

2.4 Traces

In [Bar13], Bartholdi defines the trace of an automorphism φ of F_n by:

$$\mathrm{Tr}(\varphi) := \mathrm{Tr}(D\varphi - 1) \in \mathbb{Z}F_n,$$

where $D\varphi$ denotes φ 's Jacobian matrix. We will show that Tr induces a well-defined map between the graded Lie algebras, which we still call Tr :

$$\mathrm{Tr} : \mathcal{L}(\mathcal{A}_*(F_n)) \longrightarrow \mathrm{gr}(\mathbb{Z}F_n) \cong TV.$$

The aim of this paragraph is to show that this map is indeed well-defined, to investigate its behaviour with respect to Lie structures, and to get Morita's algebraic description [Mor93], used by Satoh in [Sat12].

2.4.1 The induced map between Lie algebras

Let $\varphi \in \mathcal{A}_k(F_n)$. By definition, $\varphi_i := x_i^{-1}\varphi(x_i) \in \Gamma_{k+1}$. The Jacobian matrix of φ can be described explicitly:

$$(D\varphi)_{ij} = \frac{\partial(x_i\varphi_i)}{\partial x_j} = \underbrace{\frac{\partial x_i}{\partial x_j}}_{\delta_{ij}} + x_i \frac{\partial(\varphi_i)}{\partial x_j}.$$

Hence:

$$D\varphi - 1 = \left(x_i \frac{\partial \varphi_i}{\partial x_j} \right)_{ij}.$$

Using Remark 2.23, we see that this matrix is in fact in $M_n(I^k)$ (to shorten notations, we write I for IF_n in the sequel). Moreover, x_i acts trivially on I^k/I^{k+1} . We thus get an explicit formula for the trace map:

$$\mathrm{Tr}(D\varphi - 1) = \sum_i x_i \frac{\partial \varphi_i}{\partial x_i} \equiv \sum_i \frac{\partial \varphi_i}{\partial x_i} \pmod{I^{k+1}}.$$

Let G be any group. We can apply the construction of Paragraph 1.7 to $A = \mathbb{k}G$, filtered by the powers of the augmentation ideal. This gives a strongly central filtration $GL_n(I^*G)$ on $GL_n(IG)$, which comes with an embedding of Lie algebras:

$$\mathcal{L}(GL_n(I^*G)) \hookrightarrow \mathrm{gr}(M_n(\mathbb{k}G)) \cong M_n(\mathrm{gr}(\mathbb{k}G)).$$

The next proposition replaces a formula from [Bar13, section 6]:

Proposition 2.28. *The Jacobian matrix D induces a morphism between graded modules:*

$$D : \mathcal{L}(\mathcal{A}_*(F_n)) \longrightarrow \mathcal{L}(GL_n(I^*F_n)),$$

satisfying:

$$D([f, g]) = [g, Df] + [Dg, f] + [Dg, Df].$$

In order to address the issue raised in Remark 2.9, let us introduce some notations before proving the proposition. If G is any group, we denote by G^{op} the *opposite group*, where multiplication is defined by: $g \cdot_{op} h = hg$. Let G_* be a strongly central filtration on G . Then G_*^{op} is such a filtration on G^{op} and one easily checks that:

$$\mathcal{L}(G_*^{op}) = \mathcal{L}(G_*)^{op},$$

where the bracket in $\mathcal{L}(G_*)^{op}$ is $\mathcal{L}(G_*)$'s additive inverse: $[x, y]_{op} = [y, x]$.

Proof of Proposition 2.28. Corollary 2.8 states exactly that D is a derivation from $\mathrm{Aut}(F_n)$ to $GL_n(I)^{op} \subset M_n(\mathbb{Z}F_n)^{op}$, where $M_n(\mathbb{Z}F_n)$ is endowed with the obvious $\mathrm{Aut}(F_n)$ -action. We thus can apply the results from 2.2 with $\partial = D$, $G = \mathrm{Aut}(F_n)$, $H = GL_n(I)^{op}$, and $H_* = GL_n(I^*)^{op}$.

The strongly central filtration $\mathcal{A}_*(G, H_*)$ is in fact the Andreadakis filtration $\mathcal{A}_* = \mathcal{A}_*(F_n)$ on $\mathcal{A}_1 = IA_n \subset \mathrm{Aut}(F_n)$. Indeed, there is a series of inclusions:

$$\mathcal{A}_*(G, D_*F_n) \supseteq \mathcal{A}_*(G, GL_n(I^*)) \supseteq \mathcal{A}_*(G, M_n(\mathbb{Z}F_n)) = \mathcal{A}_*(G, \mathbb{Z}F_n).$$

The first one comes from 1.49 applied to the injection of D_*F_n into $GL_n(I^*)$ defined by $w \mapsto w \cdot 1$. The second one is a particular case of 1.52. The last equality comes from the fact that G acts component-wise on matrices:

$$[g, (m_{ij})] = g \cdot (m_{ij}) - (m_{ij}) = ([g, m_{ij}]).$$

According to proposition 1.53, these inclusions are in fact equalities.

Moreover, we have seen at the beginning of the present paragraph that D sends \mathcal{A}_* to $GL_n(I^*)^{op}$. The filtration $\mathcal{A}_* \cap \partial^{-1}(H_*)$ is thus only \mathcal{A}_* . The work already done in 2.2 allows us to get the desired result. \square

The map given by Proposition 2.28 can be composed with the morphism:

$$\mathcal{L}(GL_n(I^*)) \xrightarrow{(-)-1} \text{gr}(M_n(\mathbb{Z}F_n)) \cong M_n(\text{gr}(\mathbb{Z}F_n)) = M_n(TV).$$

Thus, for φ an element of $\mathcal{A}_k/\mathcal{A}_{k+1}$, $D\varphi - 1$ is well-defined modulo $M_n(I^{k+1})$. Composing with the usual trace, we get the announced well-defined linear map induced by $\varphi \mapsto \text{Tr}(D\varphi - 1)$:

$$\text{Tr} : \mathcal{L}(\mathcal{A}) \longrightarrow TV.$$

Remark 2.29. That the map $D-1$ (hence Tr) induces a well-defined map between the Lie algebras can be seen through explicit calculation, but the behaviour with respect to the Lie bracket is much less obvious from this point of view. Indeed, let $\varphi \in \mathcal{A}_k$. If $\varphi = \psi\chi$, with $\chi \in \mathcal{A}_{k+1}$, then:

$$\psi(x_i) = \varphi(x_i\chi_i) = x_i\varphi_i\varphi(\chi_i).$$

As χ_i stands inside Γ_{k+2} , its image by φ does too. Thus:

$$\frac{\partial\psi_i}{\partial x_j} = \frac{\partial\varphi_i}{\partial x_j} + \underbrace{\varphi_i \frac{\partial\varphi(\chi_i)}{\partial x_j}}_{\in I^{k+1}} \equiv \frac{\partial\varphi_i}{\partial x_j} \pmod{I^{k+1}}.$$

2.4.2 Introducing the contraction map

Consider the evaluation map:

$$ev : \text{Der}_{(1,\varepsilon)}(TV) \otimes TV \longrightarrow TV.$$

Using the universal property of TV , as in the proof of 2.25, we get a linear isomorphism:

$$\text{Der}_{(1,\varepsilon)}(TV) \cong \text{Hom}(V, TV) \cong V^* \otimes TV.$$

The evaluation map then is:

$$\begin{cases} V^* \otimes TV \otimes TV & \longrightarrow TV \\ \omega \otimes u \otimes v & \longmapsto \Phi(\omega \otimes v)u, \end{cases}$$

where Φ is the contraction map:

$$\Phi : \begin{cases} V^* \otimes V^{\otimes k+1} & \longrightarrow V^{\otimes k} \\ \alpha \otimes X_{i_1} \cdots X_{i_{k+1}} & \longmapsto X_{i_1} \cdots X_{i_k} \alpha(X_{i_{k+1}}), \end{cases}$$

extended by zero on $\mathbb{k} \cdot 1$. This follows from the fact that any $(1, \varepsilon)$ -derivation ∂ verifies:

$$\partial(uv) = u \cdot \partial v,$$

when the degree of v is at least 1 (that is, when $\varepsilon(v) = 0$), and $\partial(1) = 0$.

We sum this up in the following:

Proposition 2.30. *Let $\partial \in \text{Der}_{(1,\varepsilon)}(TV)$. Then:*

$$\partial = \Phi(\partial|_V \otimes -).$$

Let us consider the derivation $\frac{\partial}{\partial x_i}$ of $\mathbb{Z}F_n$. It induces a $(1, \varepsilon)$ -derivation of degree -1 of TV , denoted by ∂_i (any derivation of $\mathbb{k}G$ is of degree at least -1 , by Corollary 2.21). As $\partial_i|_V = X_i^*$, we get the following:

Corollary 2.31. *The $(1, \varepsilon)$ -derivation of TV induced by $\frac{\partial}{\partial x_i} \in \text{Der}(\mathbb{Z}F_n)$ is represented as:*

$$\overline{\frac{\partial}{\partial x_i}} = \partial_i = \Phi(X_i^* \otimes -) : TV \longrightarrow TV.$$

We can use these results to interpret the trace map in a way more suited to explicit calculations:

Proposition 2.32. *The trace map can be described as:*

$$\text{Tr} = \Phi \circ \iota \circ \tau,$$

where τ is the Johnson morphism (see Definition 1.26), ι denotes the inclusion of $\text{Der}_k(LV) \cong V^* \otimes L_{k+1}V$ into $\text{Der}_k(TV) \cong V^* \otimes V^{\otimes k+1}$, and Φ is the contraction map.

Proof. Let $\varphi \in \mathcal{A}_k$. Then $\tau(\varphi)$ is defined by:

$$\tau(\varphi)(X_i) = \overline{x_i^{-1}\varphi(x_i)} = \overline{\varphi_i} \in \Gamma_{k+1}/\Gamma_{k+2} \cong \mathfrak{L}_{k+1}V.$$

We have seen at the beginning of Paragraph 2.4.1 that the trace map is given by:

$$\text{Tr}(D\varphi - 1) = \sum_i \frac{\partial \varphi_i}{\partial x_i}.$$

The formula of the proposition is then equivalent to:

$$\Phi \left(\sum_i X_i^* \otimes \overline{\varphi_i} \right) = \overline{\sum_i \frac{\partial \varphi_i}{\partial x_i}}.$$

To get this formula, we evaluate the equality given by Corollary 2.31 to the elements $\overline{\frac{\varphi_i - 1}{w - 1}}$ (keeping in mind that the inclusion of $L_{k+1}V$ into $T_{k+1}V$ is given by $\overline{w} \mapsto \overline{w - 1}$). \square

2.5 Stable surjectivity

2.5.1 Vanishing of the trace map

Here, we show that the trace map takes values in brackets inside TV . This result can also be found in [MS17, Prop. 5.3], where rational methods are used to get it.

Proposition 2.33 ([BLGM90, Th. 2.1], quoted in [Bar13, Th. 6.2]). *Let $k \geq 2$, and let $J \in GL_m(I_{\mathbb{Z}}^k F_n)$. Denote by V the abelianization $V = F_n^{ab} \cong \mathbb{Z}^n$. Then:*

$$\text{Tr}(J - 1) \in [TV, TV]_k \subset V^{\otimes k} \cong I^k / I^{k+1}.$$

This result relies on the following criterion:

Proposition 2.34 ([BLGM90, prop. 2.2]). *Let $f(X_1, \dots, X_n) \in V^{\otimes k}$. Let $\mathcal{C} \subset M_k(\mathbb{Z})$ be the sub- \mathbb{Z} -module generated by the $e_{i,i+1}$. Suppose:*

$$\forall C_i \in \mathcal{C}, \quad \text{Tr}(f(C_1, \dots, C_n)) = 0.$$

Then $f \in [TV, TV]_k$.

The proof can be found in [BLGM90]. The reader is also referred to the proof of Proposition 3.16, which is the same proof, adapted to the case of positive characteristic.

Proof of Proposition 2.33. The main idea is to use evaluations into commutative algebras to be able to use Proposition 1.41, and to then get back to the non-commutative setting by using the above criterion.

Let $1 + tA_i \in GL_k(t\mathbb{k}[t])$. There is an evaluation morphism $x_i \mapsto 1 + tA_i$ from F_n to $GL_k(t\mathbb{Z}[t])$, extending to a morphism from $\mathbb{k}F_n$ to $M_k(t\mathbb{k}[t])$ sending I^* to $t^*M_k(\mathbb{k}[t])$. Taking congruence groups, we get an evaluation morphism:

$$ev_{1+tA_i} : GL_m(I^*F_n) \longrightarrow GL_m(t^*M_k(\mathbb{k}[t])) = GL_{mk}(t^*\mathbb{k}[t]).$$

There is a commutative diagram:

$$\begin{array}{ccccc} \mathcal{L}(GL_m(I^*F_n)) & \xleftarrow{(-)-1} & M_m(TV) & \xrightarrow{Tr} & TV \\ \downarrow ev_{1+tA_i} & & \downarrow ev_{tA_i} & & \downarrow ev_{tA_i} \\ \mathcal{L}(GL_{mk}(t^*\mathbb{k}[t])) & \xleftarrow{(-)-1} & M_{mk}(\mathbb{k}[t]) & \xrightarrow{Tr_m} & M_m(\mathbb{k}[t]) \\ \downarrow det & & \downarrow Tr & \swarrow Tr & \\ \mathcal{L}(\mathbb{k}[t]_*^\times) = 0 & \xleftarrow{(-)-1} & \mathbb{k}[t] & & \end{array}$$

Here, we identify $\text{gr}(I^*F_n)$ with TV by $x_i \mapsto 1 + X_i$. We also identify $\text{gr}(t^*\mathbb{k}[t])$ with $\mathbb{k}[t]$. The evaluation $x_i \mapsto 1 + tA_i$ thus induces $X_i \mapsto tA_i$ between the associated graded. The bottom-left square is just the one in Proposition 1.41. The map Tr_m is the usual trace when the base algebra is $M_m(\mathbb{k}[t])$.

Let $\mathbb{k} = \mathbb{Z}$ and $f = \text{Tr}(J - 1) \in V^{\otimes k}$. The commutativity of the above diagram gives:

$$0 = \text{Tr}(f(tA_i)) = t^k \text{Tr}(f(A_i)).$$

As a consequence, $\text{Tr}(f(A_i)) = 0$, for any $1 + tA_i \in GL_k(t\mathbb{Z}[t])$. We can then evaluate this at $t = 0$ to get: $\text{Tr}(f(\pi A_i)) = 0$. This evaluation π is the map:

$$\pi : \mathcal{L}_1(GL_k(t^*\mathbb{Z}[t])) \hookrightarrow M_k(t\mathbb{Z}[t]/t^2\mathbb{Z}[t]) \cong M_k(\mathbb{Z}).$$

Using Proposition 1.42 and Remark 1.43, we see that its image is exactly $\mathfrak{sl}_n(\mathbb{Z})$, so the conclusion follows from the above criterion (2.34). \square

Because of Proposition 2.33, we will consider the trace map as taking values in the abelianization $TV^{ab} = TV/[TV, TV]$. As $[TV, TV]_k$ is generated by the elements:

$$[X_{i_1} \otimes \dots \otimes X_{i_p}, X_{i_{p+1}} \otimes \dots \otimes X_{i_k}] = X_{i_1} \otimes \dots \otimes X_{i_k} - t^p \cdot X_{i_1} \otimes \dots \otimes X_{i_k},$$

where $t = \bar{1} \in \mathbb{Z}/k$, the module TV^{ab} is the module of *cyclic powers* C_*V :

$$(TV^{ab})_k = C_kV := V^{\otimes k}/(\mathbb{Z}/k).$$

The conclusion of Proposition 2.33 becomes, in this context: $\text{Tr}(J - 1) = 0 \in C_*V$.

2.5.2 Linear algebra

Consider the Johnson morphism $\tau' : \mathcal{L}(\Gamma_{IA_n}) \longrightarrow \text{Der}_*(\mathcal{L}(F_n))$ (cf. 1.5). The morphism τ'_1 is an isomorphism (see Proposition 1.34). Moreover, $\mathcal{L}(\Gamma_{IA_n})$ is generated in degree 1 (cf. 1.9). As a consequence, the image of τ' is exactly the sub-Lie ring generated in degree 1 inside $\text{Der}(\mathcal{L}(F_n))$. As $\mathcal{L}(F_n)$ is the free Lie ring $\mathfrak{L}V$, the study of $\text{coker}(\tau')$ is solely a problem of linear algebra.

Recall from Proposition 2.32 that the trace map can be seen as the composite of the Johnson morphism $\tau : \mathcal{L}_k(\mathcal{A}_*(F_n)) \rightarrow \text{Der}_k(\mathfrak{L}V) \cong V^* \otimes \mathfrak{L}_{k+1}V$ with:

$$\text{Tr}_M : V^* \otimes \mathfrak{L}_{k+1}V \xrightarrow{\iota} V^* \otimes V^{\otimes k+1} \xrightarrow{\Phi} V^{\otimes k} \xrightarrow{\pi} C_k V := V^{\otimes k}/(\mathbb{Z}/(k)),$$

where ι and π denote the canonical maps. All these morphisms are obviously $\text{GL}_n(\mathbb{Z})$ -equivariant (with respect to the canonical actions).

Notation 2.35. Let \mathfrak{J} denote the image of τ' , which is the sub-Lie ring generated in degree 1 inside $\text{Der}(\mathfrak{L}V)$.

The following proposition can be seen as a consequence of Proposition 2.33. Precisely, $\mathfrak{J} = \text{Im}(\tau') \subseteq \text{Im}(\tau)$, and Proposition 2.33 implies that $\text{Tr}_M \circ \tau = \text{Tr}$ vanishes.

Proposition 2.36. *For every $k \geq 2$, $\text{Tr}_M(\mathfrak{J}_k) = \{0\}$.*

2.5.3 Stable cokernel of τ' and stable surjectivity

Let $k \geq 2$. Using Proposition 2.36, we get a commutative diagram with exact rows:

$$\begin{array}{ccccc} \mathfrak{J}_k & \hookrightarrow & V^* \otimes \mathfrak{L}_{k+1}V & \twoheadrightarrow & \text{coker}(\tau'_k) \\ \downarrow \phi & & \downarrow \Phi & & \downarrow \bar{\Phi} \\ [TV, TV]_k & \hookrightarrow & V^{\otimes k} & \xrightarrow{\pi} & C_k V. \end{array}$$

In [Sat12], Satoh shows:

$$\left\{ \begin{array}{l} \text{For } n \geq k+1, \quad \bar{\Phi} \text{ is surjective,} \quad (\text{lemma 3.2}) \\ \text{For } n \geq k+2, \quad \phi \text{ is surjective,} \quad (\text{prop. 3.2}) \\ \text{For } n \geq k+2, \quad \ker \bar{\Phi} \subseteq \mathfrak{J}. \quad (\text{Prop. 3.3}) \end{array} \right.$$

His Theorem 3.1 is still true over \mathbb{Z} :

Proposition 2.37. *Let $k \geq 2$ and $n \geq k+2$ be integers. Then $\bar{\Phi}$ is a $\text{GL}_n(\mathbb{Z})$ -equivariant isomorphism:*

$$\text{coker}(\tau'_k) \cong C_k V.$$

Proof. Let us denote by K (resp. L) the kernel of $\bar{\Phi}$ (resp. its cokernel). There is a commutative diagram in $\text{GL}_n(\mathbb{Z}) - \text{Mod}_{\mathbb{Z}}$:

$$\begin{array}{ccccc} & & \ker \Phi & \overset{0}{\dashrightarrow} & K \\ & & \downarrow & & \downarrow \\ \mathfrak{J}_k & \hookrightarrow & V^* \otimes \mathfrak{L}_{k+1}V & \twoheadrightarrow & \text{coker}(\tau') \\ \downarrow \phi & & \downarrow \Phi & & \downarrow \bar{\Phi} \\ [TV, TV]_k & \hookrightarrow & V^{\otimes k} & \xrightarrow{\pi} & C_k V \\ \downarrow & & \downarrow & & \downarrow \\ 0 & & 0 & & L. \end{array}$$

The snake lemma ensures that K and L are zero: $\overline{\Phi}$ is an isomorphism. \square

We can now state our main result:

Theorem 2.38. *Let $k + 2 \leq n$. Then the canonical morphism*

$$\mathcal{L}_k(IA_n) \longrightarrow \mathcal{L}_k(\mathcal{A}_*(F_n))$$

is surjective, and τ induces an isomorphism: $\mathcal{L}_k(\mathcal{A}_(F_n)) \cong \mathfrak{J}_k$.*

Remark 2.39. Basis being chosen, there is an injection of F_n in $F_{n+1} \cong F_n * \mathbb{Z}$. An automorphism φ of F_n can be extended to an automorphism $\varphi * 1$ of F_{n+1} . This induces injections $IA_n \hookrightarrow IA_{n+1}$ which in turn induce morphisms $\mathcal{L}(\mathcal{A}_*(F_n)) \rightarrow \mathcal{L}(\mathcal{A}_*(F_{n+1}))$. Taking the colimit over n , we can define a Lie ring $\mathcal{L}^{st}(\mathcal{A}_*)$. In the same way, we can define injections from $\text{Der}(\mathcal{L}(F_n^{ab}))$ into $\text{Der}(\mathcal{L}(F_{n+1}^{ab}))$ and take the colimit \mathfrak{J}^{st} of the sub-algebras generated in degree one. With this point of view, the isomorphisms of Theorem 2.38 give an isomorphism between graded Lie algebras:

$$\tau^{st} : \mathcal{L}^{st}(\mathcal{A}_*) \cong \mathfrak{J}^{st},$$

meaning exactly that $\mathcal{L}^{st}(\mathcal{A})$ is generated in degree one.

In fact, all the constructions appearing here are functors on the category denoted by $\mathbf{S}(\mathbb{Z})$ in [Dja16b, section 7], where it is shown (using methods similar to the ones of [CEFN14]) that these functors are finitely supported. This implies the equivalence between τ_k^{st} being an isomorphism and τ_k being one for n big enough.

Proof of theorem 2.38. Consider the commutative diagram:

$$\begin{array}{ccc} \mathcal{L}(IA_n) & \xrightarrow{i_*} & \mathcal{L}(\mathcal{A}_*) \\ & \searrow \tau' & \downarrow \tau \\ & & \text{Der}(\mathfrak{L}V). \end{array}$$

The image \mathfrak{J} of τ' is the sub-Lie ring generated in degree one inside $\text{Der}(\mathfrak{L}V)$. Using the results quoted in Paragraph 2.5.3, we see that in degrees $k \leq n - 2$, it also is the kernel of the trace map. Proposition (2.33) tells us exactly that $\text{Tr} \circ \tau = 0$, so that $\text{Im } \tau_k \subseteq \ker \text{Tr}_k = \text{Im } \tau'_k$ when $k \geq n - 2$. As a consequence, $\text{Im } \tau = \text{Im } \tau'$. As τ is injective (1.32), it is an isomorphism onto its image, hence the result. \square

2.6 Automorphisms of free nilpotent groups

Automorphisms of nilpotent groups are easy to deal with, due to the following classical fact:

Lemma 2.40. *Let G be a finite-type nilpotent group. An endomorphism $\varphi \in \text{End}(G)$ is an automorphism if and only if the induced morphism $\varphi^{ab} \in \text{End}(G^{ab})$ is.*

Proof. If φ is an automorphism then φ^{ab} has to be, with $(\varphi^{ab})^{-1} = (\varphi^{-1})^{ab}$. Conversely, suppose that φ^{ab} is an automorphism. This means that $\mathcal{L}_1(\varphi)$ is. Since $\mathcal{L}(G)$ is generated in degree one, $\mathcal{L}(\varphi)$ is surjective. But each $\mathcal{L}_k(G)$ is abelian of finite type, so each $\mathcal{L}_k(\varphi)$, being surjective, has to be bijective (it is obviously the case on the torsion

part, which is finite, and it also is on the free abelian part, for reasons of rank). The lemma then follows by induction from the five-lemma applied to:

$$\begin{array}{ccccc} \mathcal{L}_k(G) & \hookrightarrow & G/\Gamma_{k+1}G & \twoheadrightarrow & G/\Gamma_kG \\ \downarrow \mathcal{L}_k(\varphi) & & \downarrow \tilde{\varphi} & & \downarrow \tilde{\varphi} \\ \mathcal{L}_k(G) & \hookrightarrow & G/\Gamma_{k+1}G & \twoheadrightarrow & G/\Gamma_kG. \end{array}$$

This induction process stops since there is a c such that $G = G/\Gamma_{c+1}G$. \square

Definition 2.41. The *pro-nilpotent completion* of a group G is: $\widehat{G} := \varprojlim (G/\Gamma_kG)$.

The completion \widehat{G} is canonically filtered by the $\bar{\Gamma}_jG := \varprojlim (\Gamma_jG/\Gamma_kG)$. This filtration is its *closed lower central series*, defined as the closure of the lower central series. It is minimal amongst *closed* strongly central filtrations on \widehat{G} . An endomorphism of \widehat{G} is continuous if and only if it preserve this filtration.

Lemma 2.42. *Let G be a group. A continuous endomorphism φ of \widehat{G} is an automorphism if and only if the induced morphism $\varphi^{ab} \in \text{End}(G^{ab})$ is.*

Proof. Such an endomorphism is an automorphism if and only if the associated morphism between projective system is. These are the induced endomorphisms of the G/Γ_kG , which are nilpotent groups. This condition amounts to φ inducing an isomorphism on $\widehat{G}/\bar{\Gamma}_2 = G^{ab}$, by Lemma 2.40. \square

In fact, we can readily deduce the following explicit description of the group $\text{Aut}_{c^0}(\widehat{G})$ of continuous automorphisms of \widehat{G} :

Proposition 2.43. *The canonical map is an isomorphism:*

$$\text{Aut}_{c^0}(\widehat{G}) \cong \varprojlim (\text{Aut}(G/\Gamma_kG)).$$

Let $G = F_n$ be a free group of finite type. It is residually nilpotent, so it embeds into its completion \widehat{F}_n . A continuous endomorphism of \widehat{F}_n is uniquely determined by its (arbitrary) values on the topological generators x_i .

Let us denote by $F_{n,c}$ the free c -nilpotent group $F_n/\Gamma_{c+1}F_n$. Using Lemmas 2.40 and 2.42, we can show two surjectivity results for automorphisms of free nilpotent groups:

Proposition 2.44. *The canonical morphisms $\text{Aut}(F_{n,c}) \rightarrow \text{Aut}(F_{n,c-1})$ are surjective.*

Proof. Let (x_i) be a free basis of F_n . Let $\varphi \in \text{Aut}(F_n/\Gamma_c(F_n))$. Lift the elements $\varphi(\bar{x}_i)$ to elements t_i of $F_n/\Gamma_{c+1}(F_n)$. Then define the endomorphism $\tilde{\varphi}$ of $F_n/\Gamma_{c+1}(F_n)$ by $\bar{x}_i \mapsto t_i$. Since φ and $\tilde{\varphi}$ induce the same endomorphism of F_n^{ab} , Lemma 2.40 implies that $\tilde{\varphi}$ is an isomorphism. \square

Proposition 2.45 ([Bar13, th. 5.1]). *The Johnson morphism associated to the universal action on $\bar{\Gamma}_*(F_n)$ is an isomorphism:*

$$\tau : \mathcal{L}(\mathcal{A}_*(\bar{\Gamma}_*F_n)) \cong \text{Der}(LV).$$

Proof. Because of Lemma 1.28, we know that τ is injective. We need to show that it is surjective. Let us first remark that $\mathcal{L}(\bar{\Gamma}_*(F_n)) = \mathcal{L}(\Gamma_*(F_n)) \cong LV$. Let $\partial \in \text{Der}_k(LV)$. Lift each $\partial(X_i) \in \Gamma_k/\Gamma_{k+1} \cong \bar{\Gamma}_k/\bar{\Gamma}_{k+1}$ to an element $t_i \in \bar{\Gamma}_k$. We can define a continuous endomorphism of \widehat{F}_n by $\varphi : x_i \mapsto t_i x_i$. Then φ acts trivially on F_n^{ab} , so it is an isomorphism by Lemma 2.42. As a consequence, $\varphi \in \mathcal{A}_k(\bar{\Gamma}_*(F_n))$ satisfies $\tau(\varphi) = \partial$. This concludes the proof. \square

We can translate this last result into a statement about automorphisms of free nilpotent groups : because of Propositions 2.43 and 2.44, $\text{Aut}(F_{n,c})$ is a quotient of $\text{Aut}_{c^0}(\widehat{F}_n)$. Moreover, the kernel of the canonical surjection is $\mathcal{A}_c(\bar{\Gamma}_*F_n)$, by definition. Thus, this projection induces the c -truncations at the level of the associated graded objects :

Corollary 2.46. *The Johnson morphism associated to the universal action on $\Gamma_*(F_{n,c})$ is an isomorphism:*

$$\tau : \mathcal{L}(\mathcal{A}_*(F_{n,c})) \cong \text{Der}(L_{\leq c}V).$$

Consider the Johnson morphism $\tau' : \mathcal{L}(\Gamma_*(IA_{F_{n,c}})) \rightarrow \text{Der}(L_{\leq c}V)$. Using the identification of corollary 2.46, we see that τ' identifies with the Andreadakis morphism $i_* : \mathcal{L}(\Gamma_*(IA_{F_{n,c}})) \rightarrow \mathcal{L}(\mathcal{A}_*(F_{n,c}))$. Its image is the subalgebra generated in degree one inside $\text{Der}(L_{\leq c}V)$, as was the case in paragraph 2.5.2. This subalgebra is exactly the truncation $\mathfrak{J}_{<c}$, so is inside (and stably equal to) the kernel of the trace map. As a consequence, **the Andreadakis equality never holds for free nilpotent groups**. Moreover, in this context, our stable surjectivity result translates as :

Corollary 2.47. *The following sequence always is a complex, and is exact for $n \geq c+1$:*

$$\mathcal{L}(\Gamma_*(IA_{F_{n,c}})) \xrightarrow{i_*} \mathcal{L}(\mathcal{A}_*(F_{n,c})) \cong \text{Der}(L_{\leq c}V) \xrightarrow{\text{Tr}} C_{<c}V \longrightarrow 0.$$

Remark 2.48 (Non-tame automorphisms). The canonical morphism $p : \text{Aut}(F_n) \rightarrow \text{Aut}(F_{n,c})$ is in general not surjective: some basis of the free nilpotent group do not lift to basis of the free group *via* $F_n \rightarrow F_{n,c}$. Automorphisms of $F_{n,c}$ induced by automorphisms of F_n are called *tame*. This was the original motivation of [BLGM90] for considering the trace map. In this regard, our stable surjectivity result could be re-stated as follows : in the stable range, the trace is the only obstruction for an automorphism to be tame.

3 The case of positive characteristic

3.1 Dark's theorem

Let w be a word in a free group F_S . Let $w(\mathbf{r})$ be the word obtained from w by replacing each generator $s \in S$ by some power s^{r_s} . Dark's theorem [Dar68] describes how to decompose $w(\mathbf{r})$ as a product of commutators. This gives very useful universal formulas, that can then be evaluated in any group. The reader is referred to [Pas79, chap. IV, Th. 1.11] for a precise statement and a proof of the theorem. Here we recall two corollaries, obtained by taking $w = [x, y]$ and $w = xy$ in $F_{\{x,y\}}$.

The case $w = [x, y]$ is [Pas79, IV, cor. 1.16]:

Corollary 3.1. *There exists a unique map $\theta : (\mathbb{N}^*)^2 \rightarrow F_2 = \langle x, y \rangle$ verifying:*

$$\forall \alpha, \beta \in \mathbb{N}, [x^\alpha, y^\beta] = \prod_{r,s \geq 1} \theta(r, s) \binom{\alpha}{r} \binom{\beta}{s}.$$

Each $\theta(r, s)$ is a product of $\{x^{\pm 1}, y^{\pm 1}\}$ -commutators such that $x^{\pm 1}$ appears at least r times and $y^{\pm 1}$ at least s times in each factor.

The case $w = xy$ has been known for a long time (quoted in [Pas85, chap. 11, Th. 1.14], it already appears for instance in [Hal34]):

Corollary 3.2. *There exists a unique map $\theta : \mathbb{N} \rightarrow F_2 = \langle x, y \rangle$ such that:*

$$\forall \alpha \in \mathbb{N}, x^\alpha y^\alpha = \prod_{r \geq 0} \theta(r)^{\binom{\alpha}{r}},$$

Each $\theta(r)$ is a product of $\{x^{\pm 1}, y^{\pm 1}\}$ -commutators of length at least r .

Remark 3.3. In both cases, uniqueness of the map θ is obvious: it can be defined by induction.

3.2 p -Restricted strongly central series

Definition 3.4. Let p be a prime number. A strongly central series G_* is said to be p -restricted if:

$$\forall i, G_i^p \subseteq G_{ip}.$$

Let G_* be a p -restricted strongly central series. Using [Pas79, Th. III.1.7], we see that the morphism:

$$\mathcal{L}(G_*) \longrightarrow \text{gr}(\mathfrak{a}_*^{\mathbb{F}_p}(G_*))$$

induced by $g \mapsto g - 1$ (see Theorem 1.37 and Proposition 1.40) is injective. We can identify $\mathcal{L}(G_*)$ with its image, which is stable by the p -th power operation in the associative \mathcal{F}_p -algebra $\text{gr}(\mathfrak{a}_*^{\mathbb{F}_p}(G_*))$, since $(g - 1)^p = g^p - 1$. From this we deduce that $\mathcal{L}(G_*)$ is a p -restricted Lie algebra with p -th power operation induced by $g \mapsto g^p$ in G_1 . The reader is referred to the classical [Jac41] for a discussion of p -restricted Lie algebras. We also can deduce from [Pas79, Th. III.1.7] that any p -restricted strongly central series has to contain the dimension series $D_*^{\mathbb{F}_p}G$ defined in Example 1.39 (because $\mathfrak{a}_*^{\mathbb{F}_p}(G_*)$ contains $I_{\mathbb{F}_p}^*G$):

Proposition 3.5. *The filtration $D_*^{\mathbb{F}_p}G$ is the minimal p -restricted strongly central series, on any group G .*

The filtration $D^{\mathbb{F}_p}$, also denoted by $\Gamma_*^{[p]}$ also admits a description by induction, or the more explicit description:

$$\Gamma_k^{[p]}G = \prod_{ip^j \geq n} (\Gamma_i G)^{p^j}.$$

These can be found in [Laz54, Th. 5.6] or [Pas79, Th. IV.1.9]. We also refer to [CE16] for a nice discussion of filtrations defined by induction. This description has a nice consequence, similar to Proposition 1.9 (we abbreviate $\mathcal{L}(\Gamma_*^{[p]}G)$ to $\mathcal{L}^{[p]}(G)$):

Proposition 3.6. *The p -restricted Lie ring $\mathcal{L}^{[p]}(G)$ is generated in degree 1. Precisely, it is generated (as a p -restricted Lie ring) by $\mathcal{L}_1^{[p]}(G) = G^{ab} \otimes \mathbb{F}_p$.*

Example 3.7. If G is a free group, then $\mathcal{L}^{[p]}(G) = \mathcal{L}(D_*^{\mathbb{F}_p}G)$ is the p -restricted Lie algebra generated by the degree-one part inside $\text{gr}(\mathbb{F}_p G) \cong T_{\mathbb{F}_p}(G^{ab})$ so, using PBW over \mathbb{F}_p [Jac41, Th. 1] it is the free p -restricted Lie algebra over the \mathbb{F}_p -module $G^{ab} \otimes \mathbb{F}_p$ [Laz54, Th. 6.5].

Remark 3.8. The filtration $\Gamma_*^{[p]}G$ (already defined in [Zas39]) is to be distinguished from Stallings' filtration $\Gamma_*^{(p)}G$, defined in [Sta65]. The latter is the minimal p -torsion strongly central filtration on a group G , where a strongly central filtration is p -torsion when $G_i^p \subseteq G_{i+1}$ (for all i). See remark 3.14 for more on q -torsion strongly central series.

3.3 The p -restricted Andreadakis problem

Let us denote $\mathcal{A}_*(\Gamma_*^{[p]}G)$ by $\mathcal{A}_*^{[p]}(G)$. Remark that $\mathcal{A}_1^{[p]}(G)$ is the group $IA_G^{[p]}$ of automorphisms acting trivially on $\mathcal{L}_1(\Gamma_*^{[p]}G) = G^{ab} \otimes \mathbb{F}_p$, hence on all of $\mathcal{L}(\Gamma_*^{[p]}G)$ (because of Proposition 3.6). If we show that $\mathcal{A}_*^{[p]}(G)$ is p -restricted (and we will – see Proposition 3.10) then we get an inclusion:

$$\Gamma_*^{[p]}(IA_G^{[p]}) \subseteq \mathcal{A}_*^{[p]}(G).$$

We are thus led to consider a p -restricted version of the Andreadakis problem:

Problem 3 (Andreadakis – p -restricted version). *What is the difference between the p -restricted strongly central series $\mathcal{A}_*^{[p]}(G)$ and $\Gamma_*^{[p]}(IA_G^{[p]})$?*

Remark 3.9. The group $IA_G^{[p]}$ contains IA_G as a normal subgroup. Moreover, the quotient $IA_G^{[p]}/IA_G$ is a subgroup of $\text{Aut}(G)/IA_G$, hence of $GL(G^{ab})$. In fact, by definition of $IA_G^{[p]}$, it is contained in the congruence group:

$$GL(pG^{ab}) = \ker(GL(G^{ab}) \rightarrow GL(G^{ab} \otimes \mathbb{F}_p)).$$

When $G = F_n$ is a free group of finite type, then $\text{Aut}(G)/IA_G \cong GL_n(\mathbb{Z})$, and $IA_G^{[p]}/IA_G$ is exactly $GL_n(p\mathbb{Z})$.

Proposition 3.10 ([HM17], Prop. 8.5). *Let G_* be a p -restricted strongly central filtration, and K be a group acting on G_* . Then $\mathcal{A}_*(K, G_*)$ is a p -restricted strongly central filtration.*

Proof. Let $\kappa \in \mathcal{A}_j(K, G_*)$ and $g \in G_i$. Using Corollary 3.1, we get:

$$[\kappa^p, g] = \prod_{k=1}^p \theta(k, 1)^{\binom{p}{k}},$$

with $\theta(k, 1) \in G_{i+kj}$, for all k . If $k < p$, then $\theta(k, 1)^{\binom{p}{k}} \in G_{p(i+kj)} \subseteq G_{i+pj+1}$. As $\theta(p, 1)$ also is in G_{i+pj} , we have:

$$[\kappa^p, g] \in G_{i+pj}.$$

This is true for every $g \in G_i$, for all i . Hence $\kappa^p \in \mathcal{A}_{pj}(K, G_*)$, which completes the proof. \square

Proposition 3.10 can be refined:

Proposition 3.11. *Under the same hypothesis as Proposition 3.10, $\mathcal{A}_*(K, G_*) \rtimes G_*$ is a p -restricted strongly central series.*

Proof. Denote $\mathcal{A}_*(K, G_*) \rtimes G_*$ by \mathcal{K}_* . An element of $\mathcal{K}_j = \mathcal{A}_j \rtimes G_j$ is a product $\kappa \cdot g$, with $\kappa \in \mathcal{A}_j$ and $g \in G_j$. Using Corollary 3.2, we get:

$$\kappa^p g^p = \prod_{k=1}^p \theta(k)^{\binom{p}{k}} = (\kappa g)^p \cdot \theta(2)^{\binom{p}{2}} \cdots \theta(p-1)^p \cdot \theta(p), \quad (3.11.1)$$

with $\theta(1) = \kappa g$ and $\theta(k) \in \mathcal{K}_{kj}$ for any k , as \mathcal{K}_* is strongly central. We use this formula to show, by induction on $d \leq p$, the following result:

$$\forall j, \mathcal{K}_j^p \subseteq \mathcal{K}_{dj}.$$

This is true for $d = 1$, obviously. Let us assume that it holds for $d - 1$. Let $\kappa g \in \mathcal{K}_j$. For the sake of clarity, let us rewrite the formula (3.11.1):

$$(\kappa g)^p = \kappa^p g^p \cdot \theta(p)^{-1} \cdot \prod_{k=p-1}^2 \theta(k)^{-\binom{p}{k}}.$$

Using, respectively, that \mathcal{A}_* is p -restricted (Proposition 3.10), that G_* is (by definition) and that $\mathcal{K}_* = \mathcal{A}_* \times G_*$ is strongly central, we get:

$$\kappa^p, g^p, \theta(p) \in \mathcal{K}_{pj} \subseteq \mathcal{K}_{dj},$$

where the inclusion comes from the inequality $d \leq p$. If $2 \leq k < p$, then $\theta(k) \in \mathcal{K}_{kj}$. As p divides $\binom{p}{k}$, the induction hypothesis implies:

$$\theta(k)^{\binom{p}{k}} \in \mathcal{K}_{kj}^p \subseteq \mathcal{K}_{(d-1)kj} \subseteq \mathcal{K}_{dj},$$

because $(d-1)kj \geq dj$. Finally, we get what we were looking for:

$$(\kappa g)^p \in \mathcal{K}_{dj},$$

which completes the induction step, and the proof of the proposition. \square

Let \mathcal{SCF}_p be the full subcategory of \mathcal{SCF} given by p -restricted strongly central series. As a consequence of Propositions 3.10 and 3.11, we get:

Corollary 3.12. *The category \mathcal{SCF}_p is action-representative, the universal action on G_* being $\mathcal{A}_*(G_*) \circlearrowleft G_*$.*

This allows us to answer [HM17, rk. 8.6]. Indeed, the Lie functor restricts to a functor $\mathcal{L} : \mathcal{SCF}_p \rightarrow p\mathcal{L}ie$ with values in the category $p\mathcal{L}ie$ of p -restricted Lie algebras (over \mathbb{F}_p). Actions in $p\mathcal{L}ie$ are represented by *p -restricted derivations*, in the sense of Jacobson [Jac41]. As in Paragraph 1.4, an action $K_* \circlearrowleft G_*$ in \mathcal{SCF}_p induces, by exactness of the Lie functor, an action $\mathcal{L}(K_*) \circlearrowleft \mathcal{L}(G_*)$ in $p\mathcal{L}ie$, which is encoded by a morphism between p -restricted Lie algebras:

$$\tau : \mathcal{L}(K_*) \longrightarrow \text{Der}^{[p]}(\mathcal{L}(G_*)),$$

where $\text{Der}^{[p]} \subseteq \text{Der}$ is the p -restricted sub-algebra of p -restricted derivations, *i.e.* derivations ∂ satisfying:

$$\partial(a^p) = \text{ad}_a^{p-1}(\partial a).$$

Let us stress that for $\mathfrak{g} \in p\mathcal{L}ie$, the Lie algebra $\text{Der}(\mathfrak{g})$ is indeed a p -restricted sub-algebra of $\text{End}_{\mathbb{F}_p}(\mathfrak{g})$, but it does not act on \mathfrak{g} in $p\mathcal{L}ie$: the Lie algebra $\mathfrak{g} \rtimes \text{Der}(\mathfrak{g})$ bears no natural p -restricted structure.

Remark 3.13. Using Proposition 3.6 instead of 1.9, and replacing derivations by p -restricted ones in the proof, we can get an analogous of Lemma 1.30 for $\mathcal{A}_*^{[p]}(G)$:

$$\mathcal{A}_j^{[p]}(G) = \left\{ \sigma \in \text{Aut}(G) \mid [\sigma, G] \subseteq \Gamma_{j+1}^{[p]}(G) \right\}.$$

In other words $\mathcal{A}_*^{[p]}(G)$ is the subgroup of automorphisms acting trivially on $G/\Gamma_{j+1}^{[p]}(G)$. This is exactly the definition used by Cooper [Coo15, def 3.2].

Remark 3.14 (On the q -torsion case). The same statements are true when considering q -torsion strongly central filtrations (q does not have to be a prime number here), except that they are easier to show, because the condition $G_i^q \subseteq G_{i+1}$ is equivalent to the fact that $\mathcal{L}(G_*)$ is q -torsion. Precisely, if $\mathcal{L}(G_*)$ is q -torsion, then $\text{Der}(\mathcal{L}(G_*))$ is too, and the injectivity of the Johnson morphism $\mathcal{L}(\mathcal{A}_*(G_*)) \hookrightarrow \text{Der}(\mathcal{L}(G_*))$ implies that $\mathcal{L}(\mathcal{A}_*(G_*))$ also is. Hence $\mathcal{A}_*(G_*)$, and $\mathcal{A}_*(G_*) \times G_*$ are q -torsion, so that these give a universal action on G_* in the category of q -torsion strongly central series.

Moreover, $\mathcal{L}(G_*)$ also gets some kind of q -th power operation, induced by q -th powers in $G = G_1$. If $G_* = \Gamma_*^{(q)}G$ is Stallings' filtration on G , then these operations, together with the Lie algebra structure, generate $\mathcal{L}(G_*)$ from its degree one part, which allow us to get an analogue of Lemma 1.30: $\mathcal{A}_j(\Gamma_*^{(q)}G)$ is the subgroup of automorphisms acting trivially on $G/\Gamma_{j+1}^{(p)}(G)$. Using only this definition, Cooper managed to get the above results on the q -torsion case [Coo15]. However, his claim that the p -restricted case worked similarly [Coo15, Lem. 3.7] seems flawed, and we do not see how to get it without the technical work done above (Proposition 3.10).

The minimality of Stallings' filtration also gives an inclusion:

$$\Gamma_*^{(q)} \left(IA_G^{[q]} \right) \subseteq \mathcal{A}_* \left(\Gamma_*^{(q)} G \right),$$

and a corresponding Andreadakis problem. Nevertheless, our methods in studying the Andreadakis problems so far rely heavily on algebraic structures associated to the dimension subgroups $D_*^{\mathbb{Z}}(F_n) = \Gamma_*(F_n)$ and $D_*^{\mathbb{F}_p}(F_n) = \Gamma_*^{[p]}(F_n)$, so they are not suited to the study of this particular problem.

3.4 The stable p -restricted Andreadakis problem

3.4.1 Vanishing of the trace map

In the p -restricted context, Proposition 2.33 is replaced by:

Proposition 3.15. *Let $k \geq 2$, and let $J \in GL_m(I_{\mathbb{F}_p}^k F_n)$. Then:*

$$\text{Tr}(J - 1) \in [TV, TV]_k + (TV)^p \subset V^{\otimes k} \cong I^k / I^{k+1},$$

Where $V = F_n^{ab} \otimes_{\mathbb{F}_p} \cong \mathbb{F}_p^n$.

The proof is exactly the same as the proof of proposition 2.33, over \mathbb{F}_p instead of \mathbb{Z} , Proposition 2.34 being replaced by:

Proposition 3.16. *Let $f(X_1, \dots, X_n) \in V^{\otimes k}$. Let $\mathcal{C} \subset M_k(\mathbb{F}_p)$ be the sub- \mathbb{Z} -module generated by the $e_{i, i+1}$. Suppose:*

$$\forall C_i \in \mathcal{C}, \text{Tr}(f(C_1, \dots, C_n)) = 0.$$

Then $f \in [TV, TV]_k + (TV)^p$.

Proof. We say that two elements u and u' of $V^{\otimes k}$ are *cyclically equivalent*, and we write $u \sim u'$ if they are conjugate under the action of \mathbb{Z}/p . We want to show that f is cyclically equivalent to a p -th power. Let us decompose f , up to cyclical equivalence, as a sum of pairwise non-cyclically equivalent monomials: $f \sim \sum \mu_g g$, where each g is of the form $g = X_{i_1} \cdots X_{i_k}$. If g is such that the i_α are pairwise distinct, evaluate each X_{i_α} as $C'_{i_\alpha} = e_{\alpha, \alpha+1}$, and all X_i not appearing in g as $C_i = 0$. Then $\mu_g =$

$\text{Tr}(f(C_1, \dots, C_n)) = 0$. As a consequence, no such g can appear in our decomposition of f .

Let λ be the algebra morphism from TV to $T(V \otimes V)$ sending X_i to $\sum_j X_{ij}$ (where $X_{ij} = X_i \otimes X_j$). Take a monomial $g = X_{i_1} \cdots X_{i_k}$ as above. Its image is $\lambda(g) = \sum_{\mathbf{j}} X_{i_1 j_1} \cdots X_{i_k j_k}$, the sum being taken over every $\mathbf{j} = (j_1, \dots, j_k) \in \{1, \dots, k\}^k$. Let r be the number of monomials cyclically equivalent to $h = X_{i_1 1} \cdots X_{i_k k}$ in this sum. Then r is exactly the number of elements of \mathbb{Z}/k stabilizing g . It is a multiple of p if and only if g is a p -th power. If we decompose $\lambda(f)$ up to cyclic equivalence, as we did earlier for f , the only occurrences of h must come from $\lambda(g)$, hence the coefficient of h must be $r\mu_g$. Note that $\lambda(f)$ satisfies the same hypothesis as f , because \mathcal{C} is stable under addition. Since the X_{i_α} are pairwise distinct, we can apply the above argument and find that $r\mu_g = 0$. Thus, $\mu_g = 0$ or g is a p -th power. Whence the result. \square

Remark 3.17. Any bracket and any p -th power satisfies the condition of Proposition 3.16. For brackets, it follows from the fact that the trace of a bracket is itself a sum of brackets. For p -th power, remark that if $M = \sum m_i \mathbf{e}_{i, i+1} \in M_n(R)$, where R is an associative ring of characteristic p , then $M^p = (\prod m_i) \cdot 1_p$, and $\text{Tr}(1_p) = p \cdot 1 = 0$ in \mathbb{k} . If $k = pl$ and $f = (X_{i_1} \cdots X_{i_l})^p$, apply this to $M = C_{i_1} \cdots C_{i_l}$ (where the C_{i_α} are in \mathcal{C}), seen as a $p \times p$ -matrix with coefficients in $M_l(\mathbb{k})$.

Because of Proposition 3.15, in characteristic p , we will consider the trace map as taking values in $C_*^{[p]}V = TV/([TV, TV] + (TV)^p)$, which is the quotient of the cyclic power C_*V by p -th powers. The conclusion of Proposition 3.15 then becomes: $\text{Tr}(J - 1) = 0 \in C_*^{[p]}V$.

3.4.2 Linear algebra

Consider the Johnson morphism

$$(\tau^{[p]})' : \mathcal{L}^{[p]}(IA_n^{[p]}) \longrightarrow \text{Der}_*^{[p]}(\mathcal{L}^{[p]}(F_n)) \cong V^* \otimes \mathfrak{L}^{[p]}V,$$

the last isomorphism being obtained as in Example 1.33, using Example 3.7 instead of Example 1.8, and replacing derivations by p -restricted ones. When $p \neq 2$, the morphism $(\tau_1^{[p]})'$ is surjective. Indeed, the free \mathbb{F}_p -Lie algebra $\mathfrak{L}V$ is a sub-algebra of the free p -restricted algebra $\mathfrak{L}^{[p]}V$, and this inclusion is an isomorphism in degrees prime to p , in particular in degree 2; thus we can lift the generators of $V^* \otimes \Lambda^2 V$ by the generators of IA_n used in the proof of Proposition 1.34. Moreover, $\mathcal{L}^{[p]}(IA_n)$ is generated in degree 1 as a p -restricted Lie algebra (cf. 3.6). As a consequence, the image of τ' is exactly the p -restricted Lie algebra generated in degree 1 inside $\text{Der}_*^{[p]}(\mathfrak{L}^{[p]}V)$.

The reader can easily check that the obvious p -restricted version of Proposition 2.32 does hold: the trace map obtained from free differential calculus can be seen as the composite of the Johnson morphism $\tau : \mathcal{L}_k(\mathcal{A}_*^{[p]}(F_n)) \rightarrow \text{Der}_k^{[p]}(\mathfrak{L}^{[p]}V) \cong V^* \otimes \mathfrak{L}_{k+1}^{[p]}V$ with:

$$\text{Tr}_M : V^* \otimes \mathfrak{L}_{k+1}^{[p]}V \xrightarrow{\iota} V^* \otimes V^{\otimes k+1} \xrightarrow{\Phi} V^{\otimes k} \xrightarrow{\pi} C_k^{[p]}V,$$

where ι and π again denote the canonical maps.

Notation 3.18. Let $\mathfrak{J}^{[p]}$ denote the image of $(\tau^{[p]})'$, the p -restricted Lie algebra generated in degree 1 inside $\text{Der}(\mathfrak{L}V)$.

The following proposition can be seen as a direct consequence of Proposition 3.15.

Proposition 3.19. *For every $k \geq 2$, $\text{Tr}_M(\mathfrak{J}_k^{[p]}) = \{0\}$.*

Consider be the subspace of p -restricted derivations stabilizing the free \mathbb{F}_p -Lie algebra $\mathcal{L}V \subset \mathcal{L}^{[p]}V$. It is a p -restricted Lie sub-algebra of $\text{Der}^{[p]}(\mathcal{L}^{[p]}V)$. Since each derivation of $\mathcal{L}V$ extends to a unique p -restricted derivation of $\mathcal{L}^{[p]}V$, this sub-algebra is isomorphic to $\text{Der}(\mathcal{L}V)$. Under the identification with the graded module $V^* \otimes \mathcal{L}^{[p]}V$, it corresponds exactly to $V^* \otimes \mathcal{L}V$. As a consequence, if $p \neq 2$, the degree one part is the same. Hence:

$$\mathfrak{J}^{[p]} \subseteq \text{Der}_{\mathcal{L}V}^{[p]}(\mathcal{L}^{[p]}V).$$

This implies that there is **no stable surjectivity here**: we can easily produce examples of automorphisms whose associated derivation does not preserve $\mathcal{L}V$. For instance, take any world in $\Gamma_k(F_n)$ not containing any occurrence of x_1 . Then the automorphism φ defined by $x_1 \mapsto w^p x_1$ and $x_i \mapsto x_i$ when $i \neq 1$ is obviously in $\mathcal{A}_{pk-1}^{[p]}$, but $\tau(\varphi) = X_1^* \otimes (\overline{w-1})^p$ sends X_1 outside of $\mathcal{L}V$.

3.4.3 Stable cokernel of i_*

We close the present paper with a quantification of the lack of stable surjectivity in the p -restricted case.

Let $k \geq 2$. Like in paragraph 2.5.3, we get a commutative diagram with exact rows:

$$\begin{array}{ccccc} \tilde{\mathfrak{J}}_k & \hookrightarrow & V^* \otimes \mathfrak{L}_{k+1}V & \twoheadrightarrow & X_k \\ \downarrow \phi & & \downarrow \Phi & & \downarrow \bar{\Phi} \\ [TV, TV]_k & \hookrightarrow & V^{\otimes k} & \xrightarrow{\pi} & C_k V. \end{array}$$

Here, V denotes $F_n^{ab} \otimes \mathbb{F}_p$, and $\tilde{\mathfrak{J}}_* = \mathfrak{J}_* \otimes \mathbb{F}_p$ is the sub-Lie algebra generated in degree one inside $\text{Der}(\mathcal{L}V)$. The space X_k is just the quotient of $V^* \otimes \mathfrak{L}_{k+1}V$ by $\tilde{\mathfrak{J}}_k$.

We will be interested in a slightly different diagram, though:

$$\begin{array}{ccccc} \mathfrak{J}_k^{[p]} & \hookrightarrow & V^* \otimes \mathfrak{L}_{k+1}V & \twoheadrightarrow & X'_k \\ \downarrow \phi & & \downarrow \Phi & & \downarrow \bar{\Phi} \\ [TV, TV]_k + (TV)^p & \hookrightarrow & V^{\otimes k} & \xrightarrow{\pi} & C_k^{[p]}V. \end{array}$$

We can still apply the calculations from [Sat12] to show that if $n \geq k+2$, then Φ is surjective and $\ker \Phi \subseteq \tilde{\mathfrak{J}}_k \subseteq \mathfrak{J}_k^{[p]}$. Only, now ϕ could have a cokernel. From [Sat12], we only get that brackets are in its image, so this cokernel can only come from p -th powers. In particular, it is concentrated in degrees divisible by p . We denote it by K . The same application of the snake lemma as in the proof of Proposition 2.37 gives that K is also the kernel of $\bar{\Phi}$, and that $\bar{\Phi}$ is surjective.

Now consider the diagram:

$$\begin{array}{ccccc} \mathfrak{J}_k^{[p]} & \hookrightarrow & V^* \otimes \mathfrak{L}_{k+1}V & \twoheadrightarrow & X'_k \\ \downarrow \iota & & \downarrow & & \downarrow \bar{\Phi} \\ \ker(\text{Tr}_M)_k & \hookrightarrow & V^* \otimes \mathfrak{L}_{k+1}^{[p]}V & \xrightarrow{\text{Tr}_M} & C_k^{[p]}V. \end{array}$$

Denote by L the cokernel of the middle inclusion, then $L = V^* \otimes (\mathfrak{L}_{k+1}^{[p]} V / \mathfrak{L}_{k+1} V)$ is concentrated in degrees $k = pl - 1$ (with $l \geq 1$). The snake lemma gives a short exact sequence:

$$0 \longrightarrow K \longrightarrow \text{coker}(\iota) \longrightarrow L \longrightarrow 0.$$

Since the trace map $\text{Tr}_M \circ \tau$ vanishes, we have a commutative diagram:

$$\begin{array}{ccc} \mathcal{L}(IA_n^{[p]}) & \xrightarrow{i_*} & \mathcal{L}(\mathcal{A}_*^{[p]}) \\ & \searrow \tau' & \downarrow \tau \\ & & \ker(\text{Tr}_M). \end{array}$$

This implies that the cokernel of i_* injects into the cokernel of ι . Thus, we have proved:

Proposition 3.20. *Fix n an integer, and consider only degrees $k \leq n-2$. The cokernel of the canonical morphism*

$$i_* : \mathcal{L}(IA_n^{[p]}) \rightarrow \mathcal{L}(\mathcal{A}_*^{[p]}(F_n))$$

is concentrated in degrees $k = pl - 1$ and $k = pl$ (for $l \geq 1$). If $k = pl - 1$, then $\text{coker}((i_)_k)$ injects into $V^* \otimes (\mathfrak{L}_{k+1}^{[p]} V / \mathfrak{L}_{k+1} V)$. If $k = pl$, it is a sub-quotient of $V^{\otimes l}$.*

Remark 3.21. The tensor power $V^{\otimes l}$ appearing in the proposition is in fact the *Frobenius twist* of $V^{\otimes l}$. This has no consequence here, as the Frobenius map is trivial on \mathbb{F}_p , but it should be kept in mind for any functorial study of this situation.

References

- [And65] Stylianos Andreadakis. On the automorphisms of free groups and free nilpotent groups. *Proc. London Math. Soc. (3)*, 15:239–268, 1965. [2](#)
- [Bar13] Laurent Bartholdi. Automorphisms of free groups. I. *New York J. Math.*, 19:395–421, 2013. [2](#), [25](#), [26](#), [28](#), [32](#)
- [Bar16] Laurent Bartholdi. Automorphisms of free groups. I—erratum [[MR3084710](#)]. *New York J. Math.*, 22:1135–1137, 2016. [2](#)
- [BB04] Francis Borceux and Dominique Bourn. *Mal’cev, protomodular, homological and semi-abelian categories*, volume 566 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 2004. [7](#), [11](#)
- [BB07] Francis Borceux and Dominique Bourn. Split extension classifier and centrality. In *Categories in algebra, geometry and mathematical physics*, volume 431 of *Contemp. Math.*, pages 85–104. Amer. Math. Soc., Providence, RI, 2007. [2](#), [8](#)
- [BBM07] Mladen Bestvina, Kai-Uwe Bux, and Dan Margalit. Dimension of the Torelli group for $\text{Out}(F_n)$. *Invent. Math.*, 170(1):1–32, 2007. [1](#), [13](#)
- [BC05] F. Borceux and Maria Manuel Clementino. Topological semi-abelian algebras. *Adv. Math.*, 190(2):425–453, 2005. [7](#)

- [BEH08] Laurent Bartholdi, Bettina Eick, and René Hartung. A nilpotent quotient algorithm for certain infinitely presented groups and its applications. *Internat. J. Algebra Comput.*, 18(8):1321–1344, 2008. [2](#)
- [BJK05] Francis Borceux, George Janelidze, and Gregory M. Kelly. On the representability of actions in a semi-abelian category. *Theory Appl. Categ.*, 14:No. 11, 244–286, 2005. [2](#), [8](#)
- [BLGM90] Roger M. Bryant, Frank Levin, Chander K. Gupta, and Horace Y. Mochizuki. Nontame automorphisms of free nilpotent groups. *Comm. Algebra*, 18(11):3619–3631, 1990. [3](#), [28](#), [29](#), [33](#)
- [BMS67] Hyman Bass, John Milnor, and Jean-Pierre Serre. Solution of the congruence subgroup problem for SL_n ($n \geq 3$) and Sp_{2n} ($n \geq 2$). *Inst. Hautes Études Sci. Publ. Math.*, 33:59–137, 1967. [16](#)
- [Bou08] Dominique Bourn. Commutator theory, action groupoids, and an intrinsic Schreier-Mac Lane extension theorem. *Adv. Math.*, 217(6):2700–2735, 2008. [2](#), [8](#)
- [CE16] Michael Chapman and Ido Efrat. Filtrations of free groups arising from the lower central series. *J. Group Theory*, 19(3):405–433, 2016. [34](#)
- [CEFN14] Thomas Church, Jordan S. Ellenberg, Benson Farb, and Rohit Nagpal. FI-modules over Noetherian rings. *Geom. Topol.*, 18(5):2951–2984, 2014. [31](#)
- [Coo15] James Cooper. Two mod- p Johnson filtrations. *J. Topol. Anal.*, 7(2):309–343, 2015. [36](#), [37](#)
- [CP17] Thomas Church and Andrew Putamn. Generating the johnson filtration ii: finite generation. arXiv:1704.01529, 2017. [1](#)
- [CV86] Marc Culler and Karen Vogtmann. Moduli of graphs and automorphisms of free groups. *Invent. Math.*, 84(1):91–119, 1986. [1](#)
- [Dar68] Rex S. Dark. *On nilpotent products of groups of prime order*. PhD thesis, University of Cambridge, 1968. [33](#)
- [Dja16a] Aurélien Djament. Décomposition de hodge pour l’homologie stable des groupes d’automorphismes des groupes libres. arXiv:1510.03546, 2016. [1](#)
- [Dja16b] Aurélien Djament. Des propriétés de finitude des foncteurs polynomiaux. *Fund. Math.*, 233(3):197–256, 2016. [31](#)
- [DP12] Matthew Day and Andrew Putman. A Birman exact sequence for $\text{Aut}(F_n)$. *Adv. Math.*, 231(1):243–275, 2012. [1](#)
- [DP16] Matthew B. Day and Andrew Putamn. On the second homology group of the Torelli subgroup of $\text{aut}(f_n)$. arXiv:1408.6242v3, 2016. [1](#), [2](#)
- [DV15] Aurélien Djament and Christine Vespa. Sur l’homologie des groupes d’automorphismes des groupes libres à coefficients polynomiaux. *Comment. Math. Helv.*, 90(1):33–58, 2015. [1](#)

- [FM12a] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012. [1](#)
- [FM12b] Stefano Francaviglia and Armando Martino. The isometry group of outer space. *Adv. Math.*, 231(3-4):1940–1973, 2012. [1](#)
- [Fox53] Ralph H. Fox. Free differential calculus. I. Derivation in the free group ring. *Ann. of Math. (2)*, 57:547–560, 1953. [19](#)
- [Fra33] Hermann Frasch. Die Erzeugenden der Hauptkongruenzgruppen für Primzahlstufen. *Math. Ann.*, 108(1):229–252, 1933. [17](#)
- [Gal11] Søren Galatius. Stable homology of automorphism groups of free groups. *Ann. of Math. (2)*, 173(2):705–768, 2011. [1](#)
- [Gro52] Emil Grosswald. On the parabolic generators of the principal congruence subgroups of the modular group. *Amer. J. Math.*, 74:435–443, 1952. [17](#)
- [Hal34] Philip Hall. A contribution to the theory of groups of prime-power order. *Proc. London Math. Soc. (2)*, 36:29–95, 1934. [33](#)
- [HL11] Manfred Hartl and Bruno Loiseau. A characterization of finite cocomplete homological and of semi-abelian categories. *Cah. Topol. Géom. Différ. Catég.*, 52(1):77–80, 2011. [7](#)
- [HM17] Kazuo Habiro and Gwenaél Massuyeau. Generalized johnson homomorphisms for extended n -series. *arXiv:1707.07428*, 2017. [2](#), [4](#), [12](#), [35](#), [36](#)
- [Jac41] Nathan Jacobson. Restricted Lie algebras of characteristic p . *Trans. Amer. Math. Soc.*, 50:15–25, 1941. [34](#), [36](#)
- [Joh83] Dennis Johnson. A survey of the Torelli group. In *Low-dimensional topology (San Francisco, Calif., 1981)*, volume 20 of *Contemp. Math.*, pages 165–179. Amer. Math. Soc., Providence, RI, 1983. [13](#)
- [JW04] Craig A. Jensen and Nathalie Wahl. Automorphisms of free groups with boundaries. *Algebr. Geom. Topol.*, 4:543–569, 2004. [1](#)
- [Kal50a] Léo Kaloujnine. Sur quelques propriétés des groupes d’automorphismes d’un groupe abstrait. *C. R. Acad. Sci. Paris*, 230:2067–2069, 1950. [8](#)
- [Kal50b] Léo Kaloujnine. Sur quelques propriétés des groupes d’automorphismes d’un groupe abstrait. (Généralisation d’un théorème de M. Ph. Hall). *C. R. Acad. Sci. Paris*, 231:400–402, 1950. [8](#)
- [Kaw06] Nariya Kawazumi. Cohomological aspects of magnus expansions. *arXiv:math/0505497*, 2006. [13](#)
- [KM97] Sava Krstić and James McCool. The non-finite presentability of $IA(F_3)$ and $GL_2(\mathbf{Z}[t, t^{-1}])$. *Invent. Math.*, 129(3):595–606, 1997. [1](#)
- [Laz54] Michel Lazard. Sur les groupes nilpotents et les anneaux de Lie. *Ann. Sci. Ecole Norm. Sup. (3)*, 71:101–190, 1954. [3](#), [6](#), [8](#), [14](#), [34](#)

- [Lop14] Jonathan Lopez. Lie algebras and cohomology of congruence subgroups for $SL_n(R)$. *J. Pure Appl. Algebra*, 218(2):256–268, 2014. [3](#), [16](#)
- [LS76] Ronnie Lee and Robert H. Szczarba. On the homology and cohomology of congruence subgroups. *Invent. Math.*, 33(1):15–53, 1976. [3](#), [17](#)
- [ML98] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998. [7](#)
- [Mor93] Shigeyuki Morita. Abelian quotients of subgroups of the mapping class group of surfaces. *Duke Math. J.*, 70(3):699–726, 1993. [3](#), [25](#)
- [MP09] Roman Mikhailov and Inder Bir Singh Passi. *Lower central and dimension series of groups*, volume 1952 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009. [14](#)
- [MS17] Gwenael Massuyeau and Takuya Sakasai. Morita’s trace maps on the group of homology cobordisms. *arXiv:1606.08244*, 2017. [2](#), [28](#)
- [Nie24] Jakob Nielsen. Die Isomorphismengruppe der freien Gruppen. *Math. Ann.*, 91(3-4):169–209, 1924. [1](#), [13](#)
- [Pas79] Inder Bir S. Passi. *Group rings and their augmentation ideals*, volume 715 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979. [14](#), [33](#), [34](#)
- [Pas85] Donald S. Passman. *The algebraic structure of group rings*. Robert E. Krieger Publishing Co., Inc., Melbourne, FL, 1985. Reprint of the 1977 original. [33](#)
- [Rip72] Eliyahu Rips. On the fourth integer dimension subgroup. *Israel J. Math.*, 12:342–346, 1972. [14](#)
- [Rot95] Joseph J. Rotman. *An introduction to the theory of groups*, volume 148 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, fourth edition, 1995. [13](#)
- [RWW17] Oscar Randal-Williams and Nathalie Wahl. Homological stability for automorphism groups. *Adv. Math.*, 318:534–626, 2017. [1](#)
- [Sat12] Takao Satoh. On the lower central series of the IA-automorphism group of a free group. *J. Pure Appl. Algebra*, 216(3):709–717, 2012. [2](#), [3](#), [25](#), [30](#), [39](#)
- [Sta65] John Stallings. Homology and central series of groups. *J. Algebra*, 2:170–181, 1965. [34](#)
- [Zas39] Hans Zassenhaus. Ein Verfahren, jeder endlichen p -Gruppe einen Lie-Ring mit der Charakteristik p zuzuordnen. *Abh. Math. Sem. Univ. Hamburg*, 13(1):200–207, 1939. [34](#)